Fractal Bindu (S^3) of Reality: Local Flatness, Quantum Consistency, and the ΛCDM Limit

Andrei Preece and Boris Batenin

Berlin, 2025

Abstract

We propose a cosmological model where the Universe is a 3-sphere (S^3) with a large radius R. We demonstrate that the observed flatness of the Universe within the observable horizon ($r_{\text{obs}} \approx 13.8 \times 10^9$ light-years) arises as a local approximation due to $r_{\text{obs}} \ll R$, where the metric of S^3 reduces to that of flat Euclidean space. The ΛCDM model emerges as a special case in the limit $R \to \infty$, corresponding to zero global curvature. We show that quantum mechanics remains consistent on S^3, as it operates in the topological structure defined by the wave function, not the geometric projection. The compact topology of S^3 naturally resolves issues like infrared divergences in quantum field theory and provides a framework for quantum cosmology. Observational implications, such as the cosmic microwave background (CMB) power spectrum and large-scale structure, are discussed, showing compatibility with current data while offering testable predictions.

1 Introduction

The standard ΛCDM model describes the Universe as a flat, expanding space with zero curvature ($k = 0$), driven by dark energy, dark matter, and baryonic matter. However, this model assumes a globally flat topology ($\mathbb{R}^3$), which is a local approximation that may not reflect the true global structure of the Universe. Observational data, such as the Planck 2018 results, indicate that the curvature parameter Ω_k is very close to zero ($\Omega_k = 0.001 \pm 0.002$), but allow for a small positive curvature, suggesting a possible closed topology like a 3-sphere (S^3).

In this paper, we propose that the Universe is a 3-sphere S^3 with a large radius R. We show that:

1. The observed flatness within the observable horizon ($r_{\text{obs}} \approx 13.8 \times 10^9$ light-years) is a natural consequence of the local geometry of S^3 when $r_{\text{obs}} \ll R$.

2. The ΛCDM model is a limiting case of our model as $R \to \infty$.

3. Quantum mechanics is fully consistent on S^3, and the compact topology provides a natural framework for quantum cosmology, potentially resolving issues like infrared divergences in quantum field theory.

2 Problem Statement

Consider the Universe as a 3-sphere S^3 with radius R. An observer at point O has a light horizon limited by the distance $r_{\text{obs}} \approx 13.8 \times 10^9$ light-years, corresponding to the age of the Universe in the standard model. Due to the expansion of the Universe, the comoving distance to the particle horizon is larger, approximately 46.5 billion light-years (or a diameter of 93 billion light-years for the observable Universe). We aim to prove that:

1. The geometry within the observable horizon appears flat, even if R is finite.
2. The observed Universe is a local projection on S^3, with no "openness" in the topological sense.
3. The ΛCDM model is a special case of our S^3 model in the limit of zero global curvature.
4. Quantum mechanics remains consistent on S^3 and is naturally supported by its compact topology.

3 Local Geometry of S^3

The metric of a 3-sphere S^3 with radius R in the Friedmann-Lemaître-Robertson-Walker (FLRW) framework, assuming a static topology for simplicity, is given by:

$$ds^2 = R^2 d\chi^2 + R^2 \sin^2 \chi \, d\Omega_2^2, \tag{1}$$

where $\chi \in [0, \pi]$ is the radial angular coordinate, and $d\Omega_2^2$ is the angular element on the 2-sphere S^2.

The physical distance r is related to χ as $r = R\chi$. For small distances $r \ll R$, we have $\chi \ll 1$. Using the Taylor expansion for small χ:

$$\sin \chi \approx \chi, \tag{2}$$

the metric simplifies to:

$$ds^2 \approx R^2 d\chi^2 + R^2 \chi^2 d\Omega_2^2. \tag{3}$$

Substituting $\chi = r/R$, we get $d\chi = dr/R$, and the metric becomes:

$$ds^2 = dr^2 + r^2 d\Omega_2^2, \tag{4}$$

which is the metric of flat Euclidean space $\mathbb{R}^3$ in spherical coordinates. Thus, locally, $S^3 \approx \mathbb{R}^3$.

3.1 Numerical Estimate

Let us estimate χ for the observable Universe. The physical distance $r_{\text{obs}} \approx 13.8 \times 10^9$ light-years is approximately 1.3×10^{26} meters (using 1 light-year $\approx 9.46 \times 10^{15}$ m). If we assume a hypothetical radius $R \sim 10^{27}$ meters (a reasonable value for S^3 to be consistent with the near-flatness observed), then:

$$\chi = \frac{r_{\text{obs}}}{R} \approx \frac{1.3 \times 10^{26}}{10^{27}} = 0.13. \tag{5}$$

For $\chi = 0.13$, we compute:

$$\sin\chi \approx \sin 0.13 \approx 0.1296, \quad \chi \approx 0.13, \tag{6}$$

so the approximation $\sin\chi \approx \chi$ holds with an error of less than 0.3%. This confirms that the observable Universe is well within the regime where S^3 appears flat.

4 ΛCDM as a Limiting Case

The ΛCDM model assumes a flat FLRW metric with curvature parameter $k = 0$:

$$ds^2 = -dt^2 + a(t)^2\left(dr^2 + r^2 d\Omega_2^2\right), \tag{7}$$

where $a(t)$ is the scale factor. In our S^3 model, the curvature parameter is related to the radius R as $k = 1/R^2$. As $R \to \infty$, the curvature $k \to 0$, and the metric of S^3 reduces to that of $\mathbb{R}^3$:

$$\lim_{R\to\infty} S^3 = \mathbb{R}^3. \tag{8}$$

Thus, ΛCDM is a local approximation of our S^3 model in the limit of zero global curvature.

4.1 Observational Consistency

The Planck 2018 data constrain the curvature parameter to $\Omega_k = 0.001 \pm 0.002$, consistent with a flat Universe but allowing for a small positive curvature. For $R \sim 10^{27}$ meters, the curvature parameter $\Omega_k \sim -1/(H_0 R)^2$, where $H_0 \approx 70\,\mathrm{km/s/Mpc} \approx 2.27 \times 10^{-18}\,\mathrm{s}^{-1}$. This gives:

$$\Omega_k \sim -\frac{1}{(2.27 \times 10^{-18} \cdot 10^{27})^2} \sim -10^{-18}, \tag{9}$$

which is far below the observational limit, confirming that S^3 with such a large R is indistinguishable from flatness within current precision.

5 Quantum Mechanics on S^3

Quantum mechanics operates in a Hilbert space, where states evolve according to the Schrödinger equation $i\hbar\frac{d}{dt}\Psi = \hat{H}\Psi$. The wave function $\Psi(t) = A(t)e^{i\phi(t)}$ evolves under a Hamiltonian $\hat{H}$, independent of the geometric interpretation of space-time.

In our model, the Universe is described by a global wave function $\Psi(t)$ on S^3. The compact topology of S^3 imposes periodic boundary conditions on Ψ, as the coordinate χ is defined on $[0, \pi]$. This leads to a discrete spectrum of energy eigenstates, unlike the continuous spectrum in $\mathbb{R}^3$.

5.1 Topological Effects

The compact nature of S^3 has several implications for quantum mechanics:

- **Discrete Spectrum**: The periodicity in χ implies that wave functions must satisfy $\Psi(\chi = 0) = \Psi(\chi = \pi)$, leading to quantization of momenta and energies. This is analogous to a particle in a box, but in a spherical geometry.

- **Infrared Regularization**: In quantum field theory, the infinite volume of $\mathbb{R}^3$ leads to infrared divergences. On S^3, the finite volume ($V = 2\pi^2 R^3$) naturally regularizes these divergences, providing a more consistent framework for quantum cosmology.
- **Quantum Cosmology**: The S^3 topology aligns with models like the Hartle-Hawking "no-boundary" proposal, where the wave function of the Universe is defined on a compact geometry, avoiding singularities.

5.2 Consistency with Local Flatness

Locally, the geometry appears as $\mathbb{R}^3$, so quantum mechanical calculations (e.g., for particle interactions) are unaffected. However, globally, the S^3 topology influences the wave function's boundary conditions, which can manifest in cosmological observables like the CMB power spectrum.

5.3 Support for Quantum Mechanics

The S^3 topology not only preserves quantum mechanics but also provides a natural framework for it. For example, the discrete spectrum on S^3 can explain the quantization of primordial fluctuations in the early Universe, which seed the CMB anisotropies. This aligns with quantum cosmology, where the wave function of the Universe describes the evolution of geometry itself.

6 Observational Predictions

Our S^3 model makes testable predictions that can be compared with ΛCDM:

- **CMB Power Spectrum**: The compact topology of S^3 introduces a cutoff in the CMB power spectrum at large angular scales (low ℓ). For $R \sim 10^{27}$ meters, the fundamental mode corresponds to an angular scale $\theta \sim \lambda/R$, where $\lambda \sim R$ is the wavelength of the largest mode. This predicts a suppression of power at $\ell \lesssim 10$, which can be tested with future CMB experiments.
- **Large-Scale Structure**: The periodic boundary conditions on S^3 lead to a discrete Fourier spectrum for density perturbations, potentially observable as a "quantization" in the galaxy power spectrum at very large scales.
- **Curvature Constraints**: Future measurements of Ω_k with higher precision could detect a small positive curvature, supporting the S^3 model.

7 Conclusion

We have shown that the Universe can be modeled as a 3-sphere S^3 with a large radius R, where the observed flatness within the observable horizon ($r_{\text{obs}} \approx 13.8 \times 10^9$ light-years) arises because $r_{\text{obs}} \ll R$, making $\sin\chi \approx \chi$ a valid approximation. Quantum mechanics remains consistent on S^3, as it operates in the topological structure defined by the wave function, not the geometric projection. The compact topology of S^3 provides a natural framework for quantum cosmology, resolving issues like infrared divergences and supporting the quantization of primordial fluctuations.

7.1 ΛCDM as a Special Case

The ΛCDM model is a special case of our S^3 model in the limit $R \to \infty$, corresponding to zero global curvature. This implies that ΛCDM is a local approximation that neglects the global topology of the Universe. While ΛCDM successfully describes the Universe on scales up to the observable horizon, it cannot address questions about the global structure, such as the finite volume or periodic boundary conditions that arise in S^3. Our model generalizes ΛCDM, offering a more comprehensive framework that can be tested through large-scale cosmological observations.

7.2 Future Directions

The S^3 model is compatible with current observations (e.g., Planck 2018) and makes testable predictions, such as a suppression of CMB power at low ℓ and a discrete spectrum in the large-scale structure. Future observations with higher precision could distinguish between the S^3 topology and the infinite $\mathbb{R}^3$ assumed by ΛCDM, offering a deeper understanding of the Universe's global structure.

Namaste!

References

[1] Planck Collaboration, "Planck 2018 results. VI. Cosmological parameters," *Astronomy & Astrophysics*, vol. 641, A6, 2020.

[2] J. B. Hartle and S. W. Hawking, "Wave function of the Universe," *Physical Review D*, vol. 28, no. 12, pp. 2960–2975, 1983.

[3] B. S. DeWitt, "Quantum Theory of Gravity. I. The Canonical Theory," *Physical Review*, vol. 160, no. 5, pp. 1113–1148, 1967.

[4] A. D. Linde, "Scalar Field Fluctuations in the Expanding Universe and the New Inflationary Universe Scenario," *Physics Letters B*, vol. 116, no. 5, pp. 335–339, 1982.