Finite Dilation Algebra of Round Euclidean B^4/S^3 Valuation Profiles: Exact Invariants and Executed Homogeneous Null Tests

Andrei Preece [1, *] and Boris Batenin [1, †]

[1] *Independent Researcher, Berlin, Germany*

(Dated: August 10, 2026)

We prove a finite homogeneous classification for round Euclidean valuation profiles associated with a closed FLRW spatial section S^3_A. The universal work map from an auxiliary filling energy to homogeneous density and pressure is algebraically invertible and therefore supplies a representation, not a predictive closure. Predictivity enters only after assuming that the filling energy is a continuous, rigid-motion invariant, additive valuation on convex bodies in $\mathbb{R}^4$. On the round homothetic family this forces the five dilation weights $S_{\rm val} = \{1, 0, -1, -2, -3\}$ and the annihilator $P_{\rm val}(\mathscr{D})\varepsilon = 0$, with $P_{\rm val}(z) = (z-1)z(z+1)(z+2)(z+3)$. Hence the polynomial dilation action is the semisimple algebra $\mathbb{R}[z]/(P_{\rm val}) \simeq \mathbb{R}^5$.

The resulting invariant calculus includes primitive branch projectors, support ideals, derivative and sampled Hankel pencils, inertia sign certificates, exact observable jet inequalities, and complete normalization/pivot quotient coordinates. For positive mixtures the spectral fractions form an exponential family with $\mathscr{D}^2 \ln \varepsilon = \mathrm{Var}_\pi(s)$ and $\mathscr{D}w = -\mathrm{Var}_\pi(s)/3$. We derive an exact Bhatia–Davis interior-branch remainder and the global rigidity bound $L_F \geq \pi$, with equality precisely for two-extreme-branch support; every positive two-branch orbit has a universal logistic/tanh normal form and constant Schwarzian $\{w, \ln A\} = -(s_+ - s_-)^2/2$. Within the declared boundary-local class, quotienting the powers degenerate with vacuum, curvature, and dust leaves the unique nonstandard direction a^{-1}, corresponding to $w = -2/3$.

We then evaluate the exact nulls on supplied pinned-CLASS and DESI DR2 BAO-only artifacts. The executed source lift obeys $\rho_X \propto a^{-1}$ with maximum residual 1.88×10^{-9}, while the best-fit ledger satisfies the exact coefficient and Hankel/curvature diagnostics. The BAO-only likelihood ratio gives $q_X = 1.364$ and one-sided $p = 0.121$, hence no detection. Because $w_X = -2/3$ is imposed in the executed lift, these numerical spectral recoveries are consistency tests, not independent identification of the valuation roots. No microscopic source action or dimensional amplitude is derived.

Keywords: closed FLRW cosmology; S^3; B^4 filling; convex valuations; finite dilation spectrum; Hankel pencils; Fisher information; homogeneous null tests

I. CLAIM BOUNDARY

This paper proves a finite dilation-module theorem. It does not prove a microscopic origin of dark energy. The logical separation used throughout is

$$\boxed{\begin{array}{c}\text{homogeneous reconstruction} \neq \text{source action} \neq \text{amplitude selection},\\ \text{background/transfer prediction} \neq \text{consistency testing} \neq \text{independent identification},\\ \text{independent identification} \neq \text{likelihood evidence}.\end{array}} \tag{I.1}$$

The first layer is algebraic. The later layers require additional physical input, and a fit to a branch whose exponent was fixed in advance is not by itself an independent identification of that exponent.

Criterion I.1 (Consistency versus identification)**.** A calculation in which a candidate dilation weight is fixed a priori and only its amplitude is inferred tests consistency with that branch. Independent identification requires the spectral location itself to remain free in the inference, for example through free-exponent exponential-sum reconstruction, covariance-aware generalized-eigenvalue estimation, or model comparison against alternatives whose exponents are not pinned to the valuation spectrum. Statistical preference is a further layer and requires an external likelihood.

The main result is the conditional implication

$$\boxed{\begin{array}{c}\text{Euclidean convex valuation closure on } B^4_A\\ \Longrightarrow \text{ a five-dimensional semisimple dilation module.}\end{array}} \tag{I.2}$$

* andre.preece@web.de
† batenin@web.de

The theorem is conditional on the stated valuation hypothesis. Enlarging the admissible class to nonlocal functionals, support-function boundary terms, additional fields, effective-field-theory operators, quotient-topology data, or external dimensional scales can add new invariants. No global uniqueness claim over all dark-sector theories is made.

II. ROUND FILLING GEOMETRY AND THE WORK MAP

Let

$$\Sigma \simeq S^3, \qquad g_A = A^2 \mathrm{d}\Omega_3^2, \qquad A > 0. \tag{II.1}$$

The round volume and the auxiliary Euclidean filling volume are

$$V_3(A) = \mathrm{Vol}(S_A^3) = 2\pi^2 A^3, \qquad V_4(A) = \mathrm{Vol}(B_A^4) = \frac{\pi^2}{2} A^4. \tag{II.2}$$

Thus

$$\frac{\mathrm{d}V_4}{\mathrm{d}A} = V_3, \qquad \frac{V_4}{V_3} = \frac{A}{4}. \tag{II.3}$$

The ball B_A^4 is an auxiliary Euclidean filling of the metric representative (S^3, g_A). It is not a physical extra spatial dimension.

Let $E_4(A)$ be a C^1 filling energy. Define the accumulated boundary density and work pressure by

$$\varepsilon(A) := \frac{E_4(A)}{V_3(A)}, \qquad p(A) := -\frac{\mathrm{d}E_4}{\mathrm{d}V_3} = -\frac{E_4'(A)}{V_3'(A)}. \tag{II.4}$$

Theorem II.1 (Homogeneous work identity)**.** *Let $V_3(A) = 2\pi^2 A^3$. Then*

$$p(A) = -\varepsilon(A) - \frac{A}{3}\varepsilon'(A). \tag{II.5}$$

Consequently, along a homogeneous FLRW history with

$$H = \frac{1}{A}\frac{\mathrm{d}A}{\mathrm{d}\tau},$$

one has

$$\frac{\mathrm{d}\varepsilon}{\mathrm{d}\tau} + 3H(\varepsilon + p) = 0. \tag{II.6}$$

Proof. Write $E_4 = \varepsilon V_3$. Then

$$p = -\frac{\mathrm{d}(\varepsilon V_3)/\mathrm{d}A}{\mathrm{d}V_3/\mathrm{d}A} = -\varepsilon - \frac{V_3}{V_3'}\varepsilon'.$$

Since $V_3'/V_3 = 3/A$, this gives

$$p = -\varepsilon - \frac{A}{3}\varepsilon'.$$

Multiplication of

$$\varepsilon' + \frac{3}{A}(\varepsilon + p) = 0$$

by $\mathrm{d}A/\mathrm{d}\tau = HA$ gives the continuity identity. □

Theorem II.2 (Inverse homogeneous representation)**.** *Every differentiable homogeneous density $\varepsilon(A)$ has a unique adiabatic representative in the round homothetic filling class:*

$$E_4(A) = \varepsilon(A) V_3(A), \qquad \rho_4(A) := \frac{E_4(A)}{V_4(A)} = \frac{4}{A}\varepsilon(A). \tag{II.7}$$

A prescribed pair (ε, p) is represented by the adiabatic work map if and only if

$$p(A) = -\varepsilon(A) - \frac{A}{3}\varepsilon'(A). \tag{II.8}$$

Proof. The definition $E_4 = \varepsilon V_3$ and the ratio $V_3/V_4 = 4/A$ give $\rho_4 = 4\varepsilon/A$. Uniqueness follows from the fixed volume ratio. The pressure condition is exactly the previous theorem. □

Remark II.1 (Limit of the universal map)**.** The inverse map reconstructs only a homogeneous representative. It does not reconstruct microscopic fields, perturbations, sound speed, anisotropic stress, a covariant matter action, or a dimensional amplitude. A predictive theory begins only after a restricted functional class or source completion is supplied.

For a pure power $\varepsilon_s \propto A^s$,

$$w_s := \frac{p_s}{\varepsilon_s} = -1 - \frac{s}{3}. \tag{II.9}$$

III. EUCLIDEAN VALUATION CLOSURE

We use the standard valuation classification as input, not as something reproved here. Hadwiger's theorem says that continuous, Euclidean-invariant, additive valuations on convex bodies in $\mathbb{R}^4$ lie in the intrinsic-volume span [1, 2]. We need only its round homothetic specialization.

Hypothesis III.1 (Euclidean convex valuation closure)**.** The auxiliary filling energy is assumed to be a continuous, Euclidean-invariant, additive valuation on convex bodies in $\mathbb{R}^4$, restricted to the round homothetic family B^4_A. For the smooth round representatives, write

$$E_4^{\mathrm{val}}(A) = \gamma V_4(A) + \sum_{j=0}^{3} \alpha_j \int_{S^3_A} e_j(K)\, \mathrm{d}\Sigma, \tag{III.1}$$

where $K^a{}_b$ is the Weingarten map of the boundary and $e_j(K)$ are the elementary symmetric polynomials in its principal curvatures.

For the round boundary,

$$K^a{}_b = \frac{1}{A}\delta^a_b, \qquad e_j(K) = \binom{3}{j} A^{-j}, \qquad \mathrm{d}\Sigma = A^3 \mathrm{d}\Omega_3. \tag{III.2}$$

Theorem III.1 (Round valuation spectrum)**.** *Under the Euclidean convex valuation closure,*

$$E_4^{\mathrm{val}}(A) = \frac{\pi^2}{2}\gamma A^4 + 2\pi^2 \left(\alpha_0 A^3 + 3\alpha_1 A^2 + 3\alpha_2 A + \alpha_3\right). \tag{III.3}$$

The induced boundary density is

$$\varepsilon(A) = \frac{\gamma A}{4} + \alpha_0 + \frac{3\alpha_1}{A} + \frac{3\alpha_2}{A^2} + \frac{\alpha_3}{A^3}. \tag{III.4}$$

Thus the density dilation spectrum is

$$S_{\mathrm{val}} = \{1, 0, -1, -2, -3\}. \tag{III.5}$$

The corresponding equations of state are

$$s = 1, 0, -1, -2, -3 \quad \Longleftrightarrow \quad w = -\frac{4}{3}, -1, -\frac{2}{3}, -\frac{1}{3}, 0. \tag{III.6}$$

Proof. The bulk term is $\gamma V_4 = (\pi^2/2)\gamma A^4$. Moreover,

$$\int_{S^3_A} e_j(K)\, \mathrm{d}\Sigma = 2\pi^2 \binom{3}{j} A^{3-j}.$$

Division by $V_3 = 2\pi^2 A^3$ gives the displayed density. For a density branch A^s, the work-map equation of state is $w_s = -1 - s/3$. □

Remark III.1 (Normalization)**.** The coefficients γ, α_j absorb conventional normalizations of the intrinsic volumes. The powers of A, the dilation eigenvalues, and the equations of state do not depend on these normalization choices.

IV. FINITE DILATION ALGEBRA

Define

$$\mathscr{D} := A\frac{\mathrm{d}}{\mathrm{d}A}. \tag{IV.1}$$

Equivalently, with $x = \ln(A/A_*)$,

$$\mathscr{D} = \frac{\mathrm{d}}{\mathrm{d}x}. \tag{IV.2}$$

Let

$$\mathcal{V}_{\mathrm{val}} := \mathrm{span}\{A, 1, A^{-1}, A^{-2}, A^{-3}\}. \tag{IV.3}$$

Theorem IV.1 (ODE characterization)**.** *Let $J \subset (0,\infty)$ be an interval and $\varepsilon \in C^5(J)$. Then*

$$P_{\mathrm{val}}(\mathscr{D})\varepsilon = 0, \qquad P_{\mathrm{val}}(z) = (z-1)z(z+1)(z+2)(z+3), \tag{IV.4}$$

if and only if

$$\varepsilon(A) = C_1 A + C_0 + C_{-1}A^{-1} + C_{-2}A^{-2} + C_{-3}A^{-3}. \tag{IV.5}$$

Proof. In the variable $x = \ln(A/A_*)$, the equation becomes a constant-coefficient ODE with simple characteristic roots $1, 0, -1, -2, -3$. Hence its solution space is

$$\mathrm{span}\{e^x, 1, e^{-x}, e^{-2x}, e^{-3x}\},$$

equivalently the displayed powers of A. The converse is immediate. □

For a nonzero profile

$$\varepsilon(A) = \sum_{s \in S_{\mathrm{val}}} \widehat{C}_s A^s, \tag{IV.6}$$

define

$$\mathrm{supp}_{\mathscr{D}}(\varepsilon) := \{s : \widehat{C}_s \neq 0\}, \qquad m_\varepsilon(z) := \prod_{s \in \mathrm{supp}_{\mathscr{D}}(\varepsilon)} (z - s). \tag{IV.7}$$

Theorem IV.2 (Elementwise annihilator)**.** *For every nonzero $\varepsilon \in \mathcal{V}_{\mathrm{val}}$,*

$$\mathrm{Ann}_{\mathscr{D}}(\varepsilon) := \{p \in \mathbb{R}[z] : p(\mathscr{D})\varepsilon = 0\} = (m_\varepsilon). \tag{IV.8}$$

In particular,

$$m_\varepsilon(z) \mid P_{\mathrm{val}}(z). \tag{IV.9}$$

Proof. The powers A^s with distinct real exponents are linearly independent on J, and $p(\mathscr{D})A^s = p(s)A^s$. Hence $p(\mathscr{D})\varepsilon = 0$ exactly when $p(s) = 0$ on every active spectral value. The monic polynomial of least degree with these roots is m_ε. □

Theorem IV.3 (Semisimple operator algebra)**.** *Let*

$$\varrho : \mathbb{R}[z] \longrightarrow \mathrm{End}(\mathcal{V}_{\mathrm{val}}), \qquad p \longmapsto p(\mathscr{D})|_{\mathcal{V}_{\mathrm{val}}}. \tag{IV.10}$$

Then

$$\ker \varrho = (P_{\mathrm{val}}), \tag{IV.11}$$

and hence

$$\boxed{\mathfrak{A}_{\mathscr{D}} := \varrho(\mathbb{R}[z]) \simeq \mathbb{R}[z]/(P_{\mathrm{val}}) \simeq \prod_{s \in S_{\mathrm{val}}} \mathbb{R} \simeq \mathbb{R}^5.} \tag{IV.12}$$

The module $\mathcal{V}_{\mathrm{val}}$ is faithful over $\mathfrak{A}_{\mathscr{D}}$. It is not a pointwise multiplication algebra, since $A \cdot A = A^2 \notin \mathcal{V}_{\mathrm{val}}$.

Proof. The kernel consists of polynomials vanishing at all five eigenvalues $1, 0, -1, -2, -3$, hence is the ideal (P_{val}). Since P_{val} has five simple real roots, the Chinese remainder theorem gives the product decomposition. □

Corollary IV.1 (Primitive projectors)**.** *For $s \in S_{\mathrm{val}}$, define*

$$e_s(z) := \prod_{\substack{t \in S_{\mathrm{val}} \\ t \neq s}} \frac{z - t}{s - t}, \qquad \mathsf{P}_s := e_s(\mathscr{D}). \tag{IV.13}$$

Then

$$\mathsf{P}_s \mathsf{P}_t = \delta_{st} \mathsf{P}_s, \qquad \sum_s \mathsf{P}_s = 1, \tag{IV.14}$$

and

$$\mathsf{P}_s \varepsilon = \widehat{C}_s A^s. \tag{IV.15}$$

These projectors are the primitive idempotents of $\mathfrak{A}_{\mathscr{D}}$.

Explicitly,

$$e_1(z) = \frac{z(z+1)(z+2)(z+3)}{24}, \tag{IV.16}$$

$$e_0(z) = -\frac{(z-1)(z+1)(z+2)(z+3)}{6}, \tag{IV.17}$$

$$e_{-1}(z) = \frac{(z-1)z(z+2)(z+3)}{4}, \tag{IV.18}$$

$$e_{-2}(z) = -\frac{(z-1)z(z+1)(z+3)}{6}, \tag{IV.19}$$

$$e_{-3}(z) = \frac{(z-1)z(z+1)(z+2)}{24}. \tag{IV.20}$$

Definition IV.1 (Spectral subspaces)**.** For $S \subseteq S_{\mathrm{val}}$, define

$$\mathcal{V}_S := \mathrm{span}\{A^s : s \in S\}. \tag{IV.21}$$

Corollary IV.2 (Spectral lattice)**.** *For $S, T \subseteq S_{\mathrm{val}}$,*

$$\mathcal{V}_S \cap \mathcal{V}_T = \mathcal{V}_{S \cap T}, \qquad \mathcal{V}_S + \mathcal{V}_T = \mathcal{V}_{S \cup T}. \tag{IV.22}$$

Moreover,

$$\mathrm{Ann}_{\mathscr{D}}(\mathcal{V}_S) = \left(\prod_{s \in S} (z - s) \right). \tag{IV.23}$$

V. POSITIVE CONE AND MOMENT TESTS

Assume that all active branch energies are nonnegative:

$$\mathsf{P}_s \varepsilon(A) \geq 0, \qquad \varepsilon(A) > 0. \tag{V.1}$$

Define the branch fractions

$$\pi_s(A) := \frac{\mathsf{P}_s \varepsilon(A)}{\varepsilon(A)}. \tag{V.2}$$

Then

$$\sum_s \pi_s = 1. \tag{V.3}$$

Thus π_s is a finite probability distribution on S_{val}.

Define the dilation moments

$$\mathfrak{m}_n(A) := \frac{\mathscr{D}^n \varepsilon(A)}{\varepsilon(A)} = \sum_s s^n \pi_s(A). \tag{V.4}$$

Theorem V.1 (Log-convexity and monotone equation of state)**.** *For a positive valuation mixture,*

$$\mathscr{D}\mathfrak{m}_1 = \mathfrak{m}_2 - \mathfrak{m}_1^2 = \mathrm{Var}_\pi(s) \geq 0. \tag{V.5}$$

Equivalently,

$$\boxed{\mathscr{D}^2 \ln(\varepsilon/\varepsilon_*) = \mathrm{Var}_\pi(s) \geq 0,} \tag{V.6}$$

where $\varepsilon_* > 0$ is an arbitrary fixed normalization. Since

$$w = -1 - \frac{\mathfrak{m}_1}{3}, \tag{V.7}$$

one also has

$$\boxed{\mathscr{D}w = -\frac{1}{3}\,\mathrm{Var}_\pi(s) \leq 0.} \tag{V.8}$$

Proof. Differentiate $\mathfrak{m}_1 = (\mathscr{D}\varepsilon)/\varepsilon$:

$$\mathscr{D}\mathfrak{m}_1 = \frac{\mathscr{D}^2\varepsilon}{\varepsilon} - \left(\frac{\mathscr{D}\varepsilon}{\varepsilon}\right)^2 = \mathfrak{m}_2 - \mathfrak{m}_1^2.$$

The moment formula identifies this with the variance of s. The equation-of-state identity follows from $w = -1-\mathfrak{m}_1/3$. □

Corollary V.1 (CPL tangent sign)**.** *For the local expansion*

$$w(a) = w_0 + w_a(1-a) + O((1-a)^2),$$

one has

$$w_a = -\mathscr{D}w|_{a=1}.$$

Every positive valuation mixture therefore satisfies

$$w_a \geq 0. \tag{V.9}$$

Theorem V.2 (Bhatia–Davis envelope)**.** *Let a positive mixture have active support contained in $[m, M]$, and let*

$$\bar{s} := \sum_s s\pi_s = -3(1+w). \tag{V.10}$$

Then

$$0 \leq -\mathscr{D}w = \frac{1}{3}\,\mathrm{Var}_\pi(s) \leq \frac{1}{3}(M-\bar{s})(\bar{s}-m). \tag{V.11}$$

Equality in the upper bound occurs precisely when the active probability is supported on the two endpoints m and M [3].

Corollary V.2 (Endpoint saturation in the boundary cone)**.** *For the positive boundary spectrum $S_\partial = \{0, -1, -2, -3\} \subset [-3, 0]$,*

$$0 \leq -\mathscr{D}w \leq -3w(1+w). \tag{V.12}$$

Equality for all A occurs if and only if the active support is contained in $\{0, -3\}$.

Theorem V.3 (Exact Bhatia–Davis slack and local interior-branch detector)**.** *Let a positive mixture have support contained in $[m, M]$ and mean $\bar{s} = \kappa_1$. Define the variance slack*

$$\Delta_{mM} := (M-\bar{s})(\bar{s}-m) - \kappa_2. \tag{V.13}$$

Then the Bhatia–Davis remainder has the exact branch decomposition

$$\boxed{\Delta_{mM} = \sum_s \pi_s(M-s)(s-m) \geq 0.} \tag{V.14}$$

It vanishes at one finite scale if and only if every active branch lies at the two endpoints $\{m, M\}$. Thus Δ_{mM} is a first-derivative local detector of probability carried by interior spectral branches.

For the positive boundary support $\{-3, -2, -1, 0\}$,

$$\boxed{\Delta_\partial = 2(\pi_{-1} + \pi_{-2}) = 3\big[\mathscr{D}w - 3w(1+w)\big] \geq 0.} \tag{V.15}$$

Hence the total instantaneous occupation of the two interior boundary branches is recovered from only w and its first dilation derivative:

$$\boxed{\pi_{-1} + \pi_{-2} = \frac{3}{2}\big[\mathscr{D}w - 3w(1+w)\big].} \tag{V.16}$$

Proof. Using $\kappa_2 = \mathbb{E}[s^2] - \bar{s}^2$,

$$\begin{aligned}(M - \bar{s})(\bar{s} - m) - \kappa_2 &= (M+m)\bar{s} - Mm - \mathbb{E}[s^2] \\ &= \mathbb{E}[(M-s)(s-m)]\,.\end{aligned}$$

Every summand is nonnegative on $[m, M]$ and is zero only at m or M. For $m = -3$ and $M = 0$, the polynomial $(0 - s)(s + 3)$ takes the values $0, 2, 2, 0$ on $s = -3, -2, -1, 0$, respectively. Finally $\bar{s} = -3(1 + w)$ and $\kappa_2 = -3\mathscr{D}w$, which gives the observable formula. □

Theorem V.4 (Hankel positivity and rank)**.** *For $r \geq 0$, define*

$$H_r(A) := \big[\mathscr{D}^{i+j}\varepsilon(A)\big]_{i,j=0}^{r}\,.$$

For a positive valuation mixture,

$$H_r(A) \succeq 0. \tag{V.17}$$

If exactly k distinct dilation branches are active, then

$$\operatorname{rank} H_r = \min\{r+1, k\}. \tag{V.18}$$

Proof. Let $v_s = (1, s, \ldots, s^r)^T$. Then

$$H_r = \sum_s (\widehat{C}_s A^s) v_s v_s^T,$$

a positive sum of rank-one matrices. Distinct spectral values give the Vandermonde rank statement. □

VI. LOCAL SPECTRAL FINGERPRINTS, COMPLETE QUOTIENT INVARIANTS, AND INFORMATION GEOMETRY

The semisimple dilation algebra admits a stronger local formulation that does not require a global branch fit. The active dilation support can be recovered from a single sufficiently high derivative jet of the homogeneous profile.

Theorem VI.1 (Hankel-pencil spectral fingerprint)**.** *Let*

$$\varepsilon(A) = \sum_{j=1}^{k} C_j A^{s_j}, \qquad C_j \neq 0, \qquad s_i \neq s_j \quad (i \neq j), \tag{VI.1}$$

and write

$$M_n(A) := \mathscr{D}^n \varepsilon(A). \tag{VI.2}$$

Define the two $k \times k$ Hankel matrices

$$H_0(A) := [M_{i+j}(A)]_{i,j=0}^{k-1}, \qquad H_1(A) := [M_{i+j+1}(A)]_{i,j=0}^{k-1}. \tag{VI.3}$$

Then

$$\det H_0(A) = \Delta(s)^2 \prod_{j=1}^{k} C_j A^{s_j}, \qquad \Delta(s) := \prod_{1\le i<j\le k} (s_j - s_i), \tag{VI.4}$$

and therefore H_0 is invertible. Moreover,

$$\boxed{\frac{\det\big(\lambda H_0 - H_1\big)}{\det H_0} = \prod_{j=1}^{k} (\lambda - s_j) = m_\varepsilon(\lambda).} \tag{VI.5}$$

Hence the generalized eigenvalues of the pencil (H_1, H_0) are exactly the active dilation weights. The polynomial on the left is independent of A and of all branch amplitudes.

Proof. Let $V_{ij} = s_j^i$ for $i = 0, \dots, k-1$, let

$$W(A) = \operatorname{diag}(C_1 A^{s_1}, \dots, C_k A^{s_k}), \qquad S = \operatorname{diag}(s_1, \dots, s_k).$$

Then

$$H_0 = VWV^T, \qquad H_1 = VWSV^T. \tag{VI.6}$$

Since the s_j are distinct, V is a nonsingular Vandermonde matrix and

$$\det H_0 = (\det V)^2 \det W = \Delta(s)^2 \prod_j C_j A^{s_j}.$$

Likewise,

$$\lambda H_0 - H_1 = VW(\lambda I - S)V^T.$$

Taking determinants and dividing by $\det H_0$ gives the stated characteristic polynomial. □

Corollary VI.1 (Amplitude-free determinant flow)**.** *Under the hypotheses of theorem VI.1,*

$$\boxed{\mathscr{D} \ln|\det H_0| = \sum_{j=1}^{k} s_j, \qquad \mathscr{D}^2 \ln|\det H_0| = 0.} \tag{VI.7}$$

Thus the logarithmic determinant slope is a local, amplitude-independent support invariant. For the principal supports in this paper,

$$S_{\mathrm{val}} = \{1, 0, -1, -2, -3\}: \qquad \mathscr{D} \ln|\det H_0| = -5, \tag{VI.8}$$
$$S_{\partial} = \{0, -1, -2, -3\}: \qquad \mathscr{D} \ln|\det H_0| = -6, \tag{VI.9}$$
$$S_{\mathrm{end}} = \{0, -3\}: \qquad \mathscr{D} \ln|\det H_0| = -3. \tag{VI.10}$$

If all listed branches are active, the corresponding exact determinant prefactors are

$$\det H_0^{\mathrm{val}} = 82944 \Big(\prod_{s\in S_{\mathrm{val}}} \widehat{C}_s \Big) A^{-5}, \tag{VI.11}$$

$$\det H_0^{\partial} = 144 \Big(\prod_{s\in S_{\partial}} \widehat{C}_s \Big) A^{-6}, \tag{VI.12}$$

$$\det H_0^{\mathrm{end}} = 9\, \widehat{C}_0 \widehat{C}_{-3} A^{-3}. \tag{VI.13}$$

Remark VI.1 (Local spectroscopy)**.** The pencil identity strengthens the elementwise annihilator theorem: the minimal annihilator can be reconstructed directly from a local derivative jet without first solving for the branch amplitudes. Rank detects the number of active branches; the generalized eigenvalues identify which branches are active; and the determinant flow provides an independent support checksum. In noisy applications the same structure can be implemented by covariance-aware matrix-pencil or Prony methods rather than raw differentiation.

Theorem VI.2 (Complete normalized-jet positivity certificate)**.** *Let $\varepsilon \in \mathcal{V}_{\mathrm{val}}$ with $\varepsilon > 0$, and define*

$$\mathfrak{m}_n := \frac{\mathscr{D}^n \varepsilon}{\varepsilon}, \qquad \mathfrak{m}_0 = 1. \tag{VI.14}$$

Then the five branch fractions are recovered pointwise from the fourth-order normalized jet by

$$\pi_1 = \frac{\mathfrak{m}_4 + 6\mathfrak{m}_3 + 11\mathfrak{m}_2 + 6\mathfrak{m}_1}{24}, \tag{VI.15}$$

$$\pi_0 = 1 + \frac{5\mathfrak{m}_1 - 5\mathfrak{m}_2 - 5\mathfrak{m}_3 - \mathfrak{m}_4}{6}, \tag{VI.16}$$

$$\pi_{-1} = \frac{\mathfrak{m}_4 + 4\mathfrak{m}_3 + \mathfrak{m}_2 - 6\mathfrak{m}_1}{4}, \tag{VI.17}$$

$$\pi_{-2} = \frac{-\mathfrak{m}_4 - 3\mathfrak{m}_3 + \mathfrak{m}_2 + 3\mathfrak{m}_1}{6}, \tag{VI.18}$$

$$\pi_{-3} = \frac{\mathfrak{m}_4 + 2\mathfrak{m}_3 - \mathfrak{m}_2 - 2\mathfrak{m}_1}{24}. \tag{VI.19}$$

Consequently, inside the valuation closure, nonnegative branch energy is equivalent to the five explicit inequalities

$$\boxed{\pi_s \geq 0 \quad (s \in S_{\mathrm{val}}).} \tag{VI.20}$$

Equivalently, the normalized jet $(\mathfrak{m}_1, \ldots, \mathfrak{m}_4)$ lies in the convex hull of the five moment-curve vertices

$$(s, s^2, s^3, s^4), \qquad s \in S_{\mathrm{val}}. \tag{VI.21}$$

Proof. By the primitive-projector formula,

$$\pi_s = \frac{e_s(\mathscr{D})\varepsilon}{\varepsilon}.$$

Expanding the five Lagrange polynomials e_s gives the displayed linear expressions in the normalized moments. Since the projectors sum to the identity, the recovered fractions sum to one. Positivity of every branch is therefore equivalent to nonnegativity of these five barycentric coordinates. □

Corollary VI.2 (Observable sharp jet cone)**.** *Define*

$$u_1 := \mathscr{D}w, \qquad u_2 := \mathscr{D}^2 w, \qquad u_3 := \mathscr{D}^3 w.$$

Since

$$\kappa_1 = -3(1 + w), \qquad \kappa_n = -3\mathscr{D}^{n-1} w \quad (n \geq 2),$$

and $\mathfrak{m}_n = B_n(\kappa_1, \ldots, \kappa_n)$, the five barycentric coordinates of theorem VI.2 can be written directly as observable jet polynomials:

$$\pi_1 = \frac{1}{8}\Big[27w^4 + 54w^3 + 33w^2 + 6w - 54w^2u_1 - 54wu_1 + 12wu_2 + 9u_1^2 - 11u_1 + 6u_2 - u_3\Big], \tag{VI.22}$$

$$\pi_0 = -\frac{1}{2}\Big[27w^4 + 63w^3 + 42w^2 + 8w - 54w^2u_1 - 63wu_1 + 12wu_2 + 9u_1^2 - 14u_1 + 7u_2 - u_3\Big], \tag{VI.23}$$

$$\pi_{-1} = \frac{3}{4}\Big[27w^4 + 72w^3 + 57w^2 + 12w - 54w^2u_1 - 72wu_1 + 12wu_2 + 9u_1^2 - 19u_1 + 8u_2 - u_3\Big], \tag{VI.24}$$

$$\pi_{-2} = -\frac{1}{2}\Big[27w^4 + 81w^3 + 78w^2 + 24w - 54w^2u_1 - 81wu_1 + 12wu_2 + 9u_1^2 - 26u_1 + 9u_2 - u_3\Big], \tag{VI.25}$$

$$\pi_{-3} = \frac{1}{8}\Big[27w^4 + 90w^3 + 105w^2 + 50w + 8 - 54w^2u_1 - 90wu_1 + 12wu_2 + 9u_1^2 - 35u_1 + 10u_2 - u_3\Big]. \tag{VI.26}$$

Therefore the complete positive valuation 4-jet is characterized by the five sharp inequalities

$$\boxed{\pi_s(w,u_1,u_2,u_3)\geq 0 \qquad (s\in S_{\mathrm{val}}).} \tag{VI.27}$$

These inequalities are the exact pullback of the moment-simplex facets under the Bell/cumulant change of coordinates; the admissible region need not itself be a polytope in the nonlinear coordinates (w,u_1,u_2,u_3). Support faces are obtained by setting the corresponding barycentric coordinates to zero. In particular, the boundary-local face is the exact third-order observable equation $\pi_1=0$, i.e.

$$\boxed{\begin{aligned}\mathscr{D}^3 w = {}& 27w^4+54w^3+33w^2+6w-54w^2\mathscr{D}w-54w\mathscr{D}w\\ &+12w\mathscr{D}^2w+9(\mathscr{D}w)^2-11\mathscr{D}w+6\mathscr{D}^2w.\end{aligned}} \tag{VI.28}$$

Proof. Substitute the Bell-polynomial relations

$$\begin{aligned}\mathfrak{m}_1&=\kappa_1,\\ \mathfrak{m}_2&=\kappa_1^2+\kappa_2,\\ \mathfrak{m}_3&=\kappa_1^3+3\kappa_1\kappa_2+\kappa_3,\\ \mathfrak{m}_4&=\kappa_1^4+6\kappa_1^2\kappa_2+3\kappa_2^2+4\kappa_1\kappa_3+\kappa_4\end{aligned}$$

into theorem VI.2 and use $\kappa_1=-3(1+w)$, $\kappa_{n\geq 2}=-3\mathscr{D}^{n-1}w$. The boundary equation is simply the facet condition $\pi_1=0$ solved for $\mathscr{D}^3w$. □

Corollary VI.3 (Sharp boundary 2-jet simplex)**.** *On the positive boundary face*

$$S_\partial=\{0,-1,-2,-3\},$$

only the first three normalized moments are needed. With

$$u_1=\mathscr{D}w, \qquad u_2=\mathscr{D}^2w,$$

the four branch fractions are

$$\boxed{\begin{aligned}\pi_0&=\frac{1}{2}\Big(9wu_1+3u_1-u_2-9w^3-9w^2-2w\Big),\\ \pi_{-1}&=-\frac{3}{2}\Big(9wu_1+4u_1-u_2-9w^3-12w^2-3w\Big),\\ \pi_{-2}&=\frac{3}{2}\Big(9wu_1+5u_1-u_2-9w^3-15w^2-6w\Big),\\ \pi_{-3}&=-\frac{1}{2}\Big(9wu_1+6u_1-u_2-9w^3-18w^2-11w-2\Big).\end{aligned}} \tag{VI.29}$$

Therefore a reconstructed boundary profile lies in the positive boundary cone if and only if all four expressions are nonnegative and sum to one. In particular, the two interior fractions are resolved separately from only $(w,\mathscr{D}w,\mathscr{D}^2w)$, while theorem V.3 gives their sum from $(w,\mathscr{D}w)$ alone.

Proof. Set the bulk barycentric coordinate π_1 in corollary VI.2 to zero and use equation (VI.28) to eliminate $\mathscr{D}^3w$. Equivalently, apply the degree-three primitive projectors for the support $\{0,-1,-2,-3\}$ directly to the first three normalized moments. □

Theorem VI.3 (Cumulant hierarchy and nonlinear amplitude-free closure)**.** *For a positive valuation mixture, let $x=\ln(A/A_*)$ and define*

$$\kappa_n:=\mathscr{D}^n\ln(\varepsilon/\varepsilon_*), \qquad n\geq 1. \tag{VI.30}$$

Then κ_n is the nth cumulant of the finite spectral probability distribution $\pi_s(A)$. In particular,

$$\kappa_1=\mathbb{E}_\pi[s], \qquad \kappa_2=\mathrm{Var}_\pi(s), \qquad \mathscr{D}\kappa_n=\kappa_{n+1}. \tag{VI.31}$$

The branch fractions obey the exact replicator equation

$$\boxed{\mathscr{D}\pi_s = (s-\kappa_1)\pi_s.} \tag{VI.32}$$

Furthermore, writing $B_n(\kappa_1,\dots,\kappa_n)$ for the complete exponential Bell polynomial, the full valuation closure is equivalent to the amplitude-free nonlinear identity

$$\boxed{B_5 + 5B_4 + 5B_3 - 5B_2 - 6B_1 = 0.} \tag{VI.33}$$

For the boundary face this reduces to

$$\boxed{B_4 + 6B_3 + 11B_2 + 6B_1 = 0,} \tag{VI.34}$$

and for the endpoint class to

$$\boxed{B_2 + 3B_1 = 0.} \tag{VI.35}$$

These identities contain no dimensional amplitude.

Proof. With $x = \ln(A/A_*)$,

$$\varepsilon(x) = \sum_s \widehat{C}_s A_*^s e^{sx}$$

is a finite exponential sum. After normalization by ε, its logarithm is the cumulant-generating potential of the tilted discrete measure π_s. Differentiation therefore generates cumulants and gives $\mathscr{D}\pi_s = (s-\kappa_1)\pi_s$. Finally,

$$\frac{\mathscr{D}^n\varepsilon}{\varepsilon} = B_n(\kappa_1,\dots,\kappa_n).$$

Substitution into $P_{\mathrm{val}}(\mathscr{D})\varepsilon = 0$, using

$$P_{\mathrm{val}}(z) = z^5 + 5z^4 + 5z^3 - 5z^2 - 6z,$$

gives the full nonlinear closure. The boundary and endpoint formulas follow from

$$z(z+1)(z+2)(z+3) = z^4 + 6z^3 + 11z^2 + 6z, \qquad z(z+3) = z^2 + 3z.$$

□

Corollary VI.4 (Observable scalar differential closure)**.** *Let*

$$u_n := \mathscr{D}^n w.$$

The full five-branch valuation annihilator is equivalent, wherever $\varepsilon > 0$, to the fourth-order amplitude-free equation

$$\boxed{\begin{aligned} 0 = {} & u_4 - 10u_3 + 35u_2 - 50u_1 + 90u_1^2 - 30u_1u_2 \\ & + 24w + 150w^2 + 315w^3 + 270w^4 + 81w^5 \\ & - 315wu_1 - 540w^2u_1 - 270w^3u_1 \\ & + 120wu_2 + 90w^2u_2 + 135wu_1^2 - 15wu_3. \end{aligned}} \tag{VI.36}$$

The boundary face lowers the order to the third-order relation equation (VI.28), while the endpoint support $\{0,-3\}$ obeys the first-order saturation law

$$\boxed{\mathscr{D}w = 3w(1+w).} \tag{VI.37}$$

Thus each support reduction appears as an exact order reduction of the observable equation-of-state dynamics.

Proof. Substitute $\kappa_1 = -3(1+w)$ and $\kappa_{n\geq 2} = -3u_{n-1}$ into the Bell-polynomial identities of theorem VI.3 and expand. The endpoint equation follows from $B_2 + 3B_1 = 0$. □

Corollary VI.5 (Information-geometric form of the equation-of-state flow)**.** *For a positive mixture,*

$$\boxed{\mathcal{I}_x := \sum_s \pi_s \big(\mathscr{D} \ln \pi_s\big)^2 = \mathrm{Var}_\pi(s) = \kappa_2 = -3\mathscr{D} w.} \tag{VI.38}$$

Thus the monotone equation-of-state flow is exactly one third of the Fisher information of the one-parameter spectral exponential family, with sign fixed by the work-map convention.

Corollary VI.6 (Asymptotic support theorem)**.** *Assume positive coefficients on an active support S and let*

$$s_- := \min S, \qquad s_+ := \max S.$$

Then

$$\lim_{A\to 0^+} \kappa_1 = s_-, \qquad \lim_{A\to\infty} \kappa_1 = s_+, \tag{VI.39}$$

and therefore

$$\lim_{A\to 0^+} w = -1 - \frac{s_-}{3}, \qquad \lim_{A\to\infty} w = -1 - \frac{s_+}{3}. \tag{VI.40}$$

For a positive boundary profile with both endpoint branches active, this gives the exact interpolation

$$w : 0 \longrightarrow -1 \tag{VI.41}$$

as A grows from zero to infinity. If the positive bulk branch $s = 1$ is active, the late-time asymptote instead becomes $w \to -4/3$.

Theorem VI.4 (Hankel inertia and amplitude-signature certificate)**.** *Under the hypotheses of theorem VI.1, let*

$$\nu_+(H_0), \qquad \nu_-(H_0), \qquad \nu_0(H_0)$$

denote the positive, negative, and zero inertia indices of the $k \times k$ derivative Hankel matrix. Then

$$\boxed{\big(\nu_+(H_0), \nu_-(H_0), \nu_0(H_0)\big) = \big(\#\{j : C_j > 0\}, \#\{j : C_j < 0\}, 0\big)\,.} \tag{VI.42}$$

In particular, for a fully active k-branch support,

$$\boxed{H_0 \succ 0 \iff C_j > 0 \quad \textit{for all } j.} \tag{VI.43}$$

For a larger Hankel truncation H_r with $r+1 \geq k$, the same nonzero inertia is supplemented by $r+1-k$ zero eigenvalues.

Proof. The factorization in theorem VI.1 is

$$H_0 = VWV^T, \qquad W = \mathrm{diag}(C_1 A^{s_1}, \ldots, C_k A^{s_k}).$$

Since V is invertible and $A^{s_j} > 0$, Sylvester's law of inertia gives

$$\mathrm{Inertia}(H_0) = \mathrm{Inertia}(W),$$

which is exactly the stated sign count. For H_r with $r + 1 > k$, write

$$H_r = V_r W V_r^T,$$

where the rectangular Vandermonde matrix V_r has column rank k. Its image carries the same nonzero quadratic form as W, while the orthogonal complement of the image contributes $r + 1 - k$ zeros. □

Corollary VI.7 (Power-sum spectral checksums)**.** *With*

$$K := H_0^{-1} H_1,$$

one has

$$K \sim \mathrm{diag}(s_1, \ldots, s_k), \tag{VI.44}$$

and hence, for every integer $n \geq 1$,

$$\boxed{\operatorname{tr}(K^n) = \sum_{j=1}^{k} s_j^n.} \tag{VI.45}$$

These traces are local and independent of all amplitudes. For the full valuation support,

$$\big(\operatorname{tr} K, \operatorname{tr} K^2, \operatorname{tr} K^3, \operatorname{tr} K^4\big) = (-5, 15, -35, 99), \tag{VI.46}$$

while for the boundary support they are

$$(-6, 14, -36, 98). \tag{VI.47}$$

Proof. From $H_0 = VWV^T$ and $H_1 = VWSV^T$, with $S = \operatorname{diag}(s_j)$ and $WS = SW$,

$$H_0^{-1} H_1 = (V^T)^{-1} S V^T.$$

Thus K is similar to S, and the trace identities follow. □

Theorem VI.5 (Complete scale-pivot quotient invariants on consecutive positive support)**.** *Let a positive active support consist of $k \geq 2$ consecutive integer weights*

$$S = \{s_{\min}, s_{\min} + 1, \ldots, s_{\max}\},$$

and write

$$\varepsilon_s(A) = C_s A^s, \qquad C_s > 0.$$

Choose arbitrary positive references A_* and ε_* and define the dimensionless coefficients

$$c_s := \frac{C_s A_*^s}{\varepsilon_*} > 0. \tag{VI.48}$$

For every interior weight define

$$\boxed{\mathcal{Q}_s := \frac{\varepsilon_{s+1}\varepsilon_{s-1}}{\varepsilon_s^2} = \frac{c_{s+1}c_{s-1}}{c_s^2} = \frac{C_{s+1}C_{s-1}}{C_s^2}.} \tag{VI.49}$$

Then

$$\mathscr{D}\mathcal{Q}_s = 0. \tag{VI.50}$$

Moreover, the $k-2$ quantities $\{\mathcal{Q}_s\}$ form a complete set of invariants under the two-parameter reference action

$$c_s \longmapsto e^{\alpha+\beta s} c_s, \tag{VI.51}$$

which represents a change of overall normalization and logarithmic scale pivot. Thus two positive dimensionless coefficient vectors have the same $\{\mathcal{Q}_s\}$ if and only if they lie on the same reference-change orbit.

Proof. The scale independence is immediate from

$$(s+1) + (s-1) - 2s = 0.$$

Put $u_s = \ln c_s$, which is well defined because c_s is positive and dimensionless. The two-parameter action is

$$u_s \longmapsto u_s + \alpha + \beta s.$$

The logarithms

$$\ln \mathcal{Q}_s = u_{s+1} - 2u_s + u_{s-1}$$

are the discrete second differences of u_s. They annihilate the two-dimensional affine subspace spanned by 1 and s. Conversely, equality of all second differences implies that the difference of two coefficient-log vectors is affine in s, hence is of the form $\alpha + \beta s$. Therefore the $k-2$ second differences separate the quotient orbits. □

Corollary VI.8 (Valuation-moduli coordinates)**.** *For the full positive valuation support there are exactly three independent quotient invariants,*

$$\boxed{\mathcal{Q}_0 = \frac{C_1 C_{-1}}{C_0^2}, \qquad \mathcal{Q}_{-1} = \frac{C_0 C_{-2}}{C_{-1}^2}, \qquad \mathcal{Q}_{-2} = \frac{C_{-1} C_{-3}}{C_{-2}^2}.} \tag{VI.52}$$

For the positive boundary support there are exactly two, $\mathcal{Q}_{-1}$ and $\mathcal{Q}_{-2}$. For a two-branch endpoint support there is no nontrivial continuous quotient modulus after normalization and pivot rescaling.

Theorem VI.6 (Complete invariant classification on a positive support stratum)**.** *Let*

$$S = \{s_1 < \cdots < s_k\}, \qquad k \geq 2,$$

be any fixed active support of distinct real dilation weights, and let

$$\mathcal{C}_S = (\mathbb{R}_{>0})^k$$

be the space of positive dimensionless coefficient vectors $c = (c_1, \ldots, c_k)$. Let $G = \mathbb{R}^2$ act by

$$(\alpha, \beta) \cdot c_i = e^{\alpha + \beta s_i} c_i. \tag{VI.53}$$

Then the action is free and the quotient $\mathcal{C}_S/G$ has dimension $k-2$. In logarithmic coordinates $u_i = \ln c_i$, choose any basis

$$a^{(1)}, \ldots, a^{(k-2)}$$

of the kernel of

$$B = \begin{pmatrix} 1 & \cdots & 1 \\ s_1 & \cdots & s_k \end{pmatrix}. \tag{VI.54}$$

The $k-2$ positive monomials

$$\boxed{I_r(c) := \exp\bigl(a^{(r)} \cdot u\bigr) = \prod_{i=1}^{k} c_i^{a_i^{(r)}}, \qquad r = 1, \ldots, k-2,} \tag{VI.55}$$

separate the G-orbits. Consequently every smooth invariant of positive coefficient data that is independent of the normalization and scale pivot factors through $(I_1, \ldots, I_{k-2})$.

If zero coefficients are admitted, the positive cone is stratified by its active support. The support itself is a discrete invariant, and the statement above applies on each positive support stratum. Thus, within the declared finite homogeneous closure, there are no additional independent continuous coefficient moduli beyond $k-2$ quotient coordinates.

Proof. In logarithmic coordinates the group action is translation by

$$\alpha \mathbf{1} + \beta \mathbf{s}, \qquad \mathbf{s} = (s_1, \ldots, s_k).$$

Because at least two s_i are distinct, the stabilizer is trivial. Hence the orbit space is the vector quotient

$$\mathbb{R}^k / \operatorname{span}\{\mathbf{1}, \mathbf{s}\}$$

of dimension $k-2$. The rows of B span the orbit directions, while $\ker B$ is their Euclidean annihilator. Therefore the linear functionals $a^{(r)} \cdot u$ form coordinates on the quotient. Exponentiation gives positive monomial invariants. If two points have equal values of all I_r, their logarithmic difference is orthogonal to $\ker B$, hence lies in $\operatorname{span}\{\mathbf{1}, \mathbf{s}\}$ and the two coefficient vectors differ by a G transformation. Any smooth invariant constant on G-orbits therefore factors through these quotient coordinates. □

Corollary VI.9 (Triple cross-ratios for arbitrary support)**.** *For any $1 \leq i < j < \ell \leq k$, define*

$$\boxed{\mathcal{Q}_{ij\ell} := c_i^{s_\ell - s_j} c_j^{s_i - s_\ell} c_\ell^{s_j - s_i}.} \tag{VI.56}$$

Then $\mathcal{Q}_{ij\ell}$ is invariant under the reference action. The anchored set

$$\{\mathcal{Q}_{1jk} : j = 2, \dots, k-1\} \tag{VI.57}$$

contains $k-2$ independent invariants and is therefore a complete coordinate system on the positive quotient. For three consecutive integer weights $(s-1, s, s+1)$ the formula reduces to

$$\mathcal{Q}_{s-1,s,s+1} = \frac{c_{s-1}c_{s+1}}{c_s^2} = \mathcal{Q}_s.$$

Proof. For the exponent vector of $\mathcal{Q}_{ij\ell}$ one has

$$\sum_r a_r = 0, \qquad \sum_r s_r a_r = 0,$$

so it lies in $\ker B$ and the monomial is invariant. For the anchored family $\mathcal{Q}_{1jk}$, the component at index j equals the same nonzero number $s_1 - s_k$ and is absent from every other anchored exponent vector. Hence these $k-2$ vectors are linearly independent and, by theorem VI.6, complete. □

Corollary VI.10 (Crossover-spacing interpretation of the quotient moduli)**.** *Write the normalized profile as*

$$\varepsilon/\varepsilon_* = \sum_i c_i e^{s_i x}, \qquad x = \ln(A/A_*),$$

and define the pairwise equality location

$$x_{ij} := \frac{\ln c_i - \ln c_j}{s_j - s_i}, \qquad i < j, \tag{VI.58}$$

so that the ith and jth branch contributions are equal at $x = x_{ij}$. Under a pivot change $x_{ij} \mapsto x_{ij} - \beta$, so differences of crossover locations are invariant. Moreover,

$$\boxed{\ln \mathcal{Q}_{ij\ell} = (s_j - s_i)(s_\ell - s_j)\big(x_{ij} - x_{j\ell}\big).} \tag{VI.59}$$

Thus the continuous quotient moduli are exactly relative spacings of branch-crossover locations in logarithmic scale.

Proof. Direct substitution of

$$x_{ij} = \frac{u_i - u_j}{s_j - s_i}$$

gives

$$(s_j - s_i)(s_\ell - s_j)(x_{ij} - x_{j\ell}) = (s_\ell - s_j)u_i + (s_i - s_\ell)u_j + (s_j - s_i)u_\ell,$$

which is $\ln \mathcal{Q}_{ij\ell}$. □

Remark VI.2 (Completeness boundary)**.** Theorem VI.6 exhausts only normalization- and pivot-independent continuous functions of the positive homogeneous coefficient vector on a fixed finite support. It does not classify invariants introduced by a source action, perturbative degrees of freedom, transfer functions, external dimensional scales, additional observables, or an enlarged functional class. Observable nulls below can be new and useful without supplying new quotient moduli: on a fixed support they are functions or consequences of the support data and the complete quotient coordinates.

Remark VI.3 (Local differential realization)**.** The quotient invariants are themselves local finite-jet invariants because

$$\varepsilon_s = e_s(\mathscr{D})\varepsilon.$$

Hence

$$\boxed{\mathcal{Q}_s[\varepsilon] = \frac{[e_{s+1}(\mathscr{D})\varepsilon]\,[e_{s-1}(\mathscr{D})\varepsilon]}{[e_s(\mathscr{D})\varepsilon]^2}, \qquad \mathscr{D} \ln \mathcal{Q}_s = 0.} \tag{VI.60}$$

They therefore test relative branch shape without fixing a dimensional amplitude.

Theorem VI.7 (Complete monomial first integrals of the replicator flow)**.** *Let a positive k-branch mixture obey*

$$\mathscr{D}\pi_s = (s - \kappa_1)\pi_s.$$

For any real vector $a = (a_s)_{s\in S}$ satisfying

$$\sum_s a_s = 0, \qquad \sum_s s\, a_s = 0, \tag{VI.61}$$

the monomial

$$\boxed{I_a := \prod_{s\in S} \pi_s^{a_s}} \tag{VI.62}$$

is constant along the dilation flow. The vector space of exponent vectors a has dimension $k-2$. For consecutive support, one may choose the adjacent second-difference exponent vectors; the corresponding I_a are precisely the quotient cross-ratios of theorem VI.5.

Proof. Taking a logarithmic derivative,

$$\mathscr{D}\ln I_a = \sum_s a_s(s - \kappa_1) = \sum_s s\, a_s - \kappa_1 \sum_s a_s = 0.$$

The two constraints are independent when at least two distinct spectral weights are active, so their common kernel has dimension $k-2$. □

Corollary VI.11 (Pairwise log-odds rigidity)**.** *For any two active branches $s_i \neq s_j$ of a positive mixture,*

$$\boxed{\mathscr{D}\ln\frac{\pi_{s_j}}{\pi_{s_i}} = s_j - s_i, \qquad \mathscr{D}^2\ln\frac{\pi_{s_j}}{\pi_{s_i}} = 0.} \tag{VI.63}$$

Thus every pairwise branch log-odds is an affine function of logarithmic scale with a slope fixed solely by the spectral gap. The intercept carries the relative-amplitude information, while differences of its zero crossings reproduce the quotient moduli of corollary VI.10.

Proof. Subtract the two replicator identities:

$$\mathscr{D}\ln\pi_{s_j} - \mathscr{D}\ln\pi_{s_i} = (s_j - \kappa_1) - (s_i - \kappa_1) = s_j - s_i.$$

A second dilation derivative therefore vanishes. □

Theorem VI.8 (Discrete Koopman–Hankel spectral fingerprint)**.** *Let $x = \ln(A/A_*)$ and choose a nonzero logarithmic step h. For a k-branch profile define*

$$y_n := \varepsilon(A_0 e^{nh}) = \sum_{j=1}^{k} d_j r_j^n, \qquad d_j := C_j A_0^{s_j}, \qquad r_j := e^{s_j h}.$$

Let

$$Y_0^{(n)} := [y_{n+i+j}]_{i,j=0}^{k-1}, \qquad Y_1^{(n)} := [y_{n+i+j+1}]_{i,j=0}^{k-1}. \tag{VI.64}$$

Then

$$\boxed{\frac{\det(\lambda Y_0^{(n)} - Y_1^{(n)})}{\det Y_0^{(n)}} = \prod_{j=1}^{k}(\lambda - r_j).} \tag{VI.65}$$

Thus the generalized eigenvalues recover the active dilation weights without differentiation:

$$\boxed{s_j = \frac{\ln r_j}{h}.} \tag{VI.66}$$

In addition,

$$\boxed{\frac{\det Y_0^{(n+1)}}{\det Y_0^{(n)}} = \prod_{j=1}^{k} r_j = \exp\left(h\sum_{j=1}^{k} s_j\right).} \tag{VI.67}$$

Proof. Let $(V_h)_{ij} = r_j^i$, $R = \operatorname{diag}(r_j)$, and

$$W_n = \operatorname{diag}(d_1 r_1^n, \dots, d_k r_k^n).$$

Then

$$Y_0^{(n)} = V_h W_n V_h^T, \qquad Y_1^{(n)} = V_h W_n R V_h^T.$$

Because $h \neq 0$ and the s_j are distinct real numbers, the r_j are distinct and V_h is invertible. The determinant-pencil identity follows exactly as in theorem VI.1. Since $W_{n+1} = W_n R$,

$$\det Y_0^{(n+1)} / \det Y_0^{(n)} = \det R.$$

□

Remark VI.4 (Koopman form)**.** The logarithmic scale-translation operator

$$(U_h f)(x) := f(x + h)$$

acts on each dilation eigenbranch by

$$U_h e^{sx} = e^{sh} e^{sx}.$$

Hence the valuation space is an exact finite-dimensional invariant subspace of the translation/Koopman operator $U_h = e^{h\mathscr{D}}$. Theorem VI.8 is the sample-space spectral reconstruction of this finite Koopman representation. This is a standard exponential-sum matrix-pencil structure [6–8]; the model-specific content is the restricted valuation spectrum and the resulting exact nulls.

Theorem VI.9 (Global Fisher-information and cumulant sum rules)**.** *Let a positive finite mixture have active extreme weights*

$$s_- := \min S, \qquad s_+ := \max S,$$

and let $x = \ln(A/A_*)$*. Then*

$$\boxed{\int_{-\infty}^{+\infty} \mathcal{I}_x \, \mathrm{d}x = \int_{-\infty}^{+\infty} \kappa_2 \, \mathrm{d}x = s_+ - s_-.} \tag{VI.68}$$

Equivalently, using $\mathcal{I}_x = -3\mathscr{D}w$*,*

$$\boxed{-3 \int_{-\infty}^{+\infty} \mathscr{D}w \, \mathrm{d} \ln A = s_+ - s_-.} \tag{VI.69}$$

More generally,

$$\boxed{\int_{-\infty}^{+\infty} \kappa_n \, \mathrm{d}x = 0, \qquad n \geq 3.} \tag{VI.70}$$

All integrals are absolutely convergent for finite positive support.

Proof. By theorem VI.3,

$$\frac{\mathrm{d}\kappa_n}{\mathrm{d}x} = \kappa_{n+1}.$$

The asymptotic support theorem gives

$$\kappa_1(-\infty) = s_-, \qquad \kappa_1(+\infty) = s_+.$$

Therefore

$$\int_{-\infty}^{+\infty} \kappa_2 \, \mathrm{d}x = \kappa_1(+\infty) - \kappa_1(-\infty) = s_+ - s_-.$$

For every $m \geq 2$, the distribution becomes a point mass at each asymptotic endpoint, so $\kappa_m(\pm\infty) = 0$. Hence

$$\int \kappa_n \, \mathrm{d}x = \kappa_{n-1}(+\infty) - \kappa_{n-1}(-\infty) = 0$$

for $n \geq 3$. Exponential approach to the extreme branch on a finite support gives absolute convergence. □

Theorem VI.10 (Universal two-branch normal form)**.** *Let a positive mixture have exactly two active weights $s_- < s_+$ and put*

$$\Delta := s_+ - s_- > 0, \qquad \bar{s} := \frac{s_- + s_+}{2}.$$

There is a unique crossover location $x_\times$ at which the two branch contributions are equal. If $p_+(x)$ denotes the fraction of the upper branch, then

$$\boxed{p_+(x) = \frac{1}{1 + e^{-\Delta(x - x_\times)}}} \tag{VI.71}$$

and the complete orbit is

$$\boxed{\begin{aligned} \kappa_1(x) &= \bar{s} + \frac{\Delta}{2} \tanh\left[\frac{\Delta}{2}(x - x_\times)\right], \\ \mathcal{I}_x = \kappa_2(x) &= \frac{\Delta^2}{4} \operatorname{sech}^2\left[\frac{\Delta}{2}(x - x_\times)\right], \\ w(x) &= -1 - \frac{\bar{s}}{3} - \frac{\Delta}{6} \tanh\left[\frac{\Delta}{2}(x - x_\times)\right]. \end{aligned}} \tag{VI.72}$$

Thus, after a translation of logarithmic scale, every positive two-branch orbit with fixed support has the same shape; its only remaining parameters are the discrete spectral midpoint and gap.

Proof. If the dimensionless branch coefficients are c_- and c_+, then

$$\frac{\pi_+}{\pi_-} = \frac{c_+}{c_-} e^{\Delta x} = e^{\Delta(x - x_\times)}, \qquad x_\times = \frac{1}{\Delta} \ln \frac{c_-}{c_+}.$$

Hence $p_+ = \pi_+$ is logistic. Since $\kappa_1 = s_- + \Delta p_+$ and $\kappa_2 = \Delta^2 p_+(1 - p_+)$, the hyperbolic forms follow, and $w = -1 - \kappa_1/3$ gives the last identity. □

Corollary VI.12 (Universal Fisher length of a positive two-branch orbit)**.** *For every positive two-branch mixture,*

$$\boxed{L_F := \int_{-\infty}^{+\infty} \sqrt{\mathcal{I}_x}\, \mathrm{d}x = \pi.} \tag{VI.73}$$

The result is independent of the two positive amplitudes, their ratio, the scale pivot, and the spectral gap. The integrated Fisher information is instead the spectral width,

$$\int_{-\infty}^{+\infty} \mathcal{I}_x\, \mathrm{d}x = \Delta. \tag{VI.74}$$

Proof. By theorem VI.10,

$$\sqrt{\mathcal{I}_x} = \frac{\Delta}{2} \operatorname{sech}\left[\frac{\Delta}{2}(x - x_\times)\right].$$

Its integral over the complete real line is π. The information integral is the $n = 2$ case of theorem VI.9. □

Corollary VI.13 (Schwarzian spectral-gap invariant for two branches)**.** *Where $\mathscr{D}w \neq 0$, define the Schwarzian derivative with respect to $x = \ln(A/A_*)$ by*

$$\{w, x\} := \frac{\mathscr{D}^3 w}{\mathscr{D} w} - \frac{3}{2}\left(\frac{\mathscr{D}^2 w}{\mathscr{D} w}\right)^2. \tag{VI.75}$$

Every positive two-branch orbit satisfies the local amplitude- and pivot-independent identity

$$\boxed{\{w, x\} = -\frac{(s_+ - s_-)^2}{2}.} \tag{VI.76}$$

Thus the spectral gap can be recovered locally from the projective differential invariant

$$\boxed{\Delta = \sqrt{-2\{w, x\}}.} \tag{VI.77}$$

Proof. The equation-of-state profile in theorem VI.10 is an affine function of $\tanh[\Delta(x - x_\times)/2]$. The Schwarzian is unchanged by an affine transformation of its dependent variable, and direct differentiation gives

$$\{\tanh(ax), x\} = -2a^2.$$

Taking $a = \Delta/2$ proves the result. □

Corollary VI.14 (Exact positive $\Lambda + X$ background normal form)**.** *Consider the positive two-branch dark-sector support*

$$S_{\Lambda X} = \{-1, 0\},$$

with $s = -1$ identified with X and $s = 0$ with vacuum. Setting $x = \ln a$, the complete homogeneous dark-sector equation of state is

$$\boxed{w_D(x) = -\frac{5}{6} - \frac{1}{6}\tanh\left(\frac{x - x_\times}{2}\right).} \tag{VI.78}$$

Equivalently it obeys the first-order amplitude-free null

$$\boxed{\mathscr{D}w_D = (1 + w_D)(2 + 3w_D), \qquad -1 < w_D < -\frac{2}{3}.} \tag{VI.79}$$

The information profile and its exact extrema are

$$\boxed{\mathcal{I}_x = -3\mathscr{D}w_D = \frac{1}{4}\operatorname{sech}^2\left(\frac{x - x_\times}{2}\right), \qquad \max(-\mathscr{D}w_D) = \frac{1}{12}.} \tag{VI.80}$$

Moreover,

$$\boxed{\int_{-\infty}^{+\infty} \mathcal{I}_x \, \mathrm{d}x = 1, \qquad L_F = \pi, \qquad \{w_D, x\} = -\frac{1}{2}.} \tag{VI.81}$$

If the present epoch $a = 1$ is chosen as the physical amplitude pivot and the dark sector contains only these two branches, then the relative present amplitude is reconstructed from $w_D(a)$ by

$$\boxed{\frac{\Omega_X}{\Omega_\Lambda} = a\,\frac{3(1 + w_D)}{-2 - 3w_D}.} \tag{VI.82}$$

The last identity is pivot-fixed rather than a pivot-free quotient invariant; it therefore does not contradict the amplitude no-go theorem.

Proof. Apply theorem VI.10 with $s_- = -1$, $s_+ = 0$, hence $\Delta = 1$ and $\bar{s} = -1/2$. Since $\kappa_1 = -\pi_X = -3(1 + w_D)$, one has

$$\pi_X = 3(1 + w_D), \qquad \pi_\Lambda = -2 - 3w_D.$$

Their ratio is

$$\frac{\pi_X}{\pi_\Lambda} = \frac{\Omega_X a^{-1}}{\Omega_\Lambda},$$

which gives equation (VI.82). The remaining identities follow directly from the two-branch normal form. □

Theorem VI.11 (Fisher-length rigidity and exact branch-excess decomposition)**.** *Let a positive finite mixture have at least two distinct active weights and extreme support values*

$$m = s_- < s_+ = M.$$

Then the Fisher length of the complete logarithmic-scale orbit satisfies

$$\boxed{L_F := \int_{-\infty}^{+\infty} \sqrt{\mathcal{I}_x}\,\mathrm{d}x \geq \pi.} \tag{VI.83}$$

Equality holds if and only if the active support is contained in the two endpoints $\{m, M\}$. Hence every genuinely multi-branch positive orbit obeys

$$\boxed{\Delta_F := L_F - \pi > 0.} \tag{VI.84}$$

More sharply, let $\mu = \kappa_1$ and define

$$B(\mu) := (M-\mu)(\mu-m), \qquad V(\mu) := \kappa_2, \qquad \Delta_{mM}(\mu) := B(\mu) - V(\mu).$$

Then

$$\boxed{\Delta_F = \int_m^M \frac{\Delta_{mM}(\mu)\,\mathrm{d}\mu}{\sqrt{B(\mu)V(\mu)}\,[\sqrt{B(\mu)}+\sqrt{V(\mu)}]}.} \tag{VI.85}$$

The integral is understood as the convergent improper endpoint limit. Thus the global Fisher excess is exactly the accumulated Bhatia–Davis slack. For fixed support it is a function of the complete quotient moduli of theorem VI.6; it introduces no additional independent coefficient modulus.

Proof. For a nontrivial positive finite mixture, $\mu = \kappa_1$ increases strictly from m to M and

$$\frac{\mathrm{d}\mu}{\mathrm{d}x} = \kappa_2 = V(\mu) > 0.$$

Therefore

$$L_F = \int_m^M \frac{\mathrm{d}\mu}{\sqrt{V(\mu)}}.$$

The Bhatia–Davis bound gives $V(\mu) \leq B(\mu)$, so

$$L_F \geq \int_m^M \frac{\mathrm{d}\mu}{\sqrt{(M-\mu)(\mu-m)}} = \pi.$$

By theorem V.3, equality in the variance bound at every finite scale occurs if and only if all active probability is carried by m and M. If any interior coefficient is positive, the slack is strictly positive for every finite x, hence $L_F > \pi$. Finally,

$$\frac{1}{\sqrt{V}} - \frac{1}{\sqrt{B}} = \frac{B-V}{\sqrt{BV}(\sqrt{B}+\sqrt{V})},$$

and integration yields equation (VI.85). □

Remark VI.5 (Prior-art and novelty boundary)**.** The underlying tools are not claimed as new in isolation. Hadwiger theory and intrinsic volumes are classical; Minkowski functionals have long been used in cosmology as morphological statistics [10]; $w = -2/3$ occurs in domain-wall cosmology [11]; quasilocal energy has been studied directly in FLRW spacetimes [12, 13]; and other cosmological dilation algebras exist, for example the CVH algebra in FLRW/LQC [14]. Likewise, exponential-family geometry, Prony reconstruction, and matrix-pencil recovery are established structures [5–8].

The model-specific result proved here is the exact chain from the declared Euclidean convex-valuation closure on the round B^4/S^3 family to the five-weight cosmological dilation spectrum, its semisimple projector algebra, complete positive-support quotient moduli, Hankel/inertia fingerprints, observable jet cone, Fisher-rigidity identities, and boundary expansion quotient. A targeted prior-art search found no direct predecessor deriving this complete chain. Accordingly, any priority statement is scoped to this linked construction; no priority is claimed for Hadwiger theory, $w = -2/3$, quasilocal energy, Prony/Hankel methods, Fisher geometry, or cosmological dilation operators separately.

VII. BOUNDARY FACE AND EXPANSION QUOTIENT

The full valuation spectrum contains the bulk branch $s = 1$, coming from V_4. The boundary curvature-measure terms give the remaining four branches.

Criterion VII.1 (Boundary-local valuation class)**.** The local boundary class used here consists of filling-relative scalar densities built from the induced boundary measure and the elementary symmetric curvatures $e_j(K)$, with coefficients independent of A. It excludes boundary integrals involving the support function $x \cdot n$, ambient-position data, or other nonlocal information.

Proposition VII.1 (Boundary face)**.** *Inside this declared boundary-local class,*

$$S_\partial = \{0, -1, -2, -3\}. \tag{VII.1}$$

Equivalently, a full valuation profile lies in the boundary face if and only if

$$\mathsf{P}_1 \varepsilon = 0, \tag{VII.2}$$

or

$$\mathscr{D}(\mathscr{D}+1)(\mathscr{D}+2)(\mathscr{D}+3)\varepsilon = 0. \tag{VII.3}$$

Proof. The bulk term has energy scaling A^4 and density scaling A, hence $s = 1$. The boundary curvature-measure terms have energy powers A^3, A^2, A, A^0, hence density powers $A^0, A^{-1}, A^{-2}, A^{-3}$. The declared boundary-local class excludes the bulk intrinsic volume term. The projector and annihilator statements are the spectral algebra already proved. □

Remark VII.1 (Conditional nature of the filter)**.** The exclusion of the bulk branch is not a topological theorem. It is conditional on the stated boundary-local class. For example, the Minkowski formula can express V_4 through a boundary integral involving $x \cdot n$; that ambient-position datum is deliberately excluded here.

Let $a = A/A_0$. At the level of the normalized expansion function $\mathcal{H}^2(a) = H^2(a)/H_0^2$, the boundary powers are

$$1, \quad a^{-1}, \quad a^{-2}, \quad a^{-3}. \tag{VII.4}$$

The standard curved-ΛCDM nuisance powers are

$$1, \quad a^{-2}, \quad a^{-3}. \tag{VII.5}$$

Theorem VII.1 (Expansion-rate quotient)**.** *At the level of $\mathcal{H}^2(a)$ alone,*

$$\mathcal{V}_\partial / \operatorname{span}\{1, a^{-2}, a^{-3}\} \simeq \operatorname{span}\{[a^{-1}]\}. \tag{VII.6}$$

The unique nonstandard quotient direction has

$$w = -\frac{2}{3}. \tag{VII.7}$$

Proof. The functions $1, a^{-1}, a^{-2}, a^{-3}$ are linearly independent on every interval $a > 0$. Quotienting by the span of $1, a^{-2}, a^{-3}$ leaves the single class $[a^{-1}]$. The equation-of-state map gives $w = -1 - (-1)/3 = -2/3$. □

VIII. GEOMETRIC CURVATURE SPLIT

The quotient theorem is only an expansion-rate statement. A physical stress component with a^{-2} scaling is not the same object as geometric spatial curvature.

Let

$$\Omega_K^{\mathrm{geom}} := -\frac{c^2}{A_0^2 H_0^2} \tag{VIII.1}$$

for the closed FLRW convention. A physical $s = -2$ valuation component is denoted $\Omega_{2,\mathrm{val}}$. The boundary-profile background takes the form

$$\mathcal{H}^2(a) = \Omega_r a^{-4} + \Omega_m a^{-3} + \big(\Omega_K^{\mathrm{geom}} + \Omega_{2,\mathrm{val}}\big) a^{-2} + \Omega_X a^{-1} + \Omega_\Lambda. \tag{VIII.2}$$

The geometric curvature sector, together with the assumed S^3 topology, fixes the curvature scale entering the FLRW distance relation and compact spatial harmonic spectrum. A physical a^{-2} stress component affects the expansion history but is not thereby geometric curvature.

Define

$$d(z) := \frac{H_0 D_M(z)}{c}. \tag{VIII.3}$$

For an FLRW geometry, the Clarkson–Bassett–Lu consistency relation gives

$$\Omega_K^{\rm geom} = \frac{[\mathcal{H}(z)d'(z)]^2 - 1}{d(z)^2}, \tag{VIII.4}$$

with the value at $z = 0$ understood by the regular limit [4].

The finite cosmological power alphabet admits exact coefficient projectors.

Theorem VIII.1 (Exact expansion-coefficient projectors)**.** *Let*

$$S_{\cos} := \{0, -1, -2, -3, -4\}, \qquad F(a) = \sum_{s \in S_{\cos}} \Omega_s a^s. \tag{VIII.5}$$

For every $s \in S_{\cos}$, define

$$\boxed{\mathcal{C}_s[F](a) := a^{-s} \prod_{\substack{t \in S_{\cos} \\ t \neq s}} \frac{\mathscr{D} - t}{s - t} F(a).} \tag{VIII.6}$$

Then

$$\boxed{\mathcal{C}_s[F](a) = \Omega_s} \tag{VIII.7}$$

for every $a > 0$. Hence $\mathscr{D}\mathcal{C}_s[F] = 0$ is an exact coefficient-level null inside this five-power expansion alphabet.

Proof. The Lagrange polynomial in $\mathscr{D}$ annihilates every branch a^t with $t \neq s$ and acts as the identity on a^s. Multiplication by a^{-s} removes the remaining scale factor. □

The two projectors needed below are

$$\mathcal{C}_{-1}[F] = -\frac{a}{6}\mathscr{D}(\mathscr{D}+2)(\mathscr{D}+3)(\mathscr{D}+4)F, \tag{VIII.8}$$

$$\mathcal{C}_{-2}[F] = \frac{a^2}{4}\mathscr{D}(\mathscr{D}+1)(\mathscr{D}+3)(\mathscr{D}+4)F. \tag{VIII.9}$$

Thus $\widehat{\Omega}_X = \mathcal{C}_{-1}[F]$, while the physical $s = -2$ coefficient is obtained only after subtracting geometric curvature,

$$\boxed{\Omega_{2,\rm val} = \mathcal{C}_{-2}[F] - \Omega_K^{\rm geom}.} \tag{VIII.10}$$

Corollary VIII.1 (Physical parameter count)**.** *Relative to curved ΛCDM, the general boundary-profile extension at background-geometry level is two-parameter:*

$$(\Omega_X, \Omega_{2,\rm val}). \tag{VIII.11}$$

The one-parameter VDE subclass is the nested restriction

$$\Omega_{2,\rm val} = 0. \tag{VIII.12}$$

IX. CANONICAL SOURCE LIFTS AND DUST CURRENT

The valuation theorem fixes homogeneous powers. It does not fix a unique covariant stress tensor. In this section only we use $c = \hbar = 1$ and $M_{\rm Pl}^{-2} = 8\pi G$. For a branch $\rho_s(a) = \rho_{s0} a^s$, define

$$\Omega_s := \frac{\rho_{s0}}{3M_{\rm Pl}^2 H_0^2}. \tag{IX.1}$$

Theorem IX.1 (Canonical scalar realization for $s < 0$)**.** *Let*

$$\rho_s(a) = \rho_{s0} a^s > 0, \qquad s < 0, \qquad w_s = -1 - \frac{s}{3}. \tag{IX.2}$$

A minimally coupled canonical scalar realizes this homogeneous branch through

$$\dot\phi_s^2 = -\frac{s}{3}\rho_s, \qquad V_s = \frac{6+s}{6}\rho_s. \tag{IX.3}$$

Equivalently, with $M_{\rm Pl}^{-2} = 8\pi G$,

$$\frac{{\rm d}\phi_s}{{\rm d}\ln a} = \pm M_{\rm Pl}\frac{\sqrt{-s\,\Omega_s a^s}}{\mathcal{H}(a)}. \tag{IX.4}$$

The realization has

$$c_{s,\rm rest}^2 = 1, \qquad \pi_{\mu\nu} = 0. \tag{IX.5}$$

Proof. For a canonical scalar,

$$\rho = \frac{1}{2}\dot\phi^2 + V, \qquad p = \frac{1}{2}\dot\phi^2 - V.$$

Hence $\dot\phi^2 = (1+w)\rho$ and $V = (1-w)\rho/2$. Substituting $w_s = -1 - s/3$ gives the displayed formulas. □

Theorem IX.2 (Canonical scalar mass-flow invariant)**.** *For the same branch,*

$$\boxed{V_{s,\phi\phi} = -\frac{6+s}{4}H^2\left[s + \mathscr{D}\ln(\mathcal{H}^2/\mathcal{H}_*^2)\right],} \tag{IX.6}$$

where $\mathcal{H}_* > 0$ is an arbitrary normalization.

Proof. Write $F = \mathcal{H}^2 = H^2/H_0^2$ and

$$V_s(a) = \frac{6+s}{2}M_{\rm Pl}^2 H_0^2 \Omega_s a^s, \qquad \mathscr{D}\phi_s = \pm M_{\rm Pl}\sqrt{\frac{-s\Omega_s a^s}{F}}.$$

Using ${\rm d}/{\rm d}\phi = (\mathscr{D}\phi)^{-1}\mathscr{D}$, differentiating twice, and simplifying gives

$$V_{s,\phi\phi} = -\frac{6+s}{4}H_0^2 F\left[s + \mathscr{D}\ln F\right].$$

This is the stated expression. □

For the two non-dust boundary branches,

$$s = -1: \qquad m_X^2 = \frac{5}{4}H^2\left[1 - \mathscr{D}\ln\mathcal{H}^2\right], \tag{IX.7}$$

$$s = -2: \qquad m_2^2 = H^2\left[2 - \mathscr{D}\ln\mathcal{H}^2\right]. \tag{IX.8}$$

Equivalently,

$$m_X^2/H_0^2 = \frac{5}{4}(F - \mathscr{D}F), \qquad m_2^2/H_0^2 = 2F - \mathscr{D}F. \tag{IX.9}$$

The $s = -3$ branch can be reproduced homogeneously by a canonical scalar, but that scalar has $c_s^2 = 1$ and is not a cold-dust completion. A CDM-like source requires a conserved pressureless current.

Theorem IX.3 (Pressureless current realization)**.** *Let $J^\mu = nu^\mu$ be a conserved timelike particle-number current,*

$$\nabla_\mu J^\mu = 0, \qquad u^\mu u_\mu = -1. \tag{IX.10}$$

The pressureless stress tensor

$$T_{\mu\nu} = \rho u_\mu u_\nu, \qquad \rho = mn, \tag{IX.11}$$

satisfies

$$p = 0, \qquad c^2_{s,\mathrm{rest}} = 0, \qquad \pi_{\mu\nu} = 0, \tag{IX.12}$$

and, on a homogeneous FLRW background,

$$\rho \propto a^{-3}. \tag{IX.13}$$

Proof. Conservation of J^μ gives $n \propto V_3^{-1} \propto a^{-3}$ in a comoving homogeneous flow. The standard pressureless perfect-fluid stress tensor has $p = 0$, no rest-frame pressure perturbation, and no anisotropic stress. □

Remark IX.1 (Source layer)**.** The scalar and current constructions are existence lifts of selected homogeneous branches. They are not derived by covariant variation of E_4^{val}. A different lift with the same background power can have different perturbations.

Proposition IX.1 (Conditional subhorizon growth consistency for a smooth lift)**.** *Let*

$$F(a) := \mathcal{H}^2(a) = H^2(a)/H_0^2$$

be a background expansion history already fixed independently. Assume general relativity, pressureless clustering matter, negligible radiation and scale-dependent massive-neutrino corrections over the redshift and wavenumber range of interest, and an additional component whose perturbations are negligible on the considered scales. Then the linear growing-mode matter contrast $D(a)$ satisfies the standard subhorizon equation

$$\boxed{\mathcal{R}_{\mathrm{growth}}[D; F] := \mathscr{D}^2 D + \left(2 + \frac{1}{2}\mathscr{D}\ln F\right)\mathscr{D}D - \frac{3}{2}\Omega_m(a)D = 0,} \tag{IX.14}$$

where

$$\Omega_m(a) = \frac{\Omega_{m0}a^{-3}}{F(a)}. \tag{IX.15}$$

For the canonical-scalar lifts above, $c^2_{s,\mathrm{rest}} = 1$ makes the smooth-component approximation appropriate only on scales well inside the scalar sound horizon and where its perturbations are dynamically negligible. Equation (IX.14) is therefore a conditional lift-level consistency relation, not an exact consequence of the valuation theorem and not a substitute for a full Boltzmann transfer calculation [9].

With

$$f(a) := \mathscr{D}\ln D,$$

a pivot-normalized prediction for redshift-space growth is

$$f\sigma_8(a) = f(a)\,\sigma_{8,0}\frac{D(a)}{D(1)}. \tag{IX.16}$$

Thus a declared source lift plus the background projectors closes a reproducible approximate growth null, while disagreement with it falsifies that lift rather than the finite valuation algebra itself.

X. ENDPOINT NORMAL FORM AND AMPLITUDE NO-GO

This section records a restricted endpoint normal form. Its hypotheses are intentionally narrow.

Hypothesis X.1 (Minimal endpoint class)**.** On a connected oriented $\Sigma \simeq S^3$, assume:

1. local metric dependence is zeroth order and $\mathrm{Diff}^+(\Sigma)$-natural;
2. there are no local endpoint degrees of freedom;
3. curvature, derivatives, embedding-position data, background connections, and nonlocal spectral functionals are excluded;
4. additivity holds on finite disjoint unions;
5. global internal data enter only through a prescribed finite-rank additive charge vector $Q \in \mathbb{Z}^r$, independent of the metric scale.

Theorem X.1 (Conditional endpoint normal form)**.** *Inside the minimal endpoint class,*

$$E_{\mathrm{end}} = \sigma\ \mathrm{Vol}(\Sigma) + \sum_{A=1}^{r} \mu_A Q_A. \tag{X.1}$$

For a single oriented degree Q_U,

$$E_{\mathrm{end}} = \sigma V_3 + \mu Q_U, \tag{X.2}$$

and hence

$$\varepsilon(a) = \sigma + \frac{\mu Q_U}{2\pi^2 A_0^3} a^{-3}, \qquad p = -\sigma. \tag{X.3}$$

Proof. Under the stated exclusions, the only local zeroth-order natural metric density is a universal constant multiple of the Riemannian volume density. The remaining metric-independent charge dependence is an additive homomorphism $f : \mathbb{Z}^r \to \mathbb{R}$, hence $f(Q) = \sum_A \mu_A Q_A$. □

No-go theorem X.1 (Scale-covariance no-go for dimensional amplitudes)**.** Purely topological or dimensionless data cannot naturally determine a nonzero quantity of nonzero length dimension while remaining covariant under global metric rescalings. In particular, topology alone cannot determine nonzero values of σ, μ, H_0, branch densities, mass scales, or cutoffs.

Proof. Under $L \mapsto \lambda L$, topological and dimensionless data are unchanged. A quantity X with length dimension L^k, $k \neq 0$, must transform as $X \mapsto \lambda^k X$. If X is determined only by the unchanged data, it must also remain unchanged. Compatibility for all $\lambda > 0$ implies $X = 0$. □

Remark X.1 (Dimensionless parameters)**.** Dimensionless density parameters such as Ω_X are not fixed by this dimensional argument alone. They become meaningful only after a normalization scale, critical density, or amplitude ledger is declared. They are then constrained by matching or data, not by topology alone.

XI. CLOSURE TESTS

At the density-profile level, the exact closure residuals are

$$\mathcal{R}_{\mathrm{val}}[\varepsilon] := (\mathscr{D}-1)\mathscr{D}(\mathscr{D}+1)(\mathscr{D}+2)(\mathscr{D}+3)\varepsilon, \tag{XI.1}$$
$$\mathcal{R}_{\partial}[\varepsilon] := \mathscr{D}(\mathscr{D}+1)(\mathscr{D}+2)(\mathscr{D}+3)\varepsilon, \tag{XI.2}$$
$$\mathcal{R}_{\mathrm{end}}[\varepsilon] := \mathscr{D}(\mathscr{D}+3)\varepsilon. \tag{XI.3}$$

The full valuation, boundary valuation, and endpoint classes satisfy respectively

$$\mathcal{R}_{\mathrm{val}} = 0, \qquad \mathcal{R}_{\partial} = 0, \qquad \mathcal{R}_{\mathrm{end}} = 0. \tag{XI.4}$$

For the nested a^{-1} expansion-rate direction, use the exact coefficient projector $\widehat{\Omega}_X(a) := \mathcal{C}_{-1}[\mathcal{H}^2](a)$ of theorem VIII.1. The exact one-parameter VDE background requires $\widehat{\Omega}_X(a)$ to be constant. Curved ΛCDM is the sharper null

$$\widehat{\Omega}_X(a) = 0. \tag{XI.5}$$

Criterion XI.1 (Practical estimation)**.** High-order numerical differentiation is noise-sensitive. Practical closure tests should use covariance-aware basis fits, direct likelihood parameterizations, Gaussian-process reconstructions with derivative covariance, or discrete Prony recurrences on logarithmically spaced samples. The analytic differential operators define the invariant nulls; they need not be evaluated by raw finite differences.

Criterion XI.2 (Free-spectrum identification protocol)**.** A reconstruction in which the generalized eigenvalues are constrained from the outset to

$$\{e^h, 1, e^{-h}, e^{-2h}, e^{-3h}\}$$

can test consistency with the valuation closure but cannot independently identify those spectral locations. For identification, the sample-Hankel or exponential-sum analysis shall first infer the active generalized eigenvalues r_j as free quantities, with the full data covariance and model-order uncertainty propagated, and only afterwards test whether

$$s_j = \frac{\ln r_j}{h}$$

are statistically compatible with a subset of S_{val}. This free-spectrum step is the operational distinction between a closure null and an independent spectral measurement.

For logarithmically equally spaced samples $x_n = \ln a_n = x_0 + nh$, the discrete counterpart is

$$\prod_{s\in S}(L - e^{sh})f_n = 0, \tag{XI.6}$$

where $Lf_n = f_{n+1}$. This is the Prony form of the same finite spectral closure. A stronger derivative-free implementation is the sample Hankel pencil of theorem VI.8: its generalized eigenvalues must lie in

$$\{e^h, 1, e^{-h}, e^{-2h}, e^{-3h}\} \tag{XI.7}$$

for the full valuation class, with the appropriate subset for a lower-dimensional support. The determinant-step ratio must simultaneously equal e^{-5h}, e^{-6h}, or e^{-3h} for the fully active valuation, boundary, or endpoint supports, respectively.

XII. EXECUTED NUMERICAL INVARIANT AUDIT

This section is deliberately downstream of the theorem layer. It evaluates the exact identities on supplied execution artifacts and separates three logically distinct questions: whether the declared source lift was implemented correctly, whether the invariant identities recover the instantiated background, and whether external data independently identify or significantly prefer the predicted branch.

A. Execution identity and scope

The transfer evidence uses unmodified CLASS/`classy` v3.3.4 at source commit

`e85808324f51fc694d12e3ed7439552a3c3f9540`, (XII.1)

with closed FLRW geometry $\Omega_K^{\mathrm{geom}} = -0.01$. CLASS is a public Einstein–Boltzmann code [15, 16]. The nested source lift is represented by the CLASS fluid sector with

$$\Omega_X \mapsto \texttt{Omega_fld}, \qquad w_X = -\frac{2}{3}, \qquad c^2_{s,\mathrm{rest}} = 1, \qquad \texttt{use_ppf=no}. \tag{XII.2}$$

The evidence bundle contains scalar CMB spectra through $\ell = 1200$, linear $P(k, z = 0)$, and scalar transfer outputs. The BAO pilot uses Cobaya 3.6.2 [17] with the official all-tracer DESI DR2 BAO likelihood [18], 13 BAO observables, fixed $\Omega_K^{\mathrm{geom}} = -0.01$ and $\omega_b = 0.02237$, and profiled H_0 and ω_{cdm}. No CMB, supernova, lensing, or growth likelihood is included in this pilot.

B. Executed transfer checks

The latest supplied validation gate reports 26/26 checks passed. The first physical closed scalar mode is expected at

$$9.434617347 \times 10^{-5}\ h\,\mathrm{Mpc}^{-1} \tag{XII.3}$$

and the transfer table begins at

$$9.43455838046 \times 10^{-5}\ h\,\mathrm{Mpc}^{-1}, \tag{XII.4}$$

a relative difference of 6.25002×10^{-6}. The executed background law $\rho_X \propto a^{-1}$ has maximum reported residual

$$\boxed{1.88064 \times 10^{-9}}. \tag{XII.5}$$

The required transfer columns `d_fld`, `t_fld`, `phi`, `psi`, and `k (h/Mpc)` are present.

A closed-safe default-versus-tight precision comparison gives the 95th-percentile relative differences

	TT	EE	$P(k)$
VDE	1.026×10^{-3}	1.098×10^{-3}	4.865×10^{-4}
control	1.003×10^{-3}	1.091×10^{-3}	4.560×10^{-4}

At $\Omega_X = 10^{-8}$ relative to a 10^{-12} anchor, the latest common-precision null comparison gives

$$TT_{\mathrm{p95}} = 3.18 \times 10^{-6}, \qquad EE_{\mathrm{p95}} = 6.30 \times 10^{-6}, \qquad P(k)_{\mathrm{p95}} = 9.22 \times 10^{-8}. \tag{XII.6}$$

For the fiducial $\Omega_X = 0.05$ run,

$$\sigma_{8,\mathrm{control}} = 0.831044, \qquad \sigma_{8,\mathrm{VDE}} = 0.822593, \tag{XII.7}$$

a -1.02% shift. These are transfer consequences of the declared source lift, not likelihood evidence for the valuation hypothesis.

C. Exact projectors, support recovery, and curvature inertia

For invariant arithmetic we use the supplied deterministic best-fit ledger

$$H_0 = 66.3703\ \mathrm{km\,s^{-1}\,Mpc^{-1}}, \quad \Omega_K^{\mathrm{geom}} = -0.01, \quad \Omega_X = 0.174541, \quad \Omega_\Lambda = 0.541216, \tag{XII.8}$$

with matter and radiation completed by the accompanying machine-readable background ledger. The projectors of theorem VIII.1 return

$$\boxed{\mathcal{C}_{-1}[F] = 0.174541}, \qquad \boxed{\mathcal{C}_{-2}[F] = -0.010000} \tag{XII.9}$$

throughout the tested scale-factor grid. Hence

$$\boxed{\Omega_{2,\mathrm{val}} = \mathcal{C}_{-2}[F] - \Omega_K^{\mathrm{geom}} = 0}, \tag{XII.10}$$

which is exactly the nested one-parameter restriction used in the execution.

At $a = 1$, the two-state dark-sector derivative pencil for

$$\varepsilon_D = \Omega_\Lambda + \Omega_X a^{-1} \tag{XII.11}$$

recovers the generalized eigenvalues

$$\boxed{\{-1, 0\}}, \tag{XII.12}$$

with positive inertia $(2, 0, 0)$. After geometric curvature is removed, the nonradiative physical ledger $\{\Lambda, X, m\}$ has support $\{-3, -1, 0\}$, inertia $(3, 0, 0)$, and positive Hankel determinant. If the negative closed-FLRW geometric coefficient is instead inserted into the same physical moment measure, the pencil recovers $\{-3, -2, -1, 0\}$ with

$$\left(\mathrm{tr}\,K, \mathrm{tr}\,K^2, \mathrm{tr}\,K^3, \mathrm{tr}\,K^4\right) = (-6, 14, -36, 98) \tag{XII.13}$$

and

$$\boxed{(\nu_+, \nu_-, \nu_0) = (3, 1, 0)}. \tag{XII.14}$$

The unique negative direction is exactly the coefficient $\Omega_K^{\mathrm{geom}} = -0.01$. This numerically demonstrates why geometric curvature must be removed before Hankel positivity is interpreted as positivity of physical branch energies.

By criteria I.1 and XI.2, these projector and pencil results are implementation-consistency tests. They do not independently discover $s = -1$, because $w_X = -2/3$ fixes that exponent before the calculation.

D. Information-geometric invariants of the supplied $\Lambda + X$ ledger

The exact two-branch normal form of corollary VI.14 gives at $a = 1$

$$\pi_X = 0.243855, \qquad \pi_\Lambda = 0.756145, \tag{XII.15}$$

and therefore

$$\boxed{w_D = -0.918715, \qquad \mathcal{I}_x = 0.184390, \qquad w_a = -\mathscr{D} w_D = 0.0614633.} \tag{XII.16}$$

The equal-weight point occurs at

$$a_\times = 0.322498, \qquad \boxed{z_\times = 2.10080}, \tag{XII.17}$$

where $w_D = -5/6$, $\mathcal{I}_x = 1/4$, and $-\mathscr{D} w_D = 1/12$. Because the instantiated support is exactly $\{-1, 0\}$, its complete-orbit invariants are

$$\boxed{L_F = \pi, \qquad \int_{-\infty}^{+\infty} \mathcal{I}_x \, \mathrm{d}x = 1, \qquad \{w_D, x\} = -\frac{1}{2}.} \tag{XII.18}$$

Relative to the full positive boundary endpoints $\{-3, 0\}$, the local Bhatia–Davis slack at the present epoch is

$$\Delta_\partial = 2\pi_X \simeq 0.487710, \tag{XII.19}$$

correctly identifying the $s = -1$ branch as interior to that larger boundary spectrum. These are derived invariants of the supplied ledger, not independently fitted CPL or Schwarzian observables.

E. Finite $s = 0, -1$ residual budget

The finite present-day energy density carried by the two dark slots is

$$\rho_{\Lambda X,0} = \frac{3H_0^2 c^2}{8\pi G}(\Omega_\Lambda + \Omega_X) = \frac{3H_0^2 c^2}{8\pi G}\left(1 - \Omega_K^{\mathrm{geom}} - \Omega_{2,\mathrm{val}} - \Omega_m - \Omega_r\right). \tag{XII.20}$$

For the supplied deterministic ledger,

$$\Omega_\Lambda + \Omega_X = 0.715757, \tag{XII.21}$$

which gives

$$\boxed{\rho_{\Lambda X,0} = 5.322665232877199 \times 10^{-10}\ \mathrm{J\,m^{-3}},} \tag{XII.22}$$

and the natural-unit fourth-root scale

$$\boxed{E_{\Lambda X} := \left[\rho_{\Lambda X,0}(\hbar c)^3\right]^{1/4} = 2.247732070576682\ \mathrm{meV}.} \tag{XII.23}$$

This is deterministic background closure arithmetic. It is not the bare ultraviolet sum $\frac{1}{2}\sum \hbar\omega$, not a topology-only amplitude selector, and not a vacuum-energy cancellation mechanism.

F. DESI DR2 BAO-only amplitude test

The latest supplied profile-likelihood summary gives

branch	Ω_X	H_0	$\omega_{\rm cdm}$	χ^2
alternative	0.174541	66.370288	0.107202	9.271990
null	0	69.237631	0.121014	10.636129

so that

$$q_X = 1.364139, \qquad \sqrt{q_X} = 1.167964, \qquad \boxed{p_{\text{one-sided}} = 0.121411}. \tag{XII.24}$$

With the one-parameter boundary calibration, this is only a 1.17σ-level BAO-only fluctuation and therefore not a detection. The accompanying positive-prior MCMC slice has posterior mean $\Omega_X \simeq 0.1521$, but because the prior enforces $\Omega_X \geq 0$, that posterior by itself is not evidence that the boundary amplitude is nonzero.

The numerical hierarchy is therefore

$$\boxed{\begin{array}{c}\text{valuation theorem} \Longrightarrow \text{exact invariant predictions},\\ \text{fixed } s=-1 \text{ source lift} \Longrightarrow \text{executed CLASS consistency},\\ \text{DESI DR2 BAO-only profile} \Longrightarrow \text{no significant amplitude detection.}\end{array}} \tag{XII.25}$$

The next genuinely independent test is the covariance-aware free-spectrum reconstruction of criterion XI.2, in which the dilation eigenvalues themselves are inferred before comparison with $S_{\rm val}$.

XIII. CONCLUSION

The universal homogeneous work map is exact but not predictive: its inverse reconstructs a filling representative for any differentiable homogeneous residual profile. The restriction enters only after the Euclidean convex-valuation hypothesis is imposed. Under that hypothesis the round homothetic profile is forced into

$$\mathcal{V}_{\rm val} = \operatorname{span}\{A, 1, A^{-1}, A^{-2}, A^{-3}\}, \tag{XIII.1}$$

with

$$P_{\rm val}(\mathscr{D})\varepsilon = 0, \qquad P_{\rm val}(z) = (z-1)z(z+1)(z+2)(z+3), \tag{XIII.2}$$

and the polynomial dilation action is the semisimple algebra

$$\mathfrak{A}_{\mathscr{D}} \simeq \mathbb{R}[z]/(P_{\rm val}) \simeq \mathbb{R}^5. \tag{XIII.3}$$

The finite spectrum supports a complete homogeneous invariant calculus. Primitive idempotents isolate branches; annihilator ideals and derivative or sampled Hankel pencils recover support; Hankel inertia records amplitude signs; normalized jets give exact barycentric positivity certificates; and on every positive fixed-support stratum the normalization/pivot quotient has exactly $k-2$ continuous moduli, represented by monomial cross-ratios. In the positive cone the branch fractions form an exponential family with $\mathscr{D}^2 \ln \varepsilon = \operatorname{Var}_\pi(s)$ and $\mathscr{D}w = -\operatorname{Var}_\pi(s)/3$. The Bhatia–Davis remainder is an exact local interior-branch detector, the observable $(w, \mathscr{D}w, \mathscr{D}^2 w, \mathscr{D}^3 w)$ jet is characterized by five explicit simplex inequalities, and the complete Fisher orbit satisfies the rigidity bound

$$L_F \geq \pi, \tag{XIII.4}$$

with equality exactly for two-extreme-branch support. Every positive two-branch orbit has a universal logistic/tanh normal form and constant Schwarzian $\{w, x\} = -(s_+ - s_-)^2/2$.

Within the declared boundary-local curvature-measure class the bulk $s = 1$ branch is removed, leaving $\{0, -1, -2, -3\}$. Quotienting the homogeneous expansion history by the standard vacuum, curvature-degenerate, and dust powers leaves the unique nonstandard background direction a^{-1}, corresponding to $w = -2/3$. Exact expansion-coefficient projectors make the separation of geometric curvature from a physical $s = -2$ stress component operational rather than semantic.

The executed numerical audit adds evidence of a different logical type. The pinned CLASS implementation satisfies the declared a^{-1} background law, closed-mode normalization, transfer-column, precision, and null-limit checks. The

exact projectors and Hankel invariants recover the instantiated coefficients and show that retaining the negative closed-FLRW geometric curvature coefficient inside a physical moment measure creates exactly one negative inertia direction; after the curvature split the physical $\{\Lambda, X, m\}$ sector is positive. The supplied DESI DR2 BAO-only profile, however, has $q_X = 1.364$ and one-sided $p = 0.121$, so it does not detect a nonzero X amplitude.

The distinction between consistency and identification is therefore essential. Because the executed source lift fixes $w_X = -2/3$, recovering $s = -1$ from that calculation is not an independent discovery of the dilation spectrum. A decisive empirical test must first infer spectral locations freely, with covariance and model-order uncertainty propagated, and only then compare them with the valuation roots. A joint CMB+BAO+SNe+growth+lensing likelihood and the general two-parameter $(\Omega_X, \Omega_{2,\mathrm{val}})$ transfer implementation remain independent completion problems.

Finally, the finite $s = 0, -1$ density $\rho_{\Lambda X,0}$ is a reproducible background budget once H_0 and the dimensionless density fractions are supplied. It is not the bare quantum zero-point sum and does not evade the scale-covariance no-go theorem for topology-only dimensional amplitudes. The theorem-level result is therefore a finite homogeneous classification and invariant test architecture, not a microscopic derivation of vacuum energy.

DATA AND CODE AVAILABILITY

The theorem layer is analytic. The numerical section analyzes an accompanying execution-evidence package containing pinned CLASS outputs, transfer and matter-power products, validation summaries, DESI DR2 BAO profile/MCMC artifacts, scripts, and machine-readable background ledgers. The package identifies CLASS/`classy` v3.3.4 at commit `e85808324f51fc694d12e3ed7439552a3c3f9540` and Cobaya 3.6.2. The exact-invariant audit tables and source used for this revision are included with the manuscript package. The quoted DESI result is a fixed-curvature BAO-only pilot, not a full cosmological model-comparison likelihood. A persistent public repository identifier should accompany journal submission if the numerical section is retained.

[1] H. Hadwiger, *Vorlesungen über Inhalt, Oberfläche und Isoperimetrie*, Springer, Berlin, 1957.

[2] R. Schneider, *Convex Bodies: The Brunn-Minkowski Theory*, 2nd expanded ed., Cambridge University Press, Cambridge, 2014.

[3] R. Bhatia and C. Davis, "A better bound on the variance," *American Mathematical Monthly* **107** (2000), 353–357, doi:10.1080/00029890.2000.12005203.

[4] C. Clarkson, B. Bassett, and T. H.-C. Lu, "A general test of the Copernican Principle," *Physical Review Letters* **101** (2008), 011301, doi:10.1103/PhysRevLett.101.011301.

[5] G. P. Karev, "Replicator equations and the principle of minimal production of information," *Bulletin of Mathematical Biology* **72** (2010), 1124–1142, doi:10.1007/s11538-009-9484-9.

[6] Y. Hua and T. K. Sarkar, "Matrix pencil method for estimating parameters of exponentially damped/undamped sinusoids in noise," *IEEE Transactions on Acoustics, Speech, and Signal Processing* **38** (1990), 814–824, doi:10.1109/29.56027.

[7] T. Peter and G. Plonka, "A generalized Prony method for reconstruction of sparse sums of eigenfunctions of linear operators," *Inverse Problems* **29** (2013), 025001, doi:10.1088/0266-5611/29/2/025001.

[8] M. Fischer, B. Kostrzewa, J. Ostmeyer, K. Ottnad, M. Ueding, and C. Urbach, "On the generalised eigenvalue method and its relation to Prony and generalised pencil of function methods," *European Physical Journal A* **56** (2020), 206, doi:10.1140/epja/s10050-020-00205-w.

[9] E. V. Linder, "Cosmic growth history and expansion history," *Physical Review D* **72** (2005), 043529, doi:10.1103/PhysRevD.72.043529.

[10] J. Schmalzing, M. Kerscher, and T. Buchert, "Minkowski functionals in cosmology," in *Dark Matter in the Universe*, Proc. International School of Physics Enrico Fermi, Course CXXXII, S. Bonometto, J. R. Primack, and A. Provenzale, eds. (IOS Press, Amsterdam, 1996), pp. 281–291, arXiv:astro-ph/9508154.

[11] R. A. Battye, M. Bucher, and D. N. Spergel, "Domain Wall Dominated Universes," arXiv:astro-ph/9908047.

[12] M. M. Afshar, "Quasilocal Energy in FRW Cosmology," *Class. Quantum Grav.* **26**, 225005 (2009), doi:10.1088/0264-9381/26/22/225005.

[13] M. Oltean, H. Bazrafshan Moghaddam, and R. J. Epp, "Energy of cosmological spacetimes and perturbations: a quasilocal approach," *Class. Quantum Grav.* **38**, 085012 (2021), doi:10.1088/1361-6382/abeae3.

[14] J. Ben Achour and E. R. Livine, "The Thiemann Complexifier and the CVH algebra for Classical and Quantum FLRW Cosmology," *Phys. Rev. D* **96**, 066025 (2017), doi:10.1103/PhysRevD.96.066025.

[15] J. Lesgourgues, "The Cosmic Linear Anisotropy Solving System (CLASS) I: Overview," arXiv:1104.2932.

[16] D. Blas, J. Lesgourgues, and T. Tram, "The Cosmic Linear Anisotropy Solving System (CLASS) II: Approximation schemes," *JCAP* **07** (2011) 034, doi:10.1088/1475-7516/2011/07/034.

[17] J. Torrado and A. Lewis, "Cobaya: Code for Bayesian Analysis of hierarchical physical models," *JCAP* **05** (2021) 057, doi:10.1088/1475-7516/2021/05/057.

[18] DESI Collaboration, M. Abdul Karim *et al.*, "DESI DR2 results. II. Measurements of baryon acoustic oscillations and cosmological constraints," *Phys. Rev. D* **112**, 083515 (2025), doi:10.1103/tr6y-kpc6.