

MASTER THESIS

me

March 15, 2018

# Contents

<b>1</b>	<b>An Introduction to the Einstein Relation</b>	<b>4</b>
1.1	Hausdorff measure and Hausdorff dimension . . . . .	4
1.2	Weyl asymptotics and spectral dimension . . . . .	6
1.3	Diffusion processes and walk dimension . . . . .	6
1.3.1	From Dirichlet forms to Markov processes . . . . .	6
1.4	Other versions of the Einstein relation . . . . .	6
<b>2</b>	<b>Examples and Non-examples</b>	<b>7</b>
	<b>Bibliography</b>	<b>8</b>

### **Abstract**

We consider  $\mathbb{R}$  as well as  $\dim_{\mathcal{S}}, \dim_{\mathcal{H}} \subseteq \dim_{\mathcal{W}}$ .

# Chapter 1

## An Introduction to the Einstein Relation

In this introductory chapter, we wish to briefly expose the ingredients of the ER - the Hausdorff dimension  $\dim_{\mathcal{H}}$ , the spectral dimension  $\dim_{\mathcal{S}}$ , and the walk dimension  $\dim_{\mathcal{W}}$  - and state some of their properties.

### 1.1 Hausdorff measure and Hausdorff dimension

Although the concepts of Hausdorff measure and dimension are well-known, we give the definitions in the interest of completeness. In what follows, let  $(E, d)$  be a metric space.

**Definition 1.1** (Hausdorff outer measure). For fixed  $s \geq 0$ , any subset  $S \subseteq E$  and any  $\delta > 0$ , let

$$\mathcal{H}_{\delta}^s(S) := \inf \left\{ \sum_{i \in I} (\text{diam } U_i)^s : |I| \leq \aleph_0, S \subseteq \bigcup_{i \in I} U_i \subseteq E, \text{diam } U_i \leq \delta \right\},$$

i.e. the infimum is taken over all countable coverings of  $S$  with diameter at most  $\delta$ . The  $s$ -dimensional Hausdorff outer measure of  $S$  is now defined to be

$$\mathcal{H}^s(S) := \lim_{\delta \searrow 0} \mathcal{H}_{\delta}^s(S). \quad (1.1)$$

Observe that the limit in (1.1) exists or equals  $\infty$ , since  $\mathcal{H}_{\delta}^s(S)$  is monotonically nonincreasing in  $\delta$ , yet bounded from below by 0. Furthermore, it can be shown that  $\mathcal{H}^s$  defines a metric outer measure on  $E$ , thus restricting to a measure on a  $\sigma$ -algebra containing the Borel  $\sigma$ -algebra  $\mathcal{B}(E)$  (cf. [Mat99, p.54ff]). By definition, the obtained measure then is the  $s$ -dimensional Hausdorff measure which we will denote by  $\mathcal{H}^s$  as well.

In the special case of  $(E, d)$  being an Euclidean space, Hausdorff measures interpolate between the usual Lebesgue measures  $\lambda^n$ : For  $s = 0$ , we have simply  $\mathcal{H}^0(S) = \#S$ , whereas for any integer  $n > 0$ , it can be shown that there exists a constant  $c_n > 0$  depending only on  $n$  such that  $\mathcal{H}^n = c_n \lambda^n$ , where the constant evaluates to the volume of the unit ball.

Since exponential functions are monotonically increasing, the Hausdorff measures' dependence on  $s$  exhibits the same behaviour for a fixed set  $S$ . At the same time, simple estimates yield that if  $\mathcal{H}^s(S)$  is finite for some  $s$ , it vanishes for all  $s' < s$ , and conversely, if  $\mathcal{H}^s(S) > 0$ , then  $\mathcal{H}^{s'}(S) = \infty$  for all  $s' > s$ . Therefore, there exists precisely one real number  $s$  where  $\mathcal{H}(S)$  jumps from 0 to  $\infty$  (by possibly attaining any value of  $[0, \infty]$  there). This motivates the following definition of Hausdorff dimension:

**Definition 1.2.** The Hausdorff dimension  $\dim_{\mathcal{H}}(S)$  of  $S \subseteq E$  is defined as

$$\dim_{\mathcal{H}}(S) := \inf\{s \geq 0 : \mathcal{H}^s(S) > 0\}.$$

Due to the above discussion, we have the following equalities:

$$\begin{aligned} \dim_{\mathcal{H}}(S) &= \inf\{s \geq 0 : \mathcal{H}^s(S) > 0\} = \inf\{s \geq 0 : \mathcal{H}^s(S) = \infty\} \\ &= \sup\{s \geq 0 : \mathcal{H}^s(S) = 0\} = \sup\{s \geq 0 : \mathcal{H}^s(S) < \infty\}, \end{aligned}$$

providing some alternative characterisations of the Hausdorff dimension.

We further collect some important facts. To this end, let  $S, S'$  and  $S_1, S_2, \dots$  be subsets of  $E$  as before. Then, the following properties hold (cf. [Fal07, p.32f] for a discussion in the Euclidean setting; however all arguments adapt to our more general situation without complication):

**Monotonicity.** If  $S \subseteq S'$  then  $\dim_{\mathcal{H}}(S) \leq \dim_{\mathcal{H}}(S')$ .

**Countable Stability.** For a sequence  $(S_n)_{n \geq 1}$ , we have the equality

$$\dim_{\mathcal{H}}\left(\bigcup_{n \geq 1} S_n\right) = \sup_{n \geq 1} \dim_{\mathcal{H}}(S_n).$$

**Countable Sets.** If  $|S| \leq \aleph_0$  then  $\dim_{\mathcal{H}}(S) = 0$ .

**Hölder continuous maps.** If  $(E', d')$  is another metric space and  $f : E \rightarrow E'$  is  $\alpha$ -Hölder continuous for some  $\alpha \in (0, 1]$  then  $\dim_{\mathcal{H}}(f(S)) \leq \alpha^{-1} \dim_{\mathcal{H}}(S)$ . In particular, the Hausdorff dimension is invariant under a bi-Lipschitz transformation (i.e. an invertible map  $f$  with Hölder exponent  $\alpha = 1$  for both  $f$  and  $f^{-1}$ ).

**Euclidean Case.** If  $(E, d)$  happens to be an Euclidean space (or more generally a continuously differentiable manifold) of dimension  $n$  and  $S$  is an open subset then  $\dim_{\mathcal{H}}(S) = n$ .

We conclude this section by citing Hutchinson's theorem about the Hausdorff dimension of self-similar sets which will provide us with a plethora of interesting examples. For this, we recall that a map  $F : E \rightarrow E$  on a metric space  $(E, d)$  is a strict contraction if its Lipschitz constant satisfies

$$\text{Lip}_F := \sup_{\substack{x, y \in E \\ x \neq y}} \frac{d(F(x), F(y))}{d(x, y)} < 1.$$

If the stronger condition  $d(F(x), F(y)) = \text{Lip}_F d(x, y)$  holds for all  $x, y \in E$ , we call  $F$  a similitude with contraction factor  $\text{Lip}_F$ .

**Theorem 1.3** (Hutchinson, [Hut81]). *Let  $\mathcal{S} = \{S_1, \dots, S_N\}$  be a finite set of strict contractions on the Euclidean space  $\mathbb{R}^n$ . Then there exists a unique nonempty compact set denoted by  $|\mathcal{S}|$  invariant under  $\mathcal{S}$ , i.e.*

$$|\mathcal{S}| = \bigcup_{i=1}^N S_i(|\mathcal{S}|).$$

*Furthermore, assume that  $|\mathcal{S}|$  satisfies the open set condition (OSC) which means that there exists a nonempty open set  $O \subseteq E$  with the properties  $S_i(O) \subseteq O$  and  $S_i(O) \cap S_j(O) = \emptyset$  for all  $i, j = 1, \dots, N$  with  $i \neq j$ . Also assume that the maps  $S_i$  are similitudes with contraction factor  $r_i \in (0, 1)$ . Then,  $s = \dim_{\mathcal{H}}(|\mathcal{S}|)$  is the unique solution to the equation*

$$\sum_{i=1}^N r_i^s = 1$$

*and we have  $0 < \mathcal{H}^s(|\mathcal{S}|) < \infty$ .*

While uniqueness and existence of  $|\mathcal{S}|$  are still ensured for maps on a complete metric space, the open set condition is not sufficient for statements about the Hausdorff dimension, see [Sch96] for further discussion.

## 1.2 Weyl asymptotics and spectral dimension

## 1.3 Diffusion processes and walk dimension

### 1.3.1 From Dirichlet forms to Markov processes

We start with the following definition (cf. [MR12, Def. IV.1.13]):

**Definition 1.4.** Given a filtered probability space  $(\Omega, \mathcal{A}, \mathcal{F} = (\mathcal{F}_t)_{t \geq 0}, \mathbf{P})$  satisfying the usual conditions, an  $\mathcal{F}$ -adapted time-homogenous Markov process  $X = (X_t)_{t \geq 0}$  with state space  $E_{\Delta}$  is called a right process if it satisfies the strong Markov property for all  $\mathcal{F}$ -stopping times and all its trajectories are right continuous.

## 1.4 Other versions of the Einstein relation

Telcs, Mandelbrot, etc...

## Chapter 2

# Examples and Non-examples

# Bibliography

- [Fal07] K. Falconer. *Fractal Geometry: Mathematical Foundations and Applications*. 2nd ed. Wiley, Hoboken, 2007.
- [Hut81] John E Hutchinson. “Fractals and self similarity”. In: *Indiana University Mathematics Journal* 30.5 (1981), pp. 713–747.
- [Mat99] P. Mattila. *Geometry of Sets and Measures in Euclidean Spaces: Fractals and Rectifiability*. Cambridge Studies in Advanced Mathematics. Cambridge University Press, Cambridge, 1999.
- [MR12] Z.M. Ma and M. Röckner. *Introduction to the Theory of (Non-Symmetric) Dirichlet Forms*. Universitext. Springer, Berlin Heidelberg, 2012.
- [Sch96] Andreas Schief. “Self-similar sets in complete metric spaces”. In: *Proceedings of the American Mathematical Society* 124.2 (1996), pp. 481–490.