

PRILL-EXCEPTIONAL ÉTALE COVERS FROM RAYNAUD BUNDLES

ABSTRACT. Let C be a smooth projective connected complex curve of genus $g \geq 2$. We prove that C admits a connected finite étale Galois cover $f: X \rightarrow C$ such that

$$h^0(X, \mathcal{O}_X(f^{-1}(p))) \geq 2$$

for every $p \in C$. Let $V = R_g \otimes M$ be a determinant-normalised twist of the rank- g^8 Raynaud bundle. Then $\deg V = 0$, $\det V \cong \mathcal{O}_C$, and $H^0(C, V) = 0$, whereas $H^0(C, V(p)) \neq 0$ for every $p \in C$. We prove that V has finite monodromy and choose an irreducible summand W with $H^0(C, W(p)) \neq 0$ for all p . For the Galois cover defined by the monodromy of W , the associated bundle $f_*\mathcal{O}_X$ contains $\mathcal{O}_C$ and W as direct summands, which yields the claimed inequality.

1. INTRODUCTION

Let $f: X \rightarrow C$ be a finite morphism of smooth projective connected complex curves. Prill asked whether a fibre divisor $f^{-1}(p)$ can move in a pencil; equivalently, whether

$$h^0(X, \mathcal{O}_X(f^{-1}(p))) > 1$$

for a general point $p \in C$. The question appears in [1, Chapter VI, Exercise D] and, for curves of genus at least three, is recorded as [6, Problem 14]. Partial affirmative results were obtained in [3, 7]. Landesman and Litt subsequently constructed, for every curve of genus two, a connected finite étale cover of degree 36 satisfying the inequality for every $p \in C$ [5].

We call f *Prill-exceptional* when the inequality holds for every $p \in C$. The construction below works for every $g \geq 2$.

Theorem 1.1. *Let C be a smooth projective connected complex curve of genus $g \geq 2$. There is a connected finite étale Galois cover $f: X \rightarrow C$ such that*

$$h^0(X, \mathcal{O}_X(f^{-1}(p))) \geq 2$$

for every $p \in C$.

For $g \geq 3$, Theorem 1.1 answers [6, Problem 14] affirmatively. For a finite flat morphism, the function

$$p \mapsto h^0(X, \mathcal{O}_X(f^{-1}(p)))$$

is upper semicontinuous. The locus where this dimension is at least two is therefore closed. If it contains a dense open subset of C , it equals C . Thus the general-fibre and pointwise formulations are equivalent. We use the latter because the construction yields it directly.

The proof uses two properties of Raynaud's construction. The restricted bundle R_g satisfies $H^0(C, R_g \otimes \alpha) \neq 0$ for every $\alpha \in \text{Pic}^0(C)$, whereas the ambient bundle E_g becomes a direct sum of copies of one line bundle after pullback by $[g]: J \rightarrow J$. We choose a twist with trivial determinant while preserving the pointwise nonvanishing. On the induced étale pullback of C , the repeated line bundle is torsion, so the twisted bundle has finite linear monodromy.

Date: 10 July 2026.

2020 Mathematics Subject Classification. Primary 14H30; Secondary 14H60, 14H40.

Key words and phrases. Prill's problem, finite étale covers, Raynaud bundles, finite monodromy.

2. THE RAYNAUD BUNDLE

Write $J = \text{Jac}(C) = \text{Pic}^0(C)$, fix a theta divisor Θ on J , and choose an Abel–Jacobi embedding $i: C \hookrightarrow J$. The form of Raynaud’s construction needed here is recorded in [8, Section 3] and [2, Section 2.3].

Theorem 2.1 (Raynaud). *For every integer $n \geq 1$, there is a vector bundle E_n on J such that*

$$[n]^* E_n \cong H^0(J, \mathcal{O}_J(n\Theta))^* \otimes_{\mathbb{C}} \mathcal{O}_J(n\Theta). \quad (1)$$

Its restriction $R_n = i^ E_n$ is semistable, has rank n^g and slope g/n , and satisfies*

$$H^0(C, R_n \otimes \alpha) \neq 0 \quad \text{for every } \alpha \in \text{Pic}^0(C). \quad (2)$$

Take $n = g$ and set

$$R := R_g, \quad r := \text{rk}(R) = g^g.$$

Then R is semistable of slope one, so $\deg(R) = r$.

3. A DETERMINANT-NORMALISED TWIST

Lemma 3.1. *There is a line bundle $M \in \text{Pic}^{-1}(C)$ such that*

$$M^{\otimes r} \cong (\det R)^{-1} \quad \text{and} \quad H^0(C, R \otimes M) = 0. \quad (3)$$

Proof. The map

$$\text{Pic}^{-1}(C) \longrightarrow \text{Pic}^{-r}(C), \quad M \longmapsto M^{\otimes r},$$

is a translate of the multiplication-by- r isogeny of J . Consequently, the fibre

$$T := \{M \in \text{Pic}^{-1}(C) : M^{\otimes r} \cong (\det R)^{-1}\}$$

has cardinality r^{2g} .

Call $M \in \text{Pic}^{-1}(C)$ *bad* if $H^0(C, R \otimes M) \neq 0$. Such a section induces a nonzero morphism $M^{-1} \rightarrow R$. If $L \subset R$ is the saturation of its image, then $L \cong M^{-1}(D)$ for an effective divisor D . Since $\deg(M^{-1}) = 1$ and R is semistable of slope one,

$$1 + \deg D = \deg L \leq 1.$$

Thus $D = 0$, and M^{-1} is a degree-one line subbundle of R .

Fix a Jordan–Hölder filtration $0 = R_0 \subset R_1 \subset \cdots \subset R_m = R$. Given a degree-one line subbundle $L \subset R$, let k be minimal with $L \subset R_k$. The induced map $L \rightarrow R_k/R_{k-1}$ is nonzero. Since its source and target are stable of slope one, it is an isomorphism. Thus every bad M makes M^{-1} isomorphic to a rank-one Jordan–Hölder factor of R . The filtration has at most r factors, so at most r isomorphism classes of M are bad.

Because $g \geq 2$ and $r = g^g > 1$, one has $r^{2g} > r$. The set T therefore contains an M that is not bad. $\square$

Fix M as in Lemma 3.1 and put $V := R \otimes M$. Then

$$\deg V = 0, \quad \det V \cong \mathcal{O}_C, \quad H^0(C, V) = 0. \quad (4)$$

For every $p \in C$, the line bundle $M(p)$ has degree zero, so (2) gives

$$H^0(C, V(p)) = H^0(C, R \otimes M(p)) \neq 0. \quad (5)$$

4. FINITE MONODROMY

Proposition 4.1. *The vector bundle V is trivialised by a connected finite étale cover of C .*

Proof. Form the cartesian square

$$\begin{array}{ccc} \tilde{C} & \xrightarrow{\tilde{i}} & J \\ q \downarrow & & \downarrow [g] \\ C & \xrightarrow{i} & J. \end{array}$$

The morphism q is finite étale and surjective. By (1),

$$q^*V \cong H^0(J, \mathcal{O}_J(g\Theta))^* \otimes_{\mathbb{C}} (\tilde{i}^* \mathcal{O}_J(g\Theta) \otimes q^*M).$$

Let $\tilde{C}_0$ be a connected component of $\tilde{C}$. Its image under q is nonempty, open, and closed, hence equals C . Define

$$N := (\tilde{i}^* \mathcal{O}_J(g\Theta) \otimes q^*M)|_{\tilde{C}_0}.$$

Since $h^0(J, \mathcal{O}_J(g\Theta)) = g^g = r$, the restriction of q^*V to $\tilde{C}_0$ is $N^{\oplus r}$. Taking determinants and using $\det V \cong \mathcal{O}_C$ yields

$$N^{\otimes r} \cong \mathcal{O}_{\tilde{C}_0}.$$

Let d be the order of N ; then $d \mid r$. A chosen isomorphism $N^{\otimes d} \cong \mathcal{O}_{\tilde{C}_0}$ defines the finite étale μ_d -torsor

$$\mathrm{Spec}_{\tilde{C}_0} \left(\bigoplus_{a=0}^{d-1} N^{-a} \right) \longrightarrow \tilde{C}_0.$$

The pullback of N to this torsor is trivial. Any connected component maps surjectively to $\tilde{C}_0$; its composite with $\tilde{C}_0 \rightarrow C$ is a connected finite étale cover $h: C' \rightarrow C$ that trivialises V . $\square$

Passing to a connected finite étale Galois cover dominating h , we may regard V as the bundle associated with a complex representation of a finite group.

Proposition 4.2. *There is a nontrivial irreducible finite-monodromy vector bundle W on C such that*

$$H^0(C, W(p)) \neq 0 \quad \text{for every } p \in C.$$

Proof. Choose a connected finite étale Galois cover $u: Y \rightarrow C$ trivialising V , and write H for its Galois group. Fix a trivialisation $u^*V \cong \mathcal{O}_Y^{\oplus r}$. Since Y is connected and projective, $H^0(Y, \mathcal{O}_Y) = \mathbb{C}$; hence the descent matrices are constant and define a representation $H \rightarrow \mathrm{GL}_r(\mathbb{C})$. Maschke's theorem gives a decomposition

$$V \cong \bigoplus_{j=1}^s W_j^{\oplus m_j},$$

where the W_j are pairwise nonisomorphic irreducible finite-monodromy bundles. No W_j is trivial, because a trivial summand would contradict $H^0(C, V) = 0$ in (4).

For $1 \leq j \leq s$, set

$$Z_j := \{p \in C \mid H^0(C, W_j(p)) \neq 0\}.$$

Let $p_1, p_2: C \times C \rightarrow C$ be the projections and let $\Delta \subset C \times C$ be the diagonal. The restriction of $p_1^*W_j \otimes \mathcal{O}_{C \times C}(\Delta)$ to $C \times \{p\}$ is $W_j(p)$. Upper semicontinuity for the proper morphism p_2 shows that each Z_j is closed. The decomposition of V and (5) give $C = \bigcup_{j=1}^s Z_j$. An irreducible variety cannot be a finite union of proper closed subsets, so $Z_j = C$ for some j . $\square$

5. THE COVERING

The vector-bundle criterion used below is the étale case of [4, Lemma 5.4]; we include the argument for completeness.

Proof of Theorem 1.1. Let W be as in Proposition 4.2, and let

$$\rho: \pi_1(C, c) \longrightarrow \mathrm{GL}(W_c)$$

be its monodromy representation. Its image G is finite. The subgroup $\ker \rho$ determines a connected finite étale Galois cover $f: X \rightarrow C$ with Galois group G .

The bundle $f_*\mathcal{O}_X$ corresponds to the regular G -module $\mathbb{C}[G]$, which contains every irreducible complex representation of G with multiplicity equal to its dimension. Hence $\mathcal{O}_C$ and W are distinct direct summands of $f_*\mathcal{O}_X$.

For $p \in C$, the fibre divisor is the pullback of p , so

$$\mathcal{O}_X(f^{-1}(p)) \cong f^*\mathcal{O}_C(p).$$

The projection formula and the summands $\mathcal{O}_C, W \subset f_*\mathcal{O}_X$ give

$$\begin{aligned} H^0(X, \mathcal{O}_X(f^{-1}(p))) &\cong H^0(C, f_*\mathcal{O}_X \otimes \mathcal{O}_C(p)) \\ &\supset H^0(C, \mathcal{O}_C(p)) \oplus H^0(C, W(p)). \end{aligned}$$

Since p is effective, $h^0(C, \mathcal{O}_C(p)) \geq 1$; Proposition 4.2 gives $h^0(C, W(p)) \geq 1$. Thus the left-hand side has dimension at least two for every $p \in C$. $\square$

Remark 5.1. The construction gives no useful degree bound. The initial Raynaud bundle has rank g^8 , and the proof subsequently takes a Kummer cover and a Galois closure. In genus two, the degree-36 cover of [5] is numerically far more efficient.

Remark 5.2. The determinant condition is essential. Formula (1) shows only that the pullback of R_g is a direct sum of copies of one line bundle. Triviality of the determinant makes that line bundle torsion. Consequently, the twisted bundle has finite linear, rather than merely finite projective, monodromy.

REFERENCES

1. Enrico Arbarello, Maurizio Cornalba, Phillip A. Griffiths, and Joe Harris, *Geometry of algebraic curves. volume i*, Grundlehren der mathematischen Wissenschaften, vol. 267, Springer, New York, 1985, <https://doi.org/10.1007/978-1-4757-5323-3>.
2. Arnaud Beauville, *Vector bundles on curves and theta functions*, Moduli Spaces and Arithmetic Geometry, Advanced Studies in Pure Mathematics, vol. 45, Mathematical Society of Japan, 2006, <https://arxiv.org/abs/math/0502179>, pp. 145–156.
3. Indranil Biswas and David C. Butler, *On prill's problem*, Communications in Algebra **33** (2005), no. 1, 323–330, <https://doi.org/10.1081/AGB-200041017>.
4. Aaron Landesman and Daniel Litt, *Applications of the algebraic geometry of the Putman–Wieland conjecture*, Proceedings of the London Mathematical Society **127** (2023), no. 1, 116–133, <https://doi.org/10.1112/plms.12539>.
5. ———, *Prill's problem*, Algebraic Geometry **11** (2024), no. 2, 290–295, <https://doi.org/10.14231/AG-2024-009>.
6. Daniel Litt, *Problem 14*, Problems I Like, <https://www.problemsilike.com/14>, 2026, Accessed 10 July 2026.
7. N. Mohan Kumar, *Some remarks on prill's problem*, Affine Algebraic Geometry, Osaka University Press, Osaka, 2007, pp. 289–294.
8. Michel Raynaud, *Sections des fibrés vectoriels sur une courbe*, Bulletin de la Société Mathématique de France **110** (1982), 103–125, <https://doi.org/10.24033/bsmf.1955>.