


The Characteristic Direction: A Geometrical Approach to Differential Expression – Part Four



Network Analysis in Systems Biology

Neil Clark, PhD

Instructor, Ma'ayan Lab

Department of Pharmacology and Systems Therapeutics

Icahn School of Medicine at Mount Sinai, New York, NY 10029




Performance on High-Throughput Sequencing Data


- ▶ Study the relationship between differential binding of STAT3 to differential gene expression in large B-cell lymphoma
 - Hardee, Jennifer, et al. "STAT3 Targets Suggest Mechanisms of Aggressive Tumorigenesis in Diffuse Large B-Cell Lymphoma." *G3: Genes/ Genomes/ Genetics* 3.12 (2013): 2173-2185.
- ▶ Compare the performance of the characteristic direction approach in the setting of high-throughput sequencing technologies
 - Compare to DESeq

Performance on High-Throughput Sequencing Data

- ▶ Examine differential binding and expression between two subtypes of Large B-cell Lymphoma:
 - germinal center B-cell-like
 - activated B-cell-like
- ▶ To what degree are genes which are differentially bound by STAT3 also differentially expressed?



Performance on Synthetic Data



Summary

- ▶ We have proposed a new approach to differential expression
- ▶ Our approach successfully:
 - Addressed the curse of dimensionality
 - Takes into account the fact that genes work together rather than in isolation
- ▶ Our approach outperforms the current most popular methods
- ▶ We have very promising natural extensions of our method in the making.