


Iron deficiency

Iron deficiency, or **sideropaenia**, is the state in which a body lacks enough iron to supply its needs. Iron is present in all cells in the human body and has several vital functions, such as carrying oxygen to the tissues from the lungs as a key component of the hemoglobin protein, acting as a transport medium for electrons within the cells in the form of cytochromes, and facilitating oxygen enzyme reactions in various tissues. Too little iron can interfere with these vital functions and lead to morbidity and death.^[1]

Total body iron averages approximately 3.8 g in men and 2.3 g in women. In blood plasma, iron is carried tightly bound to the protein transferrin. There are several mechanisms that control human iron metabolism and safeguard against iron deficiency. The main regulatory mechanism is situated in the gastrointestinal tract. When loss of iron is not sufficiently compensated by intake of iron from the diet, a state of iron deficiency develops over time. When this state is uncorrected, it leads to iron-deficiency anemia. Before anemia occurs, the medical condition of iron deficiency without anemia is called latent iron deficiency (LID) or iron-deficient erythropoiesis (IDE).

Untreated iron deficiency can lead to iron-deficiency anemia, a common type of anemia.^[1] Anemia is a condition characterized by inadequate red blood cells (erythrocytes) or hemoglobin. When the body lacks sufficient amounts of iron, production of the protein hemoglobin is reduced. Hemoglobin binds to oxygen, enabling red blood cells to supply oxygenated blood throughout the body. Children, pre-menopausal women (women of child-bearing age) and people with poor diet are most susceptible to the disease. Most cases of iron-deficiency anemia are mild, but if not treated it can cause problems like fast or irregular heartbeat, complications during pregnancy, and delayed growth in infants and children that could affect their cognitive development and their behavior.^[2]

Iron deficiency	
Other names	Sideropenia, hypoferremia
	
Iron in heme	
Specialty	Hematology

Contents

Signs and symptoms

Causes

- Athletics
- Inadequate intake

Bioavailability

Diagnosis

Treatment

Food sources

Blood transfusion

See also

Notes

References

Further reading

External links


Signs and symptoms

Symptoms of iron deficiency can occur even before the condition has progressed to iron deficiency anemia.



Symptoms of iron deficiency are not unique to iron deficiency (i.e. not pathognomonic). Iron is needed for many enzymes to function normally, so a wide range of symptoms may eventually emerge, either as the secondary result of the anemia, or as other primary results of iron deficiency. Symptoms of iron deficiency include:

- fatigue
- dizziness/lightheadedness
- pallor
- hair loss
- twitches
- irritability
- weakness
- pica
- brittle or grooved nails
- hair thinning
- Plummer–Vinson syndrome: painful atrophy of the mucous membrane covering the tongue, the pharynx and the esophagus
- impaired immune function^[4]
- pagophagia
- restless legs syndrome^[5]


Continued iron deficiency may progress to anemia and worsening fatigue. Thrombocytosis, or an elevated platelet count, can also result. A lack of sufficient iron levels in the blood is a reason that some people cannot donate blood.



Deaths due to iron-deficiency anemia per million persons in 2012



Disability-adjusted life year for iron-deficiency anemia per 100,000 inhabitants in 2004^[3]



Signs and Symptoms in Children include:

- pale skin
- Fatigue
- Slowed growth and development
- poor appetite
- behavioral problems
- Abnormal rapid breathing
- Frequent infection

Iron Requirements in young children to teenagers^[6]

Age group	Recommended amount of iron a day
7 – 12 months	11 mg
1 – 3 years	7 mg
4 – 8 years	10 mg
9 – 13 years	8 mg
14 – 18 years, girls	15 mg
14 – 18 years, boys	11 mg

Causes

- blood loss (hemoglobin contains iron)
 - donation
 - excessive menstrual bleeding
 - non-menstrual bleeding
 - bleeding from the gastrointestinal tract (ulcers, hemorrhoids, ulcerative colitis, stomach or colon cancer, etc.)
 - rarely, laryngological bleeding or from the respiratory tract
- inadequate intake (see below)
- substances (in diet or drugs) interfering with iron absorption
 - Fluoroquinolone antibiotics^[7]
- malabsorption syndromes
- inflammation where it is adaptive to limit bacterial growth in infection, but is also present in many other chronic diseases such as Inflammatory bowel disease and rheumatoid arthritis
- parasitic infection

Though genetic defects causing iron deficiency have been studied in rodents, there are no known genetic disorders of human iron metabolism that directly cause iron deficiency.

Athletics

Possible reasons that athletics may contribute to lower iron levels includes mechanical hemolysis (destruction of red blood cells from physical impact), loss of iron through sweat and urine, gastrointestinal blood loss, and haematuria (presence of blood in urine).^{[8][9]} Although small amounts of iron are excreted in sweat and urine, these losses can generally be seen as insignificant even with

increased sweat and urine production, especially considering that athletes' bodies appear to become conditioned to retain iron better.^[8] Mechanical hemolysis is most likely to occur in high-impact sports, especially among long distance runners who experience "foot-strike hemolysis" from the repeated impact of their feet with the ground. Exercise-induced gastrointestinal bleeding is most likely to occur in endurance athletes. Haematuria in athletes is most likely to occur in those that undergo repetitive impacts on the body, particularly affecting the feet (such as running on a hard road, or Kendo) and hands (e.g. Conga or Candombe drumming). Additionally, athletes in sports that emphasize weight loss (e.g. ballet, gymnastics, marathon running, and wrestling) as well as sports that emphasize high-carbohydrate, low-fat diets, may be at an increased risk for iron deficiency.^{[8][9]}

Inadequate intake

A U.S. federal survey of food consumption determined that for women and men over the age of 19, average iron consumption from foods and beverages was 13.1 and 18.0 mg/day, respectively. For women, 16% in the age range 14–50 years consumed less than the Estimated Average Requirement (EAR), for men ages 19 and up, fewer than 3%.^[10] Consumption data were updated in a 2014 U.S. government survey and reported that for men and women ages 20 and older the average iron intakes were, respectively, 16.6 and 12.6 mg/day.^[11] People in the U.S. usually obtain adequate amounts of iron from their diets. However, subgroups like infants, young children, teenaged girls, pregnant women, and premenopausal women are at risk of obtaining less than the EAR.^[12] Socio-economic and racial differences further affect the rates of iron deficiency.^[13]

Bioavailability

Iron is needed for bacterial growth making its bioavailability an important factor in controlling infection.^[14] Blood plasma as a result carries iron tightly bound to transferrin, which is taken up by cells by endocytosing transferrin, thus preventing its access to bacteria.^[15] Between 15 and 20 percent of the protein content in human milk consists of lactoferrin^[16] that binds iron. As a comparison, in cow's milk, this is only 2 percent. As a result, breast fed babies have fewer infections.^[15] Lactoferrin is also concentrated in tears, saliva and at wounds to bind iron to limit bacterial growth. Egg white contains 12% conalbumin to withhold it from bacteria that get through the egg shell (for this reason, prior to antibiotics, egg white was used to treat infections).^[17]

To reduce bacterial growth, plasma concentrations of iron are lowered in a variety of systemic inflammatory states due to increased production of hepcidin which is mainly released by the liver in response to increased production of pro-inflammatory cytokines such as Interleukin-6. This functional iron deficiency will resolve once the source of inflammation is rectified; however, if not resolved, it can progress to Anaemia of Chronic Inflammation. The underlying inflammation can be caused by fever,^[18] inflammatory bowel disease, infections, Chronic Heart Failure (CHF), carcinomas, or following surgery.

Reflecting this link between iron bioavailability and bacterial growth, the taking of oral iron supplements in excess of 200 mg/day causes a relative overabundance of iron that can alter the types of bacteria that are present within the gut. There have been concerns regarding parenteral iron being administered whilst bacteremia is present, although this has not been borne out in clinical practice. A moderate iron deficiency, in contrast, can provide protection against acute infection, especially against organisms that reside within hepatocytes and macrophages, such as malaria and tuberculosis. This is mainly beneficial in regions with a high prevalence of these diseases and where standard treatment is unavailable.

Diagnosis

- A complete blood count can reveal microcytic anemia,^[19] although this is not always present – even when iron deficiency progresses to iron-deficiency anemia.
- Low serum ferritin (see *below*)
- Low serum iron
- High TIBC (total iron binding capacity), although this can be elevated in cases of anemia of chronic inflammation.
- It is possible that the fecal occult blood test might be positive, if iron deficiency is the result of gastrointestinal bleeding; although the sensitivity of the test may mean that in some cases it will be negative even with enteral blood loss.

As always, laboratory values have to be interpreted with the lab's reference values in mind and considering all aspects of the individual clinical situation.

Serum ferritin can be elevated in inflammatory conditions; so a normal serum ferritin may not always exclude iron deficiency, and the utility is improved by taking a concurrent C-reactive protein (CRP). The level of serum ferritin that is viewed as "high" depends on the condition. For example, in inflammatory bowel disease the threshold is 100, where as in chronic heart failure (CHF) the levels are 200.

Treatment

Before commencing treatment, there should be definitive diagnosis of the underlying cause for iron deficiency. This is particularly the case in older patients, who are most susceptible to colorectal cancer and the gastrointestinal bleeding it often causes. In adults, 60% of patients with iron-deficiency anemia may have underlying gastrointestinal disorders leading to chronic blood loss.^[20] It is likely that the cause of the iron deficiency will need treatment as well.

Upon diagnosis, the condition can be treated with iron supplements. The choice of supplement will depend upon both the severity of the condition, the required speed of improvement (e.g. if awaiting elective surgery) and the likelihood of treatment being effective (e.g. if has underlying IBD, is undergoing dialysis, or is having ESA therapy).

Examples of oral iron that are often used are ferrous sulfate, ferrous gluconate, or amino acid chelate tablets. Recent research suggests the replacement dose of iron, at least in the elderly with iron deficiency, may be as little as 15 mg per day of elemental iron.^[21]

Food sources

Mild iron deficiency can be prevented or corrected by eating iron-rich foods and by cooking in an iron skillet. Because iron is a requirement for most plants and animals, a wide range of foods provide iron. Good sources of dietary iron have heme-iron, as this is most easily absorbed and is not inhibited by medication or other dietary components. Three examples are red meat, poultry, and insects.^{[22][23]} Non-heme sources do contain iron, though it has reduced bioavailability. Examples are lentils, beans, leafy vegetables, pistachios, tofu, fortified bread, and fortified breakfast cereals.

Iron from different foods is absorbed and processed differently by the body; for instance, iron in meat (heme-iron source) is more easily absorbed than iron in grains and vegetables ("non-heme" iron sources).^[24] Minerals and chemicals in one type of food may also inhibit absorption of iron from another

type of food eaten at the same time.^[25] For example, oxalates and phytic acid form insoluble complexes which bind iron in the gut before it can be absorbed.

Because iron from plant sources is less easily absorbed than the heme-bound iron of animal sources, vegetarians and vegans should have a somewhat higher total daily iron intake than those who eat meat, fish or poultry.^[26] Legumes and dark-green leafy vegetables like broccoli, kale and oriental greens are especially good sources of iron for vegetarians and vegans. However, spinach and Swiss chard contain oxalates which bind iron, making it almost entirely unavailable for absorption. Iron from non-heme sources is more readily absorbed if consumed with foods that contain either heme-bound iron or vitamin C. This is due to a hypothesised "meat factor" which enhances iron absorption.^[27]

Following are two tables showing the richest foods in heme and non-heme iron.^[28] In both tables, food serving sizes may differ from the usual 100g quantity for relevancy reasons. Arbitrarily, the guideline is set at 18 mg, which is the USDA Recommended Dietary Allowance for women aged between 19 and 50.^[29]

Abstract: richest foods in *heme iron*

Food	Serving size	Iron	% guideline
<u>clam</u> ^[a]	100g	28 mg	155%
<u>pork liver</u>	100g	18 mg	100%
<u>lamb kidney</u>	100g	12 mg	69%
<u>cooked oyster</u>	100g	12 mg	67%
<u>cuttlefish</u>	100g	11 mg	60%
<u>lamb liver</u>	100g	10 mg	57%
<u>octopus</u>	100g	9.5 mg	53%
<u>mussel</u>	100g	6.7 mg	37%
<u>beef liver</u>	100g	6.5 mg	36%
<u>beef heart</u>	100g	6.4 mg	35%

Abstract: richest foods in *non-heme iron*

Food	Serving size	Iron	% guideline
<u>raw yellow beans</u>	100g	7 mg	35%
<u>spirulina</u>	15g	4.3 mg	24%
<u>falafel</u>	140g	4.8 mg	24%
<u>soybean kernels</u>	125ml=1/2cup	4.6 mg	23%
<u>spinach</u>	125g	4.4 mg	22%
<u>lentil</u>	125ml=1/2cup	3.5 mg	17.5%
<u>treacle (CSR Australia)</u>	20ml=1Tbsp	3.4 mg	17%
<u>molasses (BlueLabel Australia)</u>	20ml=1Tbsp	1.8 mg	9%
<u>candied ginger root</u>	15g~3p	1.7 mg	8.5%
<u>toasted sesame seeds</u>	10g	1.4 mg	7%
<u>cocoa (dry powder)</u>	5g~1Tbsp	.8 mg	4%

Food Recommendations for Children

Children at 6 months should start having solid food that contains enough iron, which could be found in both heme and non-heme iron^[33]

Heme iron :

- Red meat (for example, beef, pork, lamb, goat, or venison)
- Fatty fish
- Poultry (for example, chicken or turkey)
- Eggs

Non-heme iron :

- Iron-fortified infant cereals
- Tofu
- Beans and lentils
- Dark green leafy vegetables

Iron deficiency can have serious health consequences that diet may not be able to quickly correct; hence, an iron supplement is often necessary if the iron deficiency has become symptomatic.

Blood transfusion

Blood transfusion is sometimes used to treat iron deficiency with hemodynamic instability.^[34] Sometimes transfusions are considered for people who have chronic iron deficiency or who will soon go to surgery, but even if such people have low hemoglobin, they should be given oral treatment or intravenous iron.^[34]

See also

- Bahima disease
- CO2 fertilization effect

Notes

- a. Iron content in clams can vary considerably between types and modes of preparation, and the presence of aluminium could reduce iron bioavailability.^[30] The bioaccumulation of heavy metals in clams from highly contaminated areas may make regular consumption unsafe in the long term.^{[31][32]}

References

1. Centers for Disease Control and Prevention (3 April 1998). "Recommendations to Prevent and Control Iron Deficiency in the United States" (<https://www.cdc.gov/mmwr/preview/mmwr.html/00051880.htm>). *Morbidity and Mortality Weekly Report. Recommendations and Reports*. **47** (RR-3): 1–36. PMID 9563847 (<https://pubmed.ncbi.nlm.nih.gov/9563847>).

2. "Iron and Iron Deficiency" (<https://web.archive.org/web/20140908003331/http://www.cdc.gov/nutrition/everyone/basics/vitamins/iron.html>). *Centers for Disease Control and Prevention*. 23 February 2011. Archived from the original (<https://www.cdc.gov/nutrition/everyone/basic/s/vitamins/iron.html>) on 8 September 2014. Retrieved 12 August 2014.
3. "Mortality and Burden of Disease Estimates for WHO Member States in 2002" (<http://www.who.int/entity/healthinfo/statistics/bodgbdddeathdalyestimates.xls>) (xls). *World Health Organization*. 2002.
4. Wintergerst, E. S.; Maggini, S.; Hornig, D. H. (2007). "Contribution of Selected Vitamins and Trace Elements to Immune Function" (<http://doc.rero.ch/record/304705/files/S0029665108006939.pdf>) (PDF). *Annals of Nutrition and Metabolism*. **51** (4): 301–323. doi:10.1159/000107673 (<https://doi.org/10.1159/000107673>). PMID 17726308 (<https://pubmed.ncbi.nlm.nih.gov/17726308>).
5. Rangarajan, Sunad; D'Souza, George Albert. (April 2007). "Restless legs syndrome in Indian patients having iron deficiency anemia in a tertiary care hospital". *Sleep Medicine*. **8** (3): 247–51. doi:10.1016/j.sleep.2006.10.004 (<https://doi.org/10.1016/j.sleep.2006.10.004>). PMID 17368978 (<https://pubmed.ncbi.nlm.nih.gov/17368978>).
6. "Is your child getting enough iron?" (<https://www.mayoclinic.org/healthy-lifestyle/childrens-health/in-depth/iron-deficiency/art-20045634>). *Mayo Clinic*. Retrieved 26 April 2019.
7. Badal S, Her YF, Maher LJ 3rd (September 2015). "Nonantibiotic Effects of Fluoroquinolones in Mammalian Cells" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4571980>). *J Biol Chem*. **290** (36): 22287–97. doi:10.1074/jbc.M115.671222 (<https://doi.org/10.1074/jbc.M115.671222>). PMC 4571980 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4571980>). PMID 26205818 (<https://pubmed.ncbi.nlm.nih.gov/26205818>).
8. Nielson, Peter; Nachtigall, Detlef (October 1998). "Iron supplementation in athletes: current recommendations" (<http://camilleherron.com/wp-content/uploads/2011/05/ironsupplementation.pdf>) (PDF). *Sports Med*. **26** (4): 207–216. doi:10.2165/00007256-199826040-00001 (<https://doi.org/10.2165/00007256-199826040-00001>). PMID 9820921 (<https://pubmed.ncbi.nlm.nih.gov/9820921>). Retrieved 7 July 2013.
9. Chatard, Jean-Claude; Mujika, Iñigo; Guy, Claire; Lacour, Jean-René (April 1999). "Anaemia and Iron Deficiency in Athletes Practical Recommendations for Treatment" (<http://www.pilarmartinescudero.com/pdf/asignaturabiopatologia/LecturasRecomendadas/Anaemiaandirondeficiencyinathletes.pdf>) (PDF). *Sports Med*. **27** (4): 229–240. doi:10.2165/00007256-199927040-00003 (<https://doi.org/10.2165/00007256-199927040-00003>). PMID 10367333 (<https://pubmed.ncbi.nlm.nih.gov/10367333>). Retrieved 7 July 2013.
10. What We Eat In America, NHANES 2001–2002 (<https://www.ars.usda.gov/SP2UserFiles/Place/80400530/pdf/0102/usualintaketables2001-02.pdf>) Archived (<https://web.archive.org/web/20150106010550/https://www.ars.usda.gov/SP2UserFiles/Place/80400530/pdf/0102/usualintaketables2001-02.pdf>) 6 January 2015 at the Wayback Machine. see Table A12: Iron.
11. What We Eat In America, NHANES 2013-2014 (https://www.ars.usda.gov/ARSUserFiles/80400530/pdf/1314/Table_1_NIN_GEN_13.pdf).
12. "Office of Dietary Supplements - Iron" (<https://ods.od.nih.gov/factsheets/Iron-HealthProfessional/>). *ods.od.nih.gov*. Retrieved 15 November 2019.
13. "Office of Dietary Supplements - Iron" (<https://ods.od.nih.gov/factsheets/Iron-HealthProfessional/>). *ods.od.nih.gov*. Retrieved 15 November 2019.
14. Kluger, M. J.; Rothenburg, B. A. (1979). "Fever and reduced iron: Their interaction as a host defense response to bacterial infection". *Science*. **203** (4378): 374–376. doi:10.1126/science.760197 (<https://doi.org/10.1126/science.760197>). PMID 760197 (<https://pubmed.ncbi.nlm.nih.gov/760197>).
15. Nesse, R. M.; Williams, G. C. *Why We Get Sick: The New Science of Darwinian Medicine*. New York. page 30 ISBN 0-679-74674-9.

16. T. William Hutchens, Bo Lönnerdal; Lactoferrin: Interactions and Biological Functions (1997). page 379 on Google Books (https://books.google.com/books?id=KOHK0Zi8S08C&pg=PA379&vq=milk&source=gbs_search_s&cad=0)
17. Nesse, R. M.; Williams, G. C. *Why We Get Sick: The New Science of Darwinian Medicine*. New York. page 29 ISBN 0-679-74674-9.
18. Weinberg, E. D. (1984). "Iron withholding: A defense against infection and neoplasia". *Physiological Reviews*. **64** (1): 65–102. doi:10.1152/physrev.1984.64.1.65 (<https://doi.org/10.1152/physrev.1984.64.1.65>). PMID 6420813 (<https://pubmed.ncbi.nlm.nih.gov/6420813/>).
19. Longmore, Murray; Ian B. Wilkinson; Supaj Rajagoplan (2004). *Oxford Handbook of Clinical Medicine* (6th ed.). Oxford University Press. pp. 626–628. ISBN 0-19-852558-3.
20. Rockey D, Cello J (1993). "Evaluation of the gastrointestinal tract in patients with iron-deficiency anemia". *N Engl J Med*. **329** (23): 1691–5. doi:10.1056/NEJM199312023292303 (<https://doi.org/10.1056/NEJM199312023292303>). PMID 8179652 (<https://pubmed.ncbi.nlm.nih.gov/8179652/>).
21. Rimón E, Kagansky N, Kagansky M, Mechnick L, Mashiah T, Namir M, Levy S (2005). "Are we giving too much iron? Low-dose iron therapy is effective in octogenarians". *Am J Med*. **118** (10): 1142–7. doi:10.1016/j.amjmed.2005.01.065 (<https://doi.org/10.1016/j.amjmed.2005.01.065>). PMID 16194646 (<https://pubmed.ncbi.nlm.nih.gov/16194646/>).
22. Defoliart G (1992). "Insects as Human Food". *Crop Protection*. **11** (5): 395–99. doi:10.1016/0261-2194(92)90020-6 ([https://doi.org/10.1016/0261-2194\(92\)90020-6](https://doi.org/10.1016/0261-2194(92)90020-6)).
23. Bukkens SGF (1997). "The Nutritional Value of Edible Insects". *Ecol. Food. Nutr.* **36** (2–4): 287–319. doi:10.1080/03670244.1997.9991521 (<https://doi.org/10.1080/03670244.1997.9991521>).
24. Iron deficiency (<http://www.eatwell.gov.uk/healthissues/irondeficiency/>) Archived (<https://web.archive.org/web/20060808184739/http://www.eatwell.gov.uk/healthissues/irondeficiency/>) 8 August 2006 at the Wayback Machine Food Standards Agency.
25. Iron in diet (<https://www.nlm.nih.gov/medlineplus/ency/article/002422.htm>). MedlinePlus.
26. Mangels, Reed. Iron in the vegan diet (<http://www.vrg.org/nutrition/iron.htm>). The Vegetarian Resource Group.
27. Iron (<http://www.merckmanuals.com/home/disorders-of-nutrition/minerals/iron-deficiency-and-toxicity>). The Merck Manuals Online Medical Library.
28. iron rich foods (<http://iron.rich-foods.com/>) Archived (<https://web.archive.org/web/20170518140300/http://iron.rich-foods.com/>) 18 May 2017 at the Wayback Machine. Rich Foods.
29. Dietary Reference Intakes: Recommended Intakes for Individuals (http://www.iom.edu/Activities/Nutrition/SummaryDRIs/~media/Files/Activity%20Files/Nutrition/DRIs/5_Summary%20Table%20Tables%201-4.pdf) Archived (https://web.archive.org/web/20130906180147/http://www.iom.edu/Activities/Nutrition/SummaryDRIs/~media/Files/Activity%20Files/Nutrition/DRIs/5_Summary%20Table%20Tables%201-4.pdf) 6 September 2013 at the Wayback Machine National Academy of Sciences. Institute of Medicine. Food and Nutrition Board.
30. "Evaluation of clams as a food source of iron: Total iron, heme iron, aluminum, and in vitro iron bioavailability in live and processed clams - PubAg" (<https://pubag.nal.usda.gov/catalog/567004>). *pubag.nal.usda.gov*. Retrieved 29 April 2019.
31. Rahman, Md Rezaur; Hamdan, Sinin; Hossen, Md Faruk (2015). "Review on the Risk Assessment of Heavy Metals in Malaysian Clams" (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4427851>). *The Scientific World Journal*. **2015**: 905497. doi:10.1155/2015/905497 (<https://doi.org/10.1155/2015/905497>). PMC 4427851 (<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4427851>). PMID 26060840 (<https://pubmed.ncbi.nlm.nih.gov/26060840/>).

32. Fang, Zhan-Qiang; Cheung, R. Y. H.; Wong, M. H. (January 2003). "Heavy metals in oysters, mussels and clams collected from coastal sites along the Pearl River Delta, South China". *Journal of Environmental Sciences (China)*. **15** (1): 9–24. ISSN 1001-0742 (<https://www.worldcat.org/issn/1001-0742>). PMID 12602597 (<https://pubmed.ncbi.nlm.nih.gov/12602597/>).
33. CDC (3 December 2018). "Iron - Infant and Toddler Nutrition" (<https://www.cdc.gov/nutrition/InfantandToddlerNutrition/vitamins-minerals/iron.html>). *Centers for Disease Control and Prevention*. Retrieved 26 April 2019.
34. American Association of Blood Banks (24 April 2014), "Five Things Physicians and Patients Should Question" (<http://www.choosingwisely.org/doctor-patient-lists/american-association-of-blood-banks/>), *Choosing Wisely: an initiative of the ABIM Foundation*, American Association of Blood Banks, retrieved 25 July 2014, which cites
 - AABB (2011). *Guidelines for Patient Blood Management and Blood* (<https://web.archive.org/web/20141015211139/http://marketplace.aabb.org/EbusPPROD/Default.aspx?TabID=55&productId=1845>). ISBN 978-1-56395-333-0. Archived from the original (<http://marketplace.aabb.org/EbusPPROD/Default.aspx?TabID=55&productId=1845>) on 15 October 2014. Retrieved 28 July 2014.
 - Lin, DM; Lin, ES; Tran, MH (October 2013). "Efficacy and safety of erythropoietin and intravenous iron in perioperative blood management: a systematic review". *Transfusion Medicine Reviews*. **27** (4): 221–34. doi:10.1016/j.tmr.2013.09.001 (<https://doi.org/10.1016%2Fj.tmr.2013.09.001>). PMID 24135037 (<https://pubmed.ncbi.nlm.nih.gov/24135037/>).

Further reading

- Gropper, Sareen S; Smith, Jack L; Groff, James L (2009). "Enhancers and inhibitors of iron absorption" (<https://books.google.com/books?id=rXSO9YLr72YC&pg=PA474>). *Advanced Nutrition and Human Metabolism* (<https://books.google.com/books?id=rXSO9YLr72YC>) (5th ed.). Belmont, California: Wadsworth, Cengage Learning. ISBN 978-0-495-11657-8. Retrieved 2 October 2010.
- Umbreit, Jay (2005). "Iron deficiency: A concise review". *American Journal of Hematology*. **78** (3): 225–231. doi:10.1002/ajh.20249 (<https://doi.org/10.1002%2Fajh.20249>). PMID 15726599 (<https://pubmed.ncbi.nlm.nih.gov/15726599/>).
- Nutrition, Iron (2018). *Centers for Disease Control and Prevention*.
- Is Your Child Getting Enough Iron? (2019). *Mayo clinic*

External links

- Recommendations to Prevent and Control Iron Deficiency in the United States (<https://www.cdc.gov/mmwr/preview/mmwrhtml/00051880.htm>)
- NHS.UK (<https://web.archive.org/web/20151129023838/http://www.uhs.nhs.uk/Media/Controlleddocuments/Patientinformation/Digestionandurinaryhealth/Adviceforimprovingyourironintake-patientinformation.pdf>)

Classification **ICD-10:** E61.1 (<http://apps.who.int/classifications/icd10/browse/2016/en#/E61.1>) • **DiseasesDB:** 6947 (<http://www.diseasesdatabase.com/ddb6947.htm>)

This page was last edited on 7 January 2020, at 11:19 (UTC).

Text is available under the [Creative Commons Attribution-ShareAlike License](#); additional terms may apply. By using this site, you agree to the [Terms of Use](#) and [Privacy Policy](#). Wikipedia® is a registered trademark of the [Wikimedia Foundation, Inc.](#), a non-profit organization.