





### Scientific Computing Lab

**Ordinary Differential Equations** 

**Explicit Discretization** 

Kaveh Rahnema

Equation with a function y(t) and derivatives of y(t)

$$y(t)+\dot{y}(t)=0$$

$$\sum_{i=0}^{n} a_{i} y^{(i)}(t) = 0$$

typical: development of a variable over time


$$\dot{y}(t) = f(t, y(t))$$

- Example: radioactive decay
  - Half-life: Period of time in which half of the atoms decay
  - Decay constant k: Describes rate of decay and thus half-life

$$\frac{dr(t)}{dt} = -k \cdot r(t)$$

Example: pendulum

$$\frac{d^2\alpha}{dt^2} + \frac{k}{m} \cdot \frac{d\alpha}{dt} + \frac{g}{I} \cdot \sin(\alpha) = 0$$




- Example: population growth
- populations P and Q

$$\frac{dp(t)}{dt} = (2 - p - q) \cdot p$$

$$\frac{dq(t)}{dt} = (2 - p - q) \cdot q$$


#### competition



Example: population growth populations P and Q – predator-prey

$$\frac{dp(t)}{dt} = (2-p+q) \cdot p$$

$$\frac{dq(t)}{dt} = (2 - 10p - q) \cdot q$$




- Solving = finding y(t)
- Two ways to solve the ODE:
  - Analytically
  - Numerically


- Solving = finding y(t)
- Two ways to solve the ODE:
  - Analytically
  - Numerically
    - → Needs discretization (timestepping)

Time t	0	5	10	15	20	25	30	35	40	45
$\dot{y}(t)$	5	1	0	-1	0	0	0	0	2	5


- Time in seconds,  $\dot{y}$  in m/s
- Any ideas what y describes?




Time t	0	5	10	15	20	25	30	35	40	45
$\dot{y}(t)$	5	1	0	-1	0	0	0	0	2	5




Time t	0	5	10	15	20	25	30	35	40	45
$\dot{y}(t)$	5	1	0	-1	0	0	0	0	2	5




Time t	0	5	10	15	20	25	30	35	40	45
$\dot{y}(t)$	5	1	0	-1	0	0	0	0	2	5




Time t	0	5	10	15	20	25	30	35	40	45
$\dot{y}(t)$	5	1	0	-1	0	0	0	0	2	5




	Time t	0	5	10	15	20	25	30	35	40	45
•	$\dot{y}(t)$	5	1	0	-1	0	0	0	0	2	5




	Time t	0	5	10	15	20	25	30	35	40	45
•	$\dot{y}(t)$	5	1	0	-1	0	0	0	0	2	5




Time t	0	5	10	15	20	25	30	35	40	45
$\dot{y}(t)$	5	1	0	-1	0	0	0	0	2	5



Time t	0	5	10	15	20	25	30	35	40	45
$\dot{y}(t)$	5	1	0	-1	0	0	0	0	2	5




Time t	0	5	10	15	20	25	30	35	40	45
$\dot{y}(t)$	5	1	0	-1	0	0	0	0	2	5



- explicit Euler
- method of Heun
- Runge-Kutta

$$> y_{n+1} = F(y_n, t_n, \Delta t)$$

#### explicit Euler



$$y_{n+1} = y_n + \Delta t \cdot \dot{y}_n$$
$$= y_n + \Delta t \cdot f(t_n, y_n)$$

method of Heun

$$y_{n+1} = y_n + \Delta t \cdot \frac{1}{2} (\dot{y}_n + \dot{y}_{n+1})$$

with

$$\dot{y}_{n+1} = f(t_{n+1}, y_n + \Delta t \cdot f(t_n, y_n))$$

#### Runge-Kutta

$$y_{n+1} = y_n + \Delta t \cdot \frac{1}{6} (Y_1' + 2Y_2' + 2Y_3' + Y_4')$$
 with 
$$Y_1' = f(t_n, y_n)$$
 
$$Y_2' = f(t_{n+\frac{1}{2}}, y_n + \frac{\Delta t}{2} \cdot Y_1')$$
 
$$Y_3' = f(t_{n+\frac{1}{2}}, y_n + \frac{\Delta t}{2} \cdot Y_2')$$

 $Y_{A}'=f(t_{n+1},y_n+\Delta t\cdot Y_{3}')$ 

### What is Efficiency?

- number of operations?
- runtime?
- accuracy?

### What is Efficiency?

- number of operations?
- runtime?
- accuracy?

### What is Efficiency?

- number of operations?
- runtine?
- accuracy

relation accuracy/cost !!!!!

#### Accuracy

definition of convergence:

$$||u_{exact} - u_{approx}|| = O(dt^p)$$

experimental computation?