

Representability of cohomology theories

Fernando Muro

Universidad de Sevilla
Departamento de Álgebra

Joint Mathematical Conference CSASC 2010
Prague, January 2010

Cohomology

Space = pointed simplicial (or cell) complex.

Definition

A reduced *cohomology theory* H is:

Topology \rightarrow *Algebra*

space $X \mapsto H^n(X)$ *abelian group*, $n \in \mathbb{Z}$,

map $X \xrightarrow{f} Y \mapsto H^n(X) \xleftarrow{H^n(f)} H^n(Y)$ *homomorphism*,

homotopic $f \simeq g: X \rightarrow Y \mapsto H^n(f) = H^n(g)$ *equal*,

base point union $\coprod_{i \in I} X_i \mapsto \prod_{i \in I} H^n(X_i)$ *product*,

cofiber sequence

exact sequence

$X \rightarrow Y \rightarrow C_f \rightarrow \Sigma X \mapsto H^n(X) \leftarrow H^n(Y) \leftarrow H^n(C_f) \leftarrow H^n(\Sigma X),$

$H^n(X) \cong H^{n+1}(\Sigma X)$ *suspension formula*.

Cohomology

Space = pointed simplicial (or cell) complex.

Definition

A reduced *cohomology theory* H is:

Topology \rightarrow *Algebra*

space $X \mapsto H^n(X)$ *abelian group*, $n \in \mathbb{Z}$,

map $X \xrightarrow{f} Y \mapsto H^n(X) \xleftarrow{H^n(f)} H^n(Y)$ *homomorphism*,

homotopic $f \simeq g: X \rightarrow Y \mapsto H^n(f) = H^n(g)$ *equal*,

base point union $\coprod_{i \in I} X_i \mapsto \prod_{i \in I} H^n(X_i)$ *product*,

cofiber sequence

exact sequence

$X \rightarrow Y \rightarrow C_f \rightarrow \Sigma X \mapsto H^n(X) \leftarrow H^n(Y) \leftarrow H^n(C_f) \leftarrow H^n(\Sigma X),$

$H^n(X) \cong H^{n+1}(\Sigma X)$ *suspension formula*.

Cohomology

Space = pointed simplicial (or cell) complex.

Definition

A reduced *cohomology theory* H is:

Topology \rightarrow *Algebra*

space $X \mapsto H^n(X)$ *abelian group*, $n \in \mathbb{Z}$,

map $X \xrightarrow{f} Y \mapsto H^n(X) \xleftarrow{H^n(f)} H^n(Y)$ *homomorphism*,

homotopic $f \simeq g: X \rightarrow Y \mapsto H^n(f) = H^n(g)$ *equal*,

base point union $\coprod_{i \in I} X_i \mapsto \prod_{i \in I} H^n(X_i)$ *product*,

cofiber sequence

exact sequence

$X \rightarrow Y \rightarrow C_f \rightarrow \Sigma X \mapsto H^n(X) \leftarrow H^n(Y) \leftarrow H^n(C_f) \leftarrow H^n(\Sigma X),$

$H^n(X) \cong H^{n+1}(\Sigma X)$ *suspension formula*.

Cohomology

Space = pointed simplicial (or cell) complex.

Definition

A reduced *cohomology theory* H is:

Topology \rightarrow *Algebra*

space $X \mapsto H^n(X)$ *abelian group*, $n \in \mathbb{Z}$,

map $X \xrightarrow{f} Y \mapsto H^n(X) \xleftarrow{H^n(f)} H^n(Y)$ *homomorphism*,

homotopic $f \simeq g: X \rightarrow Y \mapsto H^n(f) = H^n(g)$ *equal*,

base point union $\coprod_{i \in I} X_i \mapsto \prod_{i \in I} H^n(X_i)$ *product*,

cofiber sequence

exact sequence

$X \rightarrow Y \rightarrow C_f \rightarrow \Sigma X \mapsto H^n(X) \leftarrow H^n(Y) \leftarrow H^n(C_f) \leftarrow H^n(\Sigma X),$

$H^n(X) \cong H^{n+1}(\Sigma X)$ *suspension formula*.

Cohomology

Space = pointed simplicial (or cell) complex.

Definition

A reduced *cohomology theory* H is:

Topology \rightarrow *Algebra*

space $X \mapsto H^n(X)$ *abelian group*, $n \in \mathbb{Z}$,

map $X \xrightarrow{f} Y \mapsto H^n(X) \xleftarrow{H^n(f)} H^n(Y)$ *homomorphism*,

homotopic $f \simeq g: X \rightarrow Y \mapsto H^n(f) = H^n(g)$ *equal*,

base point union $\coprod_{i \in I} X_i \mapsto \prod_{i \in I} H^n(X_i)$ *product*,

cofiber sequence

exact sequence

$X \rightarrow Y \rightarrow C_f \rightarrow \Sigma X \mapsto H^n(X) \leftarrow H^n(Y) \leftarrow H^n(C_f) \leftarrow H^n(\Sigma X),$

$H^n(X) \cong H^{n+1}(\Sigma X)$ *suspension formula*.

Cohomology

Space = pointed simplicial (or cell) complex.

Definition

A reduced *cohomology theory* H is:

Topology \rightarrow *Algebra*

space $X \mapsto H^n(X)$ *abelian group*, $n \in \mathbb{Z}$,

map $X \xrightarrow{f} Y \mapsto H^n(X) \xleftarrow{H^n(f)} H^n(Y)$ *homomorphism*,

homotopic $f \simeq g: X \rightarrow Y \mapsto H^n(f) = H^n(g)$ *equal*,

base point union $\coprod_{i \in I} X_i \mapsto \prod_{i \in I} H^n(X_i)$ *product*,

cofiber sequence

exact sequence

$X \rightarrow Y \rightarrow C_f \rightarrow \Sigma X \mapsto H^n(X) \leftarrow H^n(Y) \leftarrow H^n(C_f) \leftarrow H^n(\Sigma X),$

$H^n(X) \cong H^{n+1}(\Sigma X)$ *suspension formula*.

Cohomology

Space = pointed simplicial (or cell) complex.

Definition

A reduced *cohomology theory* H is:

Topology \rightarrow *Algebra*

space $X \mapsto H^n(X)$ *abelian group*, $n \in \mathbb{Z}$,

map $X \xrightarrow{f} Y \mapsto H^n(X) \xleftarrow{H^n(f)} H^n(Y)$ *homomorphism*,

homotopic $f \simeq g: X \rightarrow Y \mapsto H^n(f) = H^n(g)$ *equal*,

base point union $\coprod_{i \in I} X_i \mapsto \prod_{i \in I} H^n(X_i)$ *product*,

cofiber sequence

exact sequence

$X \rightarrow Y \rightarrow C_f \rightarrow \Sigma X \mapsto H^n(X) \leftarrow H^n(Y) \leftarrow H^n(C_f) \leftarrow H^n(\Sigma X),$

$H^n(X) \cong H^{n+1}(\Sigma X)$ *suspension formula*.

Cohomology

Space = pointed simplicial (or cell) complex.

Definition

A reduced *cohomology theory* H is:

Topology \rightarrow *Algebra*

space $X \mapsto H^n(X)$ *abelian group*, $n \in \mathbb{Z}$,

map $X \xrightarrow{f} Y \mapsto H^n(X) \xleftarrow{H^n(f)} H^n(Y)$ *homomorphism*,

homotopic $f \simeq g: X \rightarrow Y \mapsto H^n(f) = H^n(g)$ *equal*,

base point union $\coprod_{i \in I} X_i \mapsto \prod_{i \in I} H^n(X_i)$ *product*,

cofiber sequence

exact sequence

$X \rightarrow Y \rightarrow C_f \rightarrow \Sigma X \mapsto H^n(X) \leftarrow H^n(Y) \leftarrow H^n(C_f) \leftarrow H^n(\Sigma X),$

$H^n(X) \cong H^{n+1}(\Sigma X)$ *suspension formula*.

Cohomology

Space = pointed simplicial (or cell) complex.

Definition

A reduced *cohomology theory* H on compact spaces is:

Topology \rightarrow *Algebra*

space $X \mapsto H^n(X)$ *abelian group*, $n \in \mathbb{Z}$,

map $X \xrightarrow{f} Y \mapsto H^n(X) \xleftarrow{H^n(f)} H^n(Y)$ *homomorphism*,

homotopic $f \simeq g: X \rightarrow Y \mapsto H^n(f) = H^n(g)$ *equal*,

base point union $\coprod_{i \in I} X_i \mapsto \prod_{i \in I} H^n(X_i)$ *product*, I *finite*,

cofiber sequence


exact sequence

$X \rightarrow Y \rightarrow C_f \rightarrow \Sigma X \mapsto H^n(X) \leftarrow H^n(Y) \leftarrow H^n(C_f) \leftarrow H^n(\Sigma X),$

$H^n(X) \cong H^{n+1}(\Sigma X)$ *suspension formula*.


$$X \xrightarrow{f} Y \xrightarrow{i} C_f \xrightarrow{q} \Sigma X$$

Cofiber sequence




$$X \xrightarrow{f} Y \xrightarrow{i} C_f \xrightarrow{q} \Sigma X$$

Cofiber sequence





$$X \xrightarrow{f} Y \xrightarrow[\text{inclusion}]{i} C_f \xrightarrow{q} \Sigma X$$

mapping cone




Cofiber sequence



$$X \xrightarrow{f} Y \xrightarrow[\text{inclusion}]{i} C_f \xrightarrow[\text{collapse}]{q} \Sigma X \quad \text{suspension}$$

mapping cone



Examples of cohomologies

Example

- Singular cohomology $H^*(X, \mathbb{Z})$, defined on *all* spaces,

$$H^n(\text{discrete}) = 0, \text{ for } n \neq 0,$$

$$H^0(X, \mathbb{Z}) = \text{pointed maps } X \rightarrow \mathbb{Z}.$$

► course

- Complex K -theory $K^*(X)$, X *compact*,

$K^0(X)$ = stable isomorphism classes of \mathbb{C} -vector bundles/ X ,

$K^n(X) \cong K^{n+2}(X)$, $n \in \mathbb{Z}$, Bott periodicity,

$K^1(X) \cong K^0(\Sigma X)$.

Examples of cohomologies

Example

- Singular cohomology $H^*(X, \mathbb{Z})$, defined on *all* spaces,

$$H^n(\text{discrete}) = 0, \text{ for } n \neq 0,$$

$$H^0(X, \mathbb{Z}) = \text{pointed maps } X \rightarrow \mathbb{Z}.$$

► course

- Complex K -theory $K^*(X)$, X *compact*,

$$K^0(X) = \text{stable isomorphism classes of } \mathbb{C}\text{-vector bundles}/X,$$

$$K^n(X) \cong K^{n+2}(X), \quad n \in \mathbb{Z}, \text{ Bott periodicity},$$

$$K^1(X) \cong K^0(\Sigma X).$$

Definition

A **spectrum** $E = \{E_0, E_1, \dots, E_n, \dots\}$ is a sequence of spaces together with **bonding maps**,

$$\Sigma E_n \longrightarrow E_{n+1}, \quad n \geq 0.$$

An **Ω -spectrum** is a spectrum E where the adjoints of the bonding maps $E_n \rightarrow \Omega E_{n+1}$ are homotopy equivalences.

Example

Any space X defines a spectrum $\Sigma^\infty X = \{X, \Sigma X, \dots, \Sigma^n X, \dots\}$, which is not an Ω -spectrum.

Definition

A **spectrum** $E = \{E_0, E_1, \dots, E_n, \dots\}$ is a sequence of spaces together with **bonding maps**,

$$\Sigma E_n \longrightarrow E_{n+1}, \quad n \geq 0.$$

An **Ω -spectrum** is a spectrum E where the adjoints of the bonding maps $E_n \rightarrow \Omega E_{n+1}$ are homotopy equivalences.

Example

Any space X defines a spectrum $\Sigma^\infty X = \{X, \Sigma X, \dots, \Sigma^n X, \dots\}$, which is not an Ω -spectrum.

Definition

A **spectrum** $E = \{E_0, E_1, \dots, E_n, \dots\}$ is a sequence of spaces together with **bonding maps**,

$$\Sigma E_n \longrightarrow E_{n+1}, \quad n \geq 0.$$

An **Ω -spectrum** is a spectrum E where the adjoints of the bonding maps $E_n \rightarrow \Omega E_{n+1}$ are homotopy equivalences.

Example

Any space X defines a spectrum $\Sigma^\infty X = \{X, \Sigma X, \dots, \Sigma^n X, \dots\}$, which is not an Ω -spectrum.

Theorem (G. W. Whitehead'62)

A spectrum E *represents* a cohomology theory H defined on *compact spaces* by

$$H^n(X) = \lim_{k \rightarrow \infty} [\Sigma^{k-n} X, E_k], \quad n \in \mathbb{Z},$$

where $[-, -]$ denotes the set of *homotopy classes* of maps.

If E is an Ω -spectrum then E *represents* a cohomology theory H defined on *all spaces* by the formula above.

Theorem (G. W. Whitehead'62)

A spectrum E *represents* a cohomology theory H defined on *compact spaces* by

$$H^n(X) = \lim_{k \rightarrow \infty} [\Sigma^{k-n} X, E_k], \quad n \in \mathbb{Z},$$

where $[-, -]$ denotes the set of *homotopy classes* of maps.

If E is an Ω -spectrum then E *represents* a cohomology theory H defined on *all spaces* by the formula above.

Classical representability theorems

Theorem (E. H. Brown'63)

*Any cohomology theory defined on **all** spaces is represented by a spectrum.*

Theorem (J. F. Adams'71)

*Any cohomology theory defined on **compact** spaces is represented by a spectrum.*

The following corollary can be applied to complex K -theory.

Corollary

*Any cohomology theory on **compact** spaces can be extended to a cohomology theory defined on **all** spaces.*

Classical representability theorems

Theorem (E. H. Brown'63)

*Any cohomology theory defined on **all** spaces is represented by a spectrum.*

Theorem (J. F. Adams'71)

*Any cohomology theory defined on **compact** spaces is represented by a spectrum.*

The following corollary can be applied to complex K -theory.

Corollary

*Any cohomology theory on **compact** spaces can be extended to a cohomology theory defined on **all** spaces.*

Classical representability theorems

Theorem (E. H. Brown'63)

*Any cohomology theory defined on **all** spaces is represented by a spectrum.*

Theorem (J. F. Adams'71)

*Any cohomology theory defined on **compact** spaces is represented by a spectrum.*

The following corollary can be applied to complex K -theory.

Corollary


*Any cohomology theory on **compact** spaces can be extended to a cohomology theory defined on **all** spaces.*

Spectrum for singular cohomology

$$E = \{\mathbb{Z}, S^1, \mathbb{C}P^\infty, \dots\} \text{ with } H^*(X) = H^*(X, \mathbb{Z}).$$


Spectrum for singular cohomology

$$E = \{\mathbb{Z}, S^1, \mathbb{C}P^\infty, \dots\} \text{ with } H^*(X) = H^*(X, \mathbb{Z}).$$




Spectrum for singular cohomology

$$E = \{\mathbb{Z}, S^1, \mathbb{C}P^\infty, \dots\} \text{ with } H^*(X) = H^*(X, \mathbb{Z}).$$




Spectrum for singular cohomology

$$E = \{\mathbb{Z}, S^1, \mathbb{C}P^\infty, \dots\} \text{ with } H^*(X) = H^*(X, \mathbb{Z}).$$




Spectrum for singular cohomology

$$E = \{\mathbb{Z}, S^1, \mathbb{C}P^\infty, \dots\} \text{ with } H^*(X) = H^*(X, \mathbb{Z}).$$




Spectrum for singular cohomology

$$E = \{\mathbb{Z}, S^1, \mathbb{C}P^\infty, \dots\} \text{ with } H^*(X) = H^*(X, \mathbb{Z}).$$




Spectrum for singular cohomology

$$E = \{\mathbb{Z}, S^1, \mathbb{C}P^\infty, \dots\} \text{ with } H^*(X) = H^*(X, \mathbb{Z}).$$




Spectrum for singular cohomology

$$E = \{\mathbb{Z}, S^1, \mathbb{C}P^\infty, \dots\} \text{ with } H^*(X) = H^*(X, \mathbb{Z}).$$




Spectrum for singular cohomology

$$E = \{\mathbb{Z}, S^1, \mathbb{C}P^\infty, \dots\} \text{ with } H^*(X) = H^*(X, \mathbb{Z}).$$




Spectrum for singular cohomology

$$E = \{\mathbb{Z}, S^1, \mathbb{C}P^\infty, \dots\} \text{ with } H^*(X) = H^*(X, \mathbb{Z}).$$



Spectrum for singular cohomology

$$E = \{\mathbb{Z}, S^1, \mathbb{C}P^\infty, \dots\} \text{ with } H^*(X) = H^*(X, \mathbb{Z}).$$

$$\Sigma S^1 = S^2 = \text{[sphere diagram]} = \mathbb{C}P^1 \xrightarrow{\text{inclusion}} \mathbb{C}P^\infty$$


Stable homotopy

Recall that cohomology regards suspension as an invertible functor:

$$H^n(X) \cong H^{n+1}(\Sigma X).$$

Definition

The *compact stable homotopy category* \mathbf{SH}^c :

- *Objects*: (X, n) , X compact space, $n \in \mathbb{Z}$,

$$(X, n) \sim \Sigma^n X.$$

- *Morphisms*: $\mathrm{Hom}((X, n), (Y, m)) = \lim_{k \rightarrow \infty} [\Sigma^{k+n} X, \Sigma^{k+m} Y].$
- *Suspension*: $\Sigma(X, n) = (X, n+1) \cong (\Sigma X, n).$
- *Exact triangles*: $(X, n) \rightarrow (Y, n) \rightarrow (C_f, n) \rightarrow \Sigma(X, n)$ coming from cofiber sequences.

Stable homotopy

Recall that cohomology regards suspension as an invertible functor:

$$H^n(X) \cong H^{n+1}(\Sigma X).$$

Definition

The *compact stable homotopy category* \mathbf{SH}^c :

- *Objects*: (X, n) , X compact space, $n \in \mathbb{Z}$,

$$(X, n) \sim \Sigma^n X.$$

- *Morphisms*: $\mathrm{Hom}((X, n), (Y, m)) = \varinjlim_{k \rightarrow \infty} [\Sigma^{k+n} X, \Sigma^{k+m} Y].$
- *Suspension*: $\Sigma(X, n) = (X, n+1) \cong (\Sigma X, n).$
- *Exact triangles*: $(X, n) \rightarrow (Y, n) \rightarrow (C_f, n) \rightarrow \Sigma(X, n)$ coming from cofiber sequences.

Stable homotopy

Recall that cohomology regards suspension as an invertible functor:

$$H^n(X) \cong H^{n+1}(\Sigma X).$$

Definition

The *compact stable homotopy category* \mathbf{SH}^c :

- *Objects*: (X, n) , X compact space, $n \in \mathbb{Z}$,

$$(X, n) \sim \Sigma^n X.$$

- *Morphisms*: $\mathrm{Hom}((X, n), (Y, m)) = \lim_{k \rightarrow \infty} [\Sigma^{k+n} X, \Sigma^{k+m} Y].$
- *Suspension*: $\Sigma(X, n) = (X, n+1) \cong (\Sigma X, n).$
- *Exact triangles*: $(X, n) \rightarrow (Y, n) \rightarrow (C_f, n) \rightarrow \Sigma(X, n)$ coming from cofiber sequences.

Stable homotopy

Recall that cohomology regards suspension as an invertible functor:

$$H^n(X) \cong H^{n+1}(\Sigma X).$$

Definition

The *compact stable homotopy category* \mathbf{SH}^c :

- *Objects*: (X, n) , X compact space, $n \in \mathbb{Z}$,

$$(X, n) \sim \Sigma^n X.$$

- *Morphisms*: $\mathrm{Hom}((X, n), (Y, m)) = \lim_{k \rightarrow \infty} [\Sigma^{k+n} X, \Sigma^{k+m} Y].$
- *Suspension*: $\Sigma(X, n) = (X, n+1) \cong (\Sigma X, n).$
- *Exact triangles*: $(X, n) \rightarrow (Y, n) \rightarrow (C_f, n) \rightarrow \Sigma(X, n)$ coming from cofiber sequences.

Stable homotopy

Recall that cohomology regards suspension as an invertible functor:

$$H^n(X) \cong H^{n+1}(\Sigma X).$$

Definition

The *compact stable homotopy category* \mathbf{SH}^c :

- *Objects*: (X, n) , X compact space, $n \in \mathbb{Z}$,

$$(X, n) \sim \Sigma^n X.$$

- *Morphisms*: $\mathrm{Hom}((X, n), (Y, m)) = \lim_{k \rightarrow \infty} [\Sigma^{k+n} X, \Sigma^{k+m} Y].$

- *Suspension*: $\Sigma(X, n) = (X, n+1) \cong (\Sigma X, n).$

- *Exact triangles*: $(X, n) \rightarrow (Y, n) \rightarrow (C_f, n) \rightarrow \Sigma(X, n)$ coming from cofiber sequences.

Stable homotopy

Recall that cohomology regards suspension as an invertible functor:

$$H^n(X) \cong H^{n+1}(\Sigma X).$$

Definition

The *compact stable homotopy category* \mathbf{SH}^c :

- *Objects*: (X, n) , X compact space, $n \in \mathbb{Z}$,

$$(X, n) \sim \Sigma^n X.$$

- *Morphisms*: $\mathrm{Hom}((X, n), (Y, m)) = \lim_{k \rightarrow \infty} [\Sigma^{k+n} X, \Sigma^{k+m} Y].$
- *Suspension*: $\Sigma(X, n) = (X, n+1) \cong (\Sigma X, n).$
- *Exact triangles*: $(X, n) \rightarrow (Y, n) \rightarrow (C_f, n) \rightarrow \Sigma(X, n)$ coming from cofiber sequences.

Triangulated categories

Definition

A *triangulated category* consists of:

- an additive category \mathbf{T} ,
- an equivalence $\Sigma: \mathbf{T} \xrightarrow{\sim} \mathbf{T}$,
- a family of *exact triangles* $X \xrightarrow{f} Y \rightarrow C_f \rightarrow \Sigma X$ in \mathbf{T} ,

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \swarrow \scriptstyle +1 & \searrow \\ & C_f & \end{array}$$

satisfying the formal properties of the compact stable homotopy category.

Triangulated categories

Definition

A *triangulated category* consists of:

- an additive category \mathbf{T} ,
- an equivalence $\Sigma: \mathbf{T} \xrightarrow{\sim} \mathbf{T}$,
- a family of *exact triangles* $X \xrightarrow{f} Y \rightarrow C_f \rightarrow \Sigma X$ in \mathbf{T} ,

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \swarrow & \searrow \\ & C_f & \end{array}$$

$+1$

satisfying the formal properties of the compact stable homotopy category.

Triangulated categories

Definition

A *triangulated category* consists of:

- an additive category \mathbf{T} ,
- an equivalence $\Sigma: \mathbf{T} \xrightarrow{\sim} \mathbf{T}$,
- a family of *exact triangles* $X \xrightarrow{f} Y \rightarrow C_f \rightarrow \Sigma X$ in \mathbf{T} ,

$$\begin{array}{ccccc} X & \xrightarrow{f} & Y & & \\ & \swarrow & \searrow & & \\ & C_f & & & \end{array}$$

$+1$

satisfying the formal properties of the compact stable homotopy category.

Triangulated categories

Definition

A *triangulated category* consists of:

- an additive category \mathbf{T} ,
- an equivalence $\Sigma: \mathbf{T} \xrightarrow{\sim} \mathbf{T}$,
- a family of *exact triangles* $X \xrightarrow{f} Y \rightarrow C_f \rightarrow \Sigma X$ in \mathbf{T} ,

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ & \swarrow \scriptstyle +1 & \searrow \\ & C_f & \end{array}$$

satisfying the formal properties of the compact stable homotopy category.

Homotopy and derived categories

Example

Let R be a ring.

- The *homotopy category* $\mathbf{K}(R)$, objects are *complexes* of R -modules,

$$C = \cdots \rightarrow C_{n+1} \xrightarrow{d_C} C_n \xrightarrow{d_C} C_{n-1} \rightarrow \cdots, \quad d_C^2 = 0,$$

$$(\Sigma C)_n = C_{n-1}, \quad d_{\Sigma C} = -d_C,$$

morphisms are chain homotopy classes of maps, and *exact triangles* come from *cofiber sequences* of complexes. 

- The *derived category* $\mathbf{D}(R) \subset \mathbf{K}(R)$ is the full subcategory spanned by ‘injective resolutions’ of complexes.
- For any Grothendieck abelian category \mathbf{A} we also have triangulated categories $\mathbf{D}(\mathbf{A}) \subset \mathbf{K}(\mathbf{A})$.

Homotopy and derived categories

Example

Let R be a ring.

- The *homotopy category* $\mathbf{K}(R)$, objects are *complexes* of R -modules,

$$C = \cdots \rightarrow C_{n+1} \xrightarrow{d_C} C_n \xrightarrow{d_C} C_{n-1} \rightarrow \cdots, \quad d_C^2 = 0,$$

$$(\Sigma C)_n = C_{n-1}, \quad d_{\Sigma C} = -d_C,$$

morphisms are chain homotopy classes of maps, and *exact triangles* come from *cofiber sequences* of complexes. ▶△

- The *derived category* $\mathbf{D}(R) \subset \mathbf{K}(R)$ is the full subcategory spanned by ‘injective resolutions’ of complexes.
- For any Grothendieck abelian category \mathbf{A} we also have triangulated categories $\mathbf{D}(\mathbf{A}) \subset \mathbf{K}(\mathbf{A})$.

Homotopy and derived categories

Example

Let R be a ring.

- The *homotopy category* $\mathbf{K}(R)$, objects are *complexes* of R -modules,

$$C = \cdots \rightarrow C_{n+1} \xrightarrow{d_C} C_n \xrightarrow{d_C} C_{n-1} \rightarrow \cdots, \quad d_C^2 = 0,$$

$$(\Sigma C)_n = C_{n-1}, \quad d_{\Sigma C} = -d_C,$$

morphisms are chain homotopy classes of maps, and *exact triangles* come from *cofiber sequences* of complexes. ▶△

- The *derived category* $\mathbf{D}(R) \subset \mathbf{K}(R)$ is the full subcategory spanned by ‘injective resolutions’ of complexes.
- For any Grothendieck abelian category \mathbf{A} we also have triangulated categories $\mathbf{D}(\mathbf{A}) \subset \mathbf{K}(\mathbf{A})$.

Example

Let R be a ring.

- The *homotopy category* $\mathbf{K}(R)$, objects are *complexes* of R -modules,

$$C = \cdots \rightarrow C_{n+1} \xrightarrow{d_C} C_n \xrightarrow{d_C} C_{n-1} \rightarrow \cdots, \quad d_C^2 = 0,$$

$$(\Sigma C)_n = C_{n-1}, \quad d_{\Sigma C} = -d_C,$$

morphisms are chain homotopy classes of maps, and *exact triangles* come from *cofiber sequences* of complexes. 

- The *derived category* $\mathbf{D}(R) \subset \mathbf{K}(R)$ is the full subcategory spanned by ‘injective resolutions’ of complexes.
- For any Grothendieck abelian category \mathbf{A} we also have triangulated categories $\mathbf{D}(\mathbf{A}) \subset \mathbf{K}(\mathbf{A})$.

Compactness in triangulated categories

Definition

A triangulated category \mathbf{T} is *cocomplete* if it has all *coproducts*,

$$\coprod_{i \in I} X_i, \quad I \text{ a set.}$$

An object Y in \mathbf{T} is *compact* if any map

$$Y \longrightarrow \coprod_{i \in I} X_i$$

factors as

$$Y \longrightarrow \coprod_{j \in J} X_j \subset \coprod_{i \in I} X_i, \quad I \supset J \text{ finite.}$$

Compactness in triangulated categories

Definition

A triangulated category \mathbf{T} is *cocomplete* if it has all *coproducts*,

$$\coprod_{i \in I} X_i, \quad I \text{ a set.}$$

An object Y in \mathbf{T} is *compact* if any map

$$Y \longrightarrow \coprod_{i \in I} X_i$$

factors as

$$Y \longrightarrow \coprod_{j \in J} X_j \subset \coprod_{i \in I} X_i, \quad I \supset J \text{ finite.}$$

Compactness in triangulated categories

Example

- $\mathbf{D}(\mathbf{A})$ and $\mathbf{K}(\mathbf{A})$ are cocomplete.
- Compact objects in $\mathbf{D}(R)$ are bounded complexes of finitely generated projective R -modules, a.k.a. *perfect complexes*.
- The compact stable homotopy category \mathbf{SH}^c admits a cocompletion \mathbf{SH} called the full *stable homotopy category*, whose objects are spectra.
- The derived category $\mathbf{D}(\text{sheaves}/M)$ of sheaves of abelian groups on an open connected manifold M of $\dim M \geq 1$ does *not* contain any non-trivial compact object [Neeman'01].

Compactness in triangulated categories

Example

- $\mathbf{D}(\mathbf{A})$ and $\mathbf{K}(\mathbf{A})$ are cocomplete.
- Compact objects in $\mathbf{D}(R)$ are bounded complexes of finitely generated projective R -modules, a.k.a. **perfect complexes**.
- The compact stable homotopy category \mathbf{SH}^c admits a cocompletion \mathbf{SH} called the full **stable homotopy category**, whose objects are spectra.
- The derived category $\mathbf{D}(\text{sheaves}/M)$ of sheaves of abelian groups on an open connected manifold M of $\dim M \geq 1$ does **not** contain any non-trivial compact object [Neeman'01].

Compactness in triangulated categories

Example

- $\mathbf{D}(\mathbf{A})$ and $\mathbf{K}(\mathbf{A})$ are cocomplete.
- Compact objects in $\mathbf{D}(R)$ are bounded complexes of finitely generated projective R -modules, a.k.a. *perfect complexes*.
- The compact stable homotopy category \mathbf{SH}^c admits a cocompletion \mathbf{SH} called the full *stable homotopy category*, whose objects are spectra.
- The derived category $\mathbf{D}(\text{sheaves}/M)$ of sheaves of abelian groups on an open connected manifold M of $\dim M \geq 1$ does *not* contain any non-trivial compact object [Neeman'01].

Example

- $\mathbf{D}(\mathbf{A})$ and $\mathbf{K}(\mathbf{A})$ are cocomplete.
- Compact objects in $\mathbf{D}(R)$ are bounded complexes of finitely generated projective R -modules, a.k.a. *perfect complexes*.
- The compact stable homotopy category \mathbf{SH}^c admits a cocompletion \mathbf{SH} called the full *stable homotopy category*, whose objects are spectra.
- The derived category $\mathbf{D}(\text{sheaves}/M)$ of sheaves of abelian groups on an open connected manifold M of $\dim M \geq 1$ does *not* contain any non-trivial compact object [Neeman'01].

Cohomology in triangulated categories

Definition

A *cohomology theory* H in a cocomplete triangulated category \mathbf{T} is an additive functor

$$H: \mathbf{T}^{op} \longrightarrow \mathbf{Ab}$$

to abelian groups taking exact triangles $X \xrightarrow{f} Y \rightarrow C_f \rightarrow \Sigma X$ in \mathbf{T} to exact sequences

$$H(X) \longleftarrow H(Y) \longleftarrow H(C_f) \longleftarrow H(\Sigma X),$$

and coproducts to products

$$H\left(\coprod_{i \in I} X_i\right) \cong \prod_{i \in I} H(X_i).$$

We set $H^n(X) = H(\Sigma^{-n} X)$.

Cohomology in triangulated categories

Definition

A *cohomology theory* H in a cocomplete triangulated category \mathbf{T} is an additive functor

$$H: \mathbf{T}^{op} \longrightarrow \mathbf{Ab}$$

to abelian groups taking exact triangles $X \xrightarrow{f} Y \rightarrow C_f \rightarrow \Sigma X$ in \mathbf{T} to exact sequences

$$H(X) \longleftarrow H(Y) \longleftarrow H(C_f) \longleftarrow H(\Sigma X),$$

and coproducts to products

$$H\left(\coprod_{i \in I} X_i\right) \cong \prod_{i \in I} H(X_i).$$

We set $H^n(X) = H(\Sigma^{-n} X)$.

Cohomology in triangulated categories

Definition

A *cohomology theory* H in a cocomplete triangulated category \mathbf{T} is an additive functor

$$H: \mathbf{T}^{op} \longrightarrow \mathbf{Ab}$$

to abelian groups taking exact triangles $X \xrightarrow{f} Y \rightarrow C_f \rightarrow \Sigma X$ in \mathbf{T} to exact sequences

$$H(X) \longleftarrow H(Y) \longleftarrow H(C_f) \longleftarrow H(\Sigma X),$$

and coproducts to products

$$H\left(\coprod_{i \in I} X_i\right) \cong \prod_{i \in I} H(X_i).$$

We set $H^n(X) = H(\Sigma^{-n} X)$.

Cohomology in triangulated categories

Definition

A *cohomology theory* H in a cocomplete triangulated category \mathbf{T} is an additive functor

$$H: \mathbf{T}^{op} \longrightarrow \mathbf{Ab}$$

to abelian groups taking exact triangles $X \xrightarrow{f} Y \rightarrow C_f \rightarrow \Sigma X$ in \mathbf{T} to exact sequences

$$H(X) \longleftarrow H(Y) \longleftarrow H(C_f) \longleftarrow H(\Sigma X),$$

and coproducts to products

$$H\left(\coprod_{i \in I} X_i\right) \cong \prod_{i \in I} H(X_i).$$

We set $H^n(X) = H(\Sigma^{-n}X)$.

Cohomology in triangulated categories

Example

- For any R -module A we have $H^n(-, A): \mathbf{K}(R)^{op} \rightarrow \mathbf{Ab}$,

$$H^n(C, A) = \frac{\text{cycles}}{\text{boundaries}} = \frac{\{\mathbf{c}: C_n \rightarrow A \mid 0 = fd_C: C_{n+1} \rightarrow C_n \rightarrow A\}}{\{\mathbf{b} = b'd_C: C_n \rightarrow C_{n-1} \rightarrow A\}}.$$

- Similarly we have $H^n(-, A): \mathbf{D}(R)^{op} \rightarrow \mathbf{Ab}$.
- Any object E in \mathbf{T} *represents* a cohomology theory $H: \mathbf{T}^{op} \rightarrow \mathbf{Ab}$ with

$$H(X) = \text{Hom}_{\mathbf{T}}(X, E).$$

Cohomology in triangulated categories

Example

- For any R -module A we have $H^n(-, A): \mathbf{K}(R)^{op} \rightarrow \mathbf{Ab}$,

$$H^n(C, A) = \frac{\text{cycles}}{\text{boundaries}} = \frac{\{\mathbf{c}: C_n \rightarrow A \mid 0 = fd_C: C_{n+1} \rightarrow C_n \rightarrow A\}}{\{\mathbf{b} = b'd_C: C_n \rightarrow C_{n-1} \rightarrow A\}}.$$

- Similarly we have $H^n(-, A): \mathbf{D}(R)^{op} \rightarrow \mathbf{Ab}$.
- Any object E in \mathbf{T} *represents* a cohomology theory $H: \mathbf{T}^{op} \rightarrow \mathbf{Ab}$ with

$$H(X) = \mathrm{Hom}_{\mathbf{T}}(X, E).$$

Cohomology in triangulated categories

Example

- For any R -module A we have $H^n(-, A): \mathbf{K}(R)^{op} \rightarrow \mathbf{Ab}$,

$$H^n(C, A) = \frac{\text{cycles}}{\text{boundaries}} = \frac{\{\mathbf{c}: C_n \rightarrow A \mid 0 = fd_C: C_{n+1} \rightarrow C_n \rightarrow A\}}{\{\mathbf{b} = b'd_C: C_n \rightarrow C_{n-1} \rightarrow A\}}.$$

- Similarly we have $H^n(-, A): \mathbf{D}(R)^{op} \rightarrow \mathbf{Ab}$.
- Any object E in \mathbf{T} **represents** a cohomology theory $H: \mathbf{T}^{op} \rightarrow \mathbf{Ab}$ with

$$H(X) = \text{Hom}_{\mathbf{T}}(X, E).$$

Cohomology in triangulated categories

Definition

A *cohomology theory* H in the full subcategory of *compact objects* $\mathbf{T}^c \subset \mathbf{T}$ is an additive functor

$$H: (\mathbf{T}^c)^{op} \longrightarrow \mathbf{Ab}$$

to abelian groups taking exact triangles $X \xrightarrow{f} Y \rightarrow C_f \rightarrow \Sigma X$ in \mathbf{T}^c to exact sequences

$$H(X) \longleftarrow H(Y) \longleftarrow H(C_f) \longleftarrow H(\Sigma X).$$

Example

Any object E in \mathbf{T} *represents* a cohomology theory $H: (\mathbf{T}^c)^{op} \rightarrow \mathbf{Ab}$ with

$$H(Y) = \mathrm{Hom}_{\mathbf{T}}(Y, E), \text{ for any } Y \text{ in } \mathbf{T}^c.$$

Cohomology in triangulated categories

Definition

A *cohomology theory* H in the full subcategory of *compact objects* $\mathbf{T}^c \subset \mathbf{T}$ is an additive functor

$$H: (\mathbf{T}^c)^{op} \longrightarrow \mathbf{Ab}$$

to abelian groups taking exact triangles $X \xrightarrow{f} Y \rightarrow C_f \rightarrow \Sigma X$ in \mathbf{T}^c to exact sequences

$$H(X) \longleftarrow H(Y) \longleftarrow H(C_f) \longleftarrow H(\Sigma X).$$

Example

Any object E in \mathbf{T} *represents* a cohomology theory $H: (\mathbf{T}^c)^{op} \rightarrow \mathbf{Ab}$ with

$$H(Y) = \mathrm{Hom}_{\mathbf{T}}(Y, E), \text{ for any } Y \text{ in } \mathbf{T}^c.$$

Compactly generated triangulated categories

Definition

A cocomplete triangulated category \mathbf{T} is **compactly generated** if any non-trivial object X in \mathbf{T} admits a non-trivial map $Y \rightarrow X$ from a compact object Y in \mathbf{T}^c .

Example

- The stable homotopy category \mathbf{SH} .
- The derived category of a ring $\mathbf{D}(R)$.
- The derived category $\mathbf{D}(\mathrm{Qcoh}/X)$ of complexes of quasi-coherent sheaves on a quasi-compact separated scheme X .
- $\mathbf{D}(\mathrm{sheaves}/M)$ is **not** compactly generated.
- $\mathbf{K}(\mathbb{Z})$ is **not** compactly generated.

Compactly generated triangulated categories

Definition

A cocomplete triangulated category \mathbf{T} is **compactly generated** if any non-trivial object X in \mathbf{T} admits a non-trivial map $Y \rightarrow X$ from a compact object Y in \mathbf{T}^c .

Example

- The stable homotopy category **SH**.
- The derived category of a ring $\mathbf{D}(R)$.
- The derived category $\mathbf{D}(\mathrm{Qcoh}/X)$ of complexes of quasi-coherent sheaves on a quasi-compact separated scheme X .
- $\mathbf{D}(\mathrm{sheaves}/M)$ is **not** compactly generated.
- $\mathbf{K}(\mathbb{Z})$ is **not** compactly generated.

Compactly generated triangulated categories

Definition

A cocomplete triangulated category \mathbf{T} is **compactly generated** if any non-trivial object X in \mathbf{T} admits a non-trivial map $Y \rightarrow X$ from a compact object Y in \mathbf{T}^c .

Example

- The stable homotopy category \mathbf{SH} .
- The derived category of a ring $\mathbf{D}(R)$.
- The derived category $\mathbf{D}(\mathrm{Qcoh}/X)$ of complexes of quasi-coherent sheaves on a quasi-compact separated scheme X .
- $\mathbf{D}(\mathrm{sheaves}/M)$ is **not** compactly generated.
- $\mathbf{K}(\mathbb{Z})$ is **not** compactly generated.

Compactly generated triangulated categories

Definition

A cocomplete triangulated category \mathbf{T} is *compactly generated* if any non-trivial object X in \mathbf{T} admits a non-trivial map $Y \rightarrow X$ from a compact object Y in \mathbf{T}^c .

Example

- The stable homotopy category \mathbf{SH} .
- The derived category of a ring $\mathbf{D}(R)$.
- The derived category $\mathbf{D}(\mathrm{Qcoh}/X)$ of complexes of quasi-coherent sheaves on a quasi-compact separated scheme X .
- $\mathbf{D}(\mathrm{sheaves}/M)$ is *not* compactly generated.
- $\mathbf{K}(\mathbb{Z})$ is *not* compactly generated.

Compactly generated triangulated categories

Definition

A cocomplete triangulated category \mathbf{T} is **compactly generated** if any non-trivial object X in \mathbf{T} admits a non-trivial map $Y \rightarrow X$ from a compact object Y in \mathbf{T}^c .

Example

- The stable homotopy category \mathbf{SH} .
- The derived category of a ring $\mathbf{D}(R)$.
- The derived category $\mathbf{D}(\mathrm{Qcoh}/X)$ of complexes of quasi-coherent sheaves on a quasi-compact separated scheme X .
- $\mathbf{D}(\mathrm{sheaves}/M)$ is **not** compactly generated.
- $\mathbf{K}(\mathbb{Z})$ is **not** compactly generated.

Compactly generated triangulated categories

Definition

A cocomplete triangulated category \mathbf{T} is **compactly generated** if any non-trivial object X in \mathbf{T} admits a non-trivial map $Y \rightarrow X$ from a compact object Y in \mathbf{T}^c .

Example

- The stable homotopy category \mathbf{SH} .
- The derived category of a ring $\mathbf{D}(R)$.
- The derived category $\mathbf{D}(\mathrm{Qcoh}/X)$ of complexes of quasi-coherent sheaves on a quasi-compact separated scheme X .
- $\mathbf{D}(\mathrm{sheaves}/M)$ is **not** compactly generated.
- $\mathbf{K}(\mathbb{Z})$ is **not** compactly generated.

Neeman's representability theorems

Theorem (Brown representability, Neeman'96)

Any cohomology theory $H: \mathbf{T}^{op} \rightarrow \mathbf{Ab}$ on a compactly generated triangulated category \mathbf{T} is representable.

Theorem (Adams representability, Neeman'97)

Let \mathbf{T} be a compactly generated triangulated category with \mathbf{T}^c countable.

- Any cohomology theory $H: (\mathbf{T}^c)^{op} \rightarrow \mathbf{Ab}$ is represented by a not necessarily compact object E in \mathbf{T} , $H = \mathrm{Hom}_{\mathbf{T}}(-, E)_{|\mathbf{T}^c}$.*
- Any natural transformation $\mathrm{Hom}_{\mathbf{T}}(-, E)_{|\mathbf{T}^c} \rightarrow \mathrm{Hom}_{\mathbf{T}}(-, E')_{|\mathbf{T}^c}$ is represented by a morphism $E \rightarrow E'$ in \mathbf{T} .*

Remark

This theorem applies to \mathbf{SH} , but it also applies to $\mathbf{D}(\mathbb{Z})$.

Neeman's representability theorems

Theorem (Brown representability, Neeman'96)

Any cohomology theory $H: \mathbf{T}^{op} \rightarrow \mathbf{Ab}$ on a compactly generated triangulated category \mathbf{T} is representable.

Theorem (Adams representability, Neeman'97)

Let \mathbf{T} be a compactly generated triangulated category with \mathbf{T}^c countable.

- Any cohomology theory $H: (\mathbf{T}^c)^{op} \rightarrow \mathbf{Ab}$ is represented by a not necessarily compact object E in \mathbf{T} , $H = \mathrm{Hom}_{\mathbf{T}}(-, E)|_{\mathbf{T}^c}$.*
- Any natural transformation $\mathrm{Hom}_{\mathbf{T}}(-, E)|_{\mathbf{T}^c} \rightarrow \mathrm{Hom}_{\mathbf{T}}(-, E')|_{\mathbf{T}^c}$ is represented by a morphism $E \rightarrow E'$ in \mathbf{T} .*

Remark

This theorem applies to \mathbf{SH} , but it also applies to $\mathbf{D}(\mathbb{Z})$.

Neeman's representability theorems

Theorem (Brown representability, Neeman'96)

Any cohomology theory $H: \mathbf{T}^{op} \rightarrow \mathbf{Ab}$ on a compactly generated triangulated category \mathbf{T} is representable.

Theorem (Adams representability, Neeman'97)

Let \mathbf{T} be a compactly generated triangulated category with \mathbf{T}^c countable.

- *Any cohomology theory $H: (\mathbf{T}^c)^{op} \rightarrow \mathbf{Ab}$ is represented by a not necessarily compact object E in \mathbf{T} , $H = \mathrm{Hom}_{\mathbf{T}}(-, E)|_{\mathbf{T}^c}$.*
- *Any natural transformation $\mathrm{Hom}_{\mathbf{T}}(-, E)|_{\mathbf{T}^c} \rightarrow \mathrm{Hom}_{\mathbf{T}}(-, E')|_{\mathbf{T}^c}$ is represented by a morphism $E \rightarrow E'$ in \mathbf{T} .*

Remark

This theorem applies to \mathbf{SH} , but it also applies to $\mathbf{D}(\mathbb{Z})$.

Neeman's representability theorems

Theorem (Brown representability, Neeman'96)

Any cohomology theory $H: \mathbf{T}^{op} \rightarrow \mathbf{Ab}$ on a compactly generated triangulated category \mathbf{T} is representable.

Theorem (Adams representability, Neeman'97)

Let \mathbf{T} be a compactly generated triangulated category with \mathbf{T}^c countable.

- *Any cohomology theory $H: (\mathbf{T}^c)^{op} \rightarrow \mathbf{Ab}$ is represented by a not necessarily compact object E in \mathbf{T} , $H = \mathrm{Hom}_{\mathbf{T}}(-, E)_{|\mathbf{T}^c}$.*
- *Any natural transformation $\mathrm{Hom}_{\mathbf{T}}(-, E)_{|\mathbf{T}^c} \rightarrow \mathrm{Hom}_{\mathbf{T}}(-, E')_{|\mathbf{T}^c}$ is represented by a morphism $E \rightarrow E'$ in \mathbf{T} .*

Remark

This theorem applies to \mathbf{SH} , but it also applies to $\mathbf{D}(\mathbb{Z})$.

Neeman's representability theorems

Theorem (Brown representability, Neeman'96)

Any cohomology theory $H: \mathbf{T}^{op} \rightarrow \mathbf{Ab}$ on a compactly generated triangulated category \mathbf{T} is representable.

Theorem (Adams representability, Neeman'97)

Let \mathbf{T} be a compactly generated triangulated category with \mathbf{T}^c countable.

- *Any cohomology theory $H: (\mathbf{T}^c)^{op} \rightarrow \mathbf{Ab}$ is represented by a not necessarily compact object E in \mathbf{T} , $H = \mathrm{Hom}_{\mathbf{T}}(-, E)_{|\mathbf{T}^c}$.*
- *Any natural transformation $\mathrm{Hom}_{\mathbf{T}}(-, E)_{|\mathbf{T}^c} \rightarrow \mathrm{Hom}_{\mathbf{T}}(-, E')_{|\mathbf{T}^c}$ is represented by a morphism $E \rightarrow E'$ in \mathbf{T} .*

Remark

This theorem applies to \mathbf{SH} , but it also applies to $\mathbf{D}(\mathbb{Z})$.

Counterexamples to Adams representability

Example (Casacuberta–Neeman'09)

Let R be a 'polynomial ring with a proper class of indeterminates', the subcategory of **acyclic complexes** in $\mathbf{K}(R)$ does **not** satisfy Brown representability.

Example (Christensen–Keller–Neeman'01)

- Whether Adams representability holds in the derived category of the complex affine plane $\mathbf{D}(\mathrm{Qcoh}/\mathbb{A}_{\mathbb{C}}^2)$ depends essentially on the **continuum hypothesis**.
- Adams representability does **not** hold in $\mathbf{D}(k[x, y])$ provided $\mathrm{card} k \geq \aleph_3$.

► transfinite

Counterexamples to Adams representability

Example (Casacuberta–Neeman'09)

Let R be a 'polynomial ring with a proper class of indeterminates', the subcategory of **acyclic complexes** in $\mathbf{K}(R)$ does **not** satisfy Brown representability.

Example (Christensen–Keller–Neeman'01)

- Whether Adams representability holds in the derived category of the complex affine plane $\mathbf{D}(\mathrm{Qcoh}/\mathbb{A}_{\mathbb{C}}^2)$ depends essentially on the **continuum hypothesis**.
- Adams representability does **not** hold in $\mathbf{D}(k[x, y])$ provided $\mathrm{card} k \geq \aleph_3$.

► transfinite

Counterexamples to Adams representability

Example (Casacuberta–Neeman'09)

Let R be a 'polynomial ring with a proper class of indeterminates', the subcategory of **acyclic complexes** in $\mathbf{K}(R)$ does **not** satisfy Brown representability.

Example (Christensen–Keller–Neeman'01)

- Whether Adams representability holds in the derived category of the complex affine plane $\mathbf{D}(\mathrm{Qcoh}/\mathbb{A}_{\mathbb{C}}^2)$ depends essentially on the **continuum hypothesis**.
- Adams representability does **not** hold in $\mathbf{D}(k[x, y])$ provided $\mathrm{card} k \geq \aleph_3$.

► transfinite

A homological criterion for Adams representability

Definition

A right \mathbf{T}^c -module A is an additive functor $A: (\mathbf{T}^c)^{op} \rightarrow \mathbf{Ab}$. The category $\mathbf{Mod}(\mathbf{T}^c)$ of right \mathbf{T}^c -modules is a Grothendieck abelian category with a set of small projective generators.

Theorem (Neeman'97, Beligiannis'00)

If \mathbf{T} is compactly generated and $H: (\mathbf{T}^c)^{op} \rightarrow \mathbf{Ab}$ is a cohomology theory then:

- $\text{proj dim } H \leq 2 \Rightarrow H$ is representable.
- $\text{proj dim } H \leq 1$ for all $H \Leftrightarrow$ Adams representability theorem for \mathbf{T} .

A homological criterion for Adams representability

Definition

A right \mathbf{T}^c -module A is an additive functor $A: (\mathbf{T}^c)^{op} \rightarrow \mathbf{Ab}$. The category $\mathbf{Mod}(\mathbf{T}^c)$ of right \mathbf{T}^c -modules is a Grothendieck abelian category with a set of small projective generators.

Theorem (Neeman'97, Beligiannis'00)

If \mathbf{T} is compactly generated and $H: (\mathbf{T}^c)^{op} \rightarrow \mathbf{Ab}$ is a cohomology theory then:

- $\text{proj dim } H \leq 2 \Rightarrow H$ is representable.
- $\text{proj dim } H \leq 1$ for all $H \Leftrightarrow$ Adams representability theorem for \mathbf{T} .

A homological criterion for Adams representability

Definition

A right \mathbf{T}^c -module A is an additive functor $A: (\mathbf{T}^c)^{op} \rightarrow \mathbf{Ab}$. The category $\mathbf{Mod}(\mathbf{T}^c)$ of right \mathbf{T}^c -modules is a Grothendieck abelian category with a set of small projective generators.

Theorem (Neeman'97, Beligiannis'00)

If \mathbf{T} is compactly generated and $H: (\mathbf{T}^c)^{op} \rightarrow \mathbf{Ab}$ is a cohomology theory then:

- $\text{proj dim } H \leq 2 \Rightarrow H$ is representable.
- $\text{proj dim } H \leq 1$ for all $H \Leftrightarrow$ Adams representability theorem for \mathbf{T} .

A homological criterion for Adams representability

Definition

A right \mathbf{T}^c -module A is an additive functor $A: (\mathbf{T}^c)^{op} \rightarrow \mathbf{Ab}$. The category $\mathbf{Mod}(\mathbf{T}^c)$ of right \mathbf{T}^c -modules is a Grothendieck abelian category with a set of small projective generators.

Theorem (Neeman'97, Beligiannis'00)

If \mathbf{T} is compactly generated and $H: (\mathbf{T}^c)^{op} \rightarrow \mathbf{Ab}$ is a cohomology theory then:

- $\text{proj dim } H \leq 2 \Rightarrow H$ is representable.
- $\text{proj dim } H \leq 1$ for all $H \Leftrightarrow$ Adams representability theorem for \mathbf{T} .

Purity is the relative homological algebra in $\mathbf{Mod}(R)$ obtained by regarding all finitely presented R -modules as projectives.

Proposition (Christensen–Keller–Neeman'01)

For any ring R ,

$$\sup_{\substack{H: (\mathbf{D}(R)^c)^{op} \rightarrow \mathbf{Ab} \\ \text{cohomology}}} \text{proj dim } H \geq \text{pure proj dim } \mathbf{Mod}(R).$$

Remark (Baer–Brune–Lenzing'82)

*If R is a finite-dimensional hereditary algebra over an uncountable algebraically closed field, then $\mathbf{D}(R)$ satisfies the Adams representability theorem \Leftrightarrow has **finite representation type**.*

Purity is the relative homological algebra in $\mathbf{Mod}(R)$ obtained by regarding all finitely presented R -modules as projectives.

Proposition (Christensen–Keller–Neeman'01)

For any ring R ,

$$\sup_{\substack{H: (\mathbf{D}(R)^c)^{op} \rightarrow \mathbf{Ab} \\ \text{cohomology}}} \text{proj dim } H \geq \text{pure proj dim } \mathbf{Mod}(R).$$

Remark (Baer–Brune–Lenzing'82)

If R is a finite-dimensional hereditary algebra over an uncountable algebraically closed field, then $\mathbf{D}(R)$ satisfies the Adams representability theorem \Leftrightarrow has *finite representation type*.

Purity is the relative homological algebra in $\mathbf{Mod}(R)$ obtained by regarding all finitely presented R -modules as projectives.

Proposition (Christensen–Keller–Neeman'01)

For any ring R ,

$$\sup_{\substack{H: (\mathbf{D}(R)^c)^{op} \rightarrow \mathbf{Ab} \\ \text{cohomology}}} \text{proj dim } H \geq \text{pure proj dim } \mathbf{Mod}(R).$$

Remark (Baer–Brune–Lenzing'82)

*If R is a finite-dimensional hereditary algebra over an uncountable algebraically closed field, then $\mathbf{D}(R)$ satisfies the Adams representability theorem \Leftrightarrow has **finite representation type**.*

Transfinite compactness in triangulated categories

Definition

Let \mathbf{T} be a cocomplete triangulated category. For any **regular cardinal** α there is a notion of **α -compact** object. These objects span a full triangulated subcategory $\mathbf{T}^\alpha \subset \mathbf{T}$. For $\alpha = \aleph_0$ this coincides with classical compactness.

If Y is in \mathbf{T}^α then any map

$$Y \longrightarrow \coprod_{i \in I} X_i$$

factors as

$$Y \longrightarrow \coprod_{j \in J} X_j \subset \coprod_{i \in I} X_i, \quad I \supset J, \text{ card } J < \alpha.$$

Transfinite compactness in triangulated categories

Definition

Let \mathbf{T} be a cocomplete triangulated category. For any **regular cardinal** α there is a notion of **α -compact** object. These objects span a full triangulated subcategory $\mathbf{T}^\alpha \subset \mathbf{T}$. For $\alpha = \aleph_0$ this coincides with classical compactness.

If Y is in \mathbf{T}^α then any map

$$Y \longrightarrow \coprod_{i \in I} X_i$$

factors as

$$Y \longrightarrow \coprod_{j \in J} X_j \subset \coprod_{i \in I} X_i, \quad I \supset J, \text{ card } J < \alpha.$$

Example

Let $\alpha > \aleph_0$.

- A spectrum E in **SH** is α -compact whenever $\text{card } \pi_n(E) < \alpha$ for all $n \in \mathbb{Z}$.
- Given a ring R , either noetherian or with $\text{card } R < \alpha$, α -compact objects in $\mathbf{D}(R)$ are complexes C such that $\text{card } H_n(C)$ has $< \alpha$ generators for all $n \in \mathbb{Z}$.

Example

Let $\alpha > \aleph_0$.

- A spectrum E in **SH** is α -compact whenever $\text{card } \pi_n(E) < \alpha$ for all $n \in \mathbb{Z}$.
- Given a ring R , either noetherian or with $\text{card } R < \alpha$, α -compact objects in **D**(R) are complexes C such that $\text{card } H_n(C)$ has $< \alpha$ generators for all $n \in \mathbb{Z}$.

Well generated categories

Definition

A cocomplete triangulated category \mathbf{T} is α -compactly generated if any non-trivial object X in \mathbf{T} admits a non-trivial map $Y \rightarrow X$ from an α -compact object Y in \mathbf{T}^α . A cocomplete triangulated category is *well generated* if it is α -compactly generated for some regular cardinal α .

Example

- All compactly generated triangulated categories are well generated.
- $\mathbf{D}(\text{sheaves}/M)$ is well generated, actually \aleph_1 -compactly generated.
- $\mathbf{K}(\mathbb{Z})$ is *not* well generated.
- If \mathbf{T} is well generated then \mathbf{T}^{op} is *never* well generated.

Well generated categories

Definition

A cocomplete triangulated category \mathbf{T} is α -compactly generated if any non-trivial object X in \mathbf{T} admits a non-trivial map $Y \rightarrow X$ from an α -compact object Y in \mathbf{T}^α . A cocomplete triangulated category is *well generated* if it is α -compactly generated for some regular cardinal α .

Example

- All compactly generated triangulated categories are well generated.
- $\mathbf{D}(\text{sheaves}/M)$ is well generated, actually \aleph_1 -compactly generated.
- $\mathbf{K}(\mathbb{Z})$ is *not* well generated.
- If \mathbf{T} is well generated then \mathbf{T}^{op} is *never* well generated.

Well generated categories

Definition

A cocomplete triangulated category \mathbf{T} is α -compactly generated if any non-trivial object X in \mathbf{T} admits a non-trivial map $Y \rightarrow X$ from an α -compact object Y in \mathbf{T}^α . A cocomplete triangulated category is *well generated* if it is α -compactly generated for some regular cardinal α .

Example

- All compactly generated triangulated categories are well generated.
- $\mathbf{D}(\text{sheaves}/M)$ is well generated, actually \aleph_1 -compactly generated.
- $\mathbf{K}(\mathbb{Z})$ is *not* well generated.
- If \mathbf{T} is well generated then \mathbf{T}^{op} is *never* well generated.

Well generated categories

Definition

A cocomplete triangulated category \mathbf{T} is α -compactly generated if any non-trivial object X in \mathbf{T} admits a non-trivial map $Y \rightarrow X$ from an α -compact object Y in \mathbf{T}^α . A cocomplete triangulated category is *well generated* if it is α -compactly generated for some regular cardinal α .

Example

- All compactly generated triangulated categories are well generated.
- $\mathbf{D}(\text{sheaves}/M)$ is well generated, actually \aleph_1 -compactly generated.
- $\mathbf{K}(\mathbb{Z})$ is *not* well generated.
- If \mathbf{T} is well generated then \mathbf{T}^{op} is *never* well generated.

Well generated categories

Definition

A cocomplete triangulated category \mathbf{T} is α -compactly generated if any non-trivial object X in \mathbf{T} admits a non-trivial map $Y \rightarrow X$ from an α -compact object Y in \mathbf{T}^α . A cocomplete triangulated category is *well generated* if it is α -compactly generated for some regular cardinal α .

Example

- All compactly generated triangulated categories are well generated.
- $\mathbf{D}(\text{sheaves}/M)$ is well generated, actually \aleph_1 -compactly generated.
- $\mathbf{K}(\mathbb{Z})$ is *not* well generated.
- If \mathbf{T} is well generated then \mathbf{T}^{op} is *never* well generated.

Representability in well generated categories

Theorem (Brown representability, Neeman'01)

Any cohomology theory $H: \mathbf{T}^{op} \rightarrow \mathbf{Ab}$ on a well generated triangulated category \mathbf{T} is representable.

Definition

A *cohomology theory* H for α -compact objects is an additive functor

$$H: (\mathbf{T}^\alpha)^{op} \longrightarrow \mathbf{Ab}$$

taking exact triangles to exact sequences and

$$H\left(\coprod_{i \in I} X_i\right) \cong \prod_{i \in I} H(X_i), \quad \text{card } I < \alpha.$$

Representability in well generated categories

Theorem (Brown representability, Neeman'01)

Any cohomology theory $H: \mathbf{T}^{op} \rightarrow \mathbf{Ab}$ on a well generated triangulated category \mathbf{T} is representable.

Definition

A *cohomology theory* H for α -compact objects is an additive functor

$$H: (\mathbf{T}^\alpha)^{op} \longrightarrow \mathbf{Ab}$$

taking exact triangles to exact sequences and

$$H\left(\coprod_{i \in I} X_i\right) \cong \prod_{i \in I} H(X_i), \quad \text{card } I < \alpha.$$

Representability in well generated categories

Theorem (Brown representability, Neeman'01)

Any cohomology theory $H: \mathbf{T}^{op} \rightarrow \mathbf{Ab}$ on a well generated triangulated category \mathbf{T} is representable.

Definition

A *cohomology theory* H for α -compact objects is an additive functor

$$H: (\mathbf{T}^\alpha)^{op} \longrightarrow \mathbf{Ab}$$

taking exact triangles to exact sequences and

$$H\left(\coprod_{i \in I} X_i\right) \cong \prod_{i \in I} H(X_i), \quad \text{card } I < \alpha.$$

What about α -Adams representability?

Representability in well generated categories

Conjecture (Rosický'05, Neeman'09)

\mathbf{T} is a well generated triangulated category \Leftrightarrow there exists a regular cardinal α such that the α -Adams representability theorem holds:

- Any cohomology theory $H: (\mathbf{T}^\alpha)^{op} \rightarrow \mathbf{Ab}$ is represented by a not necessarily α -compact object E in \mathbf{T} , $H = \text{Hom}_{\mathbf{T}}(-, E)_{|\mathbf{T}^\alpha}$.
- Any natural transformation $\text{Hom}_{\mathbf{T}}(-, E)_{|\mathbf{T}^\alpha} \rightarrow \text{Hom}_{\mathbf{T}}(-, E')_{|\mathbf{T}^\alpha}$ is represented by a morphism $E \rightarrow E'$ in \mathbf{T} .

Proposition (The 'easy' part, Rosicky'09)

\Leftarrow is true, in particular Brown representability follows from α -Adams representability.

Almost **nothing** is known about \Rightarrow for $\alpha > \aleph_0$.

► remark

► theorem

Representability in well generated categories

Conjecture (Rosický'05, Neeman'09)

\mathbf{T} is a well generated triangulated category \Leftrightarrow there exists a regular cardinal α such that the α -Adams representability theorem holds:

- Any cohomology theory $H: (\mathbf{T}^\alpha)^{op} \rightarrow \mathbf{Ab}$ is represented by a not necessarily α -compact object E in \mathbf{T} , $H = \text{Hom}_{\mathbf{T}}(-, E)_{|\mathbf{T}^\alpha}$.
- Any natural transformation $\text{Hom}_{\mathbf{T}}(-, E)_{|\mathbf{T}^\alpha} \rightarrow \text{Hom}_{\mathbf{T}}(-, E')_{|\mathbf{T}^\alpha}$ is represented by a morphism $E \rightarrow E'$ in \mathbf{T} .

Proposition (The 'easy' part, Rosický'09)

\Leftarrow is true, in particular Brown representability follows from α -Adams representability.

Almost **nothing** is known about \Rightarrow for $\alpha > \aleph_0$.

► remark

► theorem

Representability in well generated categories

Conjecture (Rosický'05, Neeman'09)

\mathbf{T} is a well generated triangulated category \Leftrightarrow there exists a regular cardinal α such that the α -Adams representability theorem holds:

- Any cohomology theory $H: (\mathbf{T}^\alpha)^{op} \rightarrow \mathbf{Ab}$ is represented by a not necessarily α -compact object E in \mathbf{T} , $H = \mathrm{Hom}_{\mathbf{T}}(-, E)_{|\mathbf{T}^\alpha}$.
- Any natural transformation $\mathrm{Hom}_{\mathbf{T}}(-, E)_{|\mathbf{T}^\alpha} \rightarrow \mathrm{Hom}_{\mathbf{T}}(-, E')_{|\mathbf{T}^\alpha}$ is represented by a morphism $E \rightarrow E'$ in \mathbf{T} .

Proposition (The 'easy' part, Rosický'09)

\Leftarrow is true, in particular Brown representability follows from α -Adams representability.

Almost **nothing** is known about \Rightarrow for $\alpha > \aleph_0$.

► remark

► theorem

Representability in well generated categories

Conjecture (Rosický'05, Neeman'09)

\mathbf{T} is a well generated triangulated category \Leftrightarrow there exists a regular cardinal α such that the α -Adams representability theorem holds:

- Any cohomology theory $H: (\mathbf{T}^\alpha)^{op} \rightarrow \mathbf{Ab}$ is represented by a not necessarily α -compact object E in \mathbf{T} , $H = \mathrm{Hom}_{\mathbf{T}}(-, E)|_{\mathbf{T}^\alpha}$.
- Any natural transformation $\mathrm{Hom}_{\mathbf{T}}(-, E)|_{\mathbf{T}^\alpha} \rightarrow \mathrm{Hom}_{\mathbf{T}}(-, E')|_{\mathbf{T}^\alpha}$ is represented by a morphism $E \rightarrow E'$ in \mathbf{T} .

Proposition (The 'easy' part, Rosicky'09)

\Leftarrow is true, in particular Brown representability follows from α -Adams representability.

Almost **nothing** is known about \Rightarrow for $\alpha > \aleph_0$.

► remark

► theorem

Representability in well generated categories

Conjecture (Rosický'05, Neeman'09)

\mathbf{T} is a well generated triangulated category \Leftrightarrow there exists a regular cardinal α such that the α -Adams representability theorem holds:

- Any cohomology theory $H: (\mathbf{T}^\alpha)^{op} \rightarrow \mathbf{Ab}$ is represented by a not necessarily α -compact object E in \mathbf{T} , $H = \mathrm{Hom}_{\mathbf{T}}(-, E)_{|\mathbf{T}^\alpha}$.
- Any natural transformation $\mathrm{Hom}_{\mathbf{T}}(-, E)_{|\mathbf{T}^\alpha} \rightarrow \mathrm{Hom}_{\mathbf{T}}(-, E')_{|\mathbf{T}^\alpha}$ is represented by a morphism $E \rightarrow E'$ in \mathbf{T} .

Proposition (The 'easy' part, Rosicky'09)

\Leftarrow is true, in particular Brown representability follows from α -Adams representability.

Almost **nothing** is known about \Rightarrow for $\alpha > \aleph_0$.

► remark

► theorem

A homological criterion for Adams representability

Definition

A right **continuous \mathbf{T}^α -module** A is an additive functor $A: (\mathbf{T}^\alpha)^{op} \rightarrow \mathbf{Ab}$ with

$$A\left(\coprod_{i \in I} X_i\right) \cong \prod_{i \in I} A(X_i), \quad \text{card } I < \alpha.$$

The category $\mathbf{Mod}_\alpha(\mathbf{T}^\alpha)$ of right continuous \mathbf{T}^α -modules is an abelian category but not Grothendieck.

Theorem (M-Raventós'09)

If \mathbf{T} is α -compactly generated and $H: (\mathbf{T}^\alpha)^{op} \rightarrow \mathbf{Ab}$ is a cohomology theory then:

- $\text{proj dim } H \leq 2 \Rightarrow H$ is representable.
- $\text{proj dim } H \leq 1$ for all $H \Leftrightarrow \alpha$ -Adams representability theorem for \mathbf{T} .

A homological criterion for Adams representability

Definition

A right **continuous \mathbf{T}^α -module** A is an additive functor $A: (\mathbf{T}^\alpha)^{op} \rightarrow \mathbf{Ab}$ with

$$A\left(\coprod_{i \in I} X_i\right) \cong \prod_{i \in I} A(X_i), \quad \text{card } I < \alpha.$$

The category $\mathbf{Mod}_\alpha(\mathbf{T}^\alpha)$ of right continuous \mathbf{T}^α -modules is an abelian category but not Grothendieck.

Theorem (M-Raventós'09)

If \mathbf{T} is α -compactly generated and $H: (\mathbf{T}^\alpha)^{op} \rightarrow \mathbf{Ab}$ is a cohomology theory then:

- $\text{proj dim } H \leq 2 \Rightarrow H$ is representable.
- $\text{proj dim } H \leq 1$ for all $H \Leftrightarrow \alpha$ -Adams representability theorem for \mathbf{T} .

A homological criterion for Adams representability

Definition

A right **continuous \mathbf{T}^α -module** A is an additive functor $A: (\mathbf{T}^\alpha)^{op} \rightarrow \mathbf{Ab}$ with

$$A\left(\coprod_{i \in I} X_i\right) \cong \prod_{i \in I} A(X_i), \quad \text{card } I < \alpha.$$

The category $\mathbf{Mod}_\alpha(\mathbf{T}^\alpha)$ of right continuous \mathbf{T}^α -modules is an abelian category but not Grothendieck.

Theorem (M-Raventós'09)

If \mathbf{T} is α -compactly generated and $H: (\mathbf{T}^\alpha)^{op} \rightarrow \mathbf{Ab}$ is a cohomology theory then:

- $\text{proj dim } H \leq 2 \Rightarrow H$ is representable.
- $\text{proj dim } H \leq 1$ for all $H \Leftrightarrow \alpha$ -Adams representability theorem for \mathbf{T} .

A homological criterion for Adams representability

Definition

A right **continuous \mathbf{T}^α -module** A is an additive functor $A: (\mathbf{T}^\alpha)^{op} \rightarrow \mathbf{Ab}$ with

$$A\left(\coprod_{i \in I} X_i\right) \cong \prod_{i \in I} A(X_i), \quad \text{card } I < \alpha.$$

The category $\mathbf{Mod}_\alpha(\mathbf{T}^\alpha)$ of right continuous \mathbf{T}^α -modules is an abelian category but not Grothendieck.

Theorem (M-Raventós'09)

If \mathbf{T} is α -compactly generated and $H: (\mathbf{T}^\alpha)^{op} \rightarrow \mathbf{Ab}$ is a cohomology theory then:

- $\text{proj dim } H \leq 2 \Rightarrow H$ is representable.
- $\text{proj dim } H \leq 1$ for all $H \Leftrightarrow \alpha$ -Adams representability theorem for \mathbf{T} .

Transfinite purity

α -purity is the relative homological algebra in $\mathbf{Mod}(R)$ obtained by regarding all R -modules with $< \alpha$ generators and relations as projectives.

Proposition (M-Raventós'09)

For any ring R ,

$$\sup_{\substack{H: (\mathbf{D}(R)^\alpha)^{op} \rightarrow \mathbf{Ab} \\ \text{cohomology}}} \text{proj dim } H \geq \alpha\text{-pure proj dim } \mathbf{Mod}(R).$$

Proposition (M-Raventós'09)

If R is a finite-dimensional *wild* hereditary k -algebra, $\text{card } k \geq \aleph_\omega$, then the α -Adams representability theorem is *false* for all $\alpha < \aleph_\omega$.

► skip remark

Transfinite purity

α -purity is the relative homological algebra in $\mathbf{Mod}(R)$ obtained by regarding all R -modules with $< \alpha$ generators and relations as projectives.

Proposition (M-Raventós'09)

For any ring R ,

$$\sup_{\substack{H: (\mathbf{D}(R)^\alpha)^{op} \rightarrow \mathbf{Ab} \\ \text{cohomology}}} \text{proj dim } H \geq \alpha\text{-pure proj dim } \mathbf{Mod}(R).$$

Proposition (M-Raventós'09)

If R is a finite-dimensional *wild* hereditary k -algebra, $\text{card } k \geq \aleph_\omega$, then the α -Adams representability theorem is *false* for all $\alpha < \aleph_\omega$.

► skip remark

Transfinite purity

α -purity is the relative homological algebra in $\mathbf{Mod}(R)$ obtained by regarding all R -modules with $< \alpha$ generators and relations as projectives.

Proposition (M-Raventós'09)

For any ring R ,

$$\sup_{\substack{H: (\mathbf{D}(R)^\alpha)^{op} \rightarrow \mathbf{Ab} \\ \text{cohomology}}} \text{proj dim } H \geq \alpha\text{-pure proj dim } \mathbf{Mod}(R).$$

Proposition (M-Raventós'09)

If R is a finite-dimensional *wild* hereditary k -algebra, $\text{card } k \geq \aleph_\omega$, then the α -Adams representability theorem is *false* for all $\alpha < \aleph_\omega$.

► skip remark

Remark

If the conjecture is true, then for any ring R there exists a cardinal α such that for any R -module A there is a short exact sequence,

$$0 \rightarrow \bigoplus_{i \in I} P_i \longrightarrow \bigoplus_{j \in J} Q_j \longrightarrow A \rightarrow 0,$$

such that the modules P_i and Q_j have $< \alpha$ generators and relations.

Moreover, if B is an R -module with $< \alpha$ generators and relations, then

A commutative diagram illustrating the relationship between modules B , A , and $\bigoplus_{j \in J} Q_j$. The diagram consists of three nodes: B at the top, $\bigoplus_{j \in J} Q_j$ at the bottom left, and A at the bottom right. Arrows connect these nodes: a vertical arrow from B down to A labeled φ ; a horizontal arrow from $\bigoplus_{j \in J} Q_j$ to A ; and a diagonal arrow from B down to $\bigoplus_{j \in J} Q_j$ labeled \exists . There is also a small arrow pointing from the diagonal arrow to the horizontal arrow.

Remark

If the conjecture is true, then for any ring R there exists a cardinal α such that for any R -module A there is a short exact sequence,

$$0 \rightarrow \bigoplus_{i \in I} P_i \longrightarrow \bigoplus_{j \in J} Q_j \longrightarrow A \rightarrow 0,$$

such that the modules P_i and Q_j have $< \alpha$ generators and relations.

Moreover, if B is an R -module with $< \alpha$ generators and relations, then

$$\begin{array}{ccc} & & B \\ & \exists \swarrow & \downarrow \forall \\ & \nwarrow & \\ \bigoplus_{j \in J} Q_j & \longrightarrow & A \end{array}$$

Adams representability for $\alpha = \aleph_1$

Under the continuum hypothesis, the following theorem can be applied to $\mathbf{D}(\text{sheaves}/M)$.

Theorem (\aleph_1 -Adams representability, M-Raventós'09)

Let \mathbf{T} be an \aleph_1 -generated triangulated category with $\text{card } \mathbf{T}^{\aleph_1} \leq \aleph_1$.

Then:

- Any cohomology theory $H: (\mathbf{T}^{\aleph_1})^{op} \rightarrow \mathbf{Ab}$ is represented by a not necessarily \aleph_1 -compact object E in \mathbf{T} , $H = \text{Hom}_{\mathbf{T}}(-, E)|_{\mathbf{T}^{\aleph_1}}$.

This is the only available non-trivial result on α -Adams representability for $\alpha > \aleph_0$.

Adams representability for $\alpha = \aleph_1$

Under the continuum hypothesis, the following theorem can be applied to $\mathbf{D}(\text{sheaves}/M)$.

Theorem (\aleph_1 -Adams representability, M-Raventós'09)

Let \mathbf{T} be an \aleph_1 -generated triangulated category with $\text{card } \mathbf{T}^{\aleph_1} \leq \aleph_1$.

Then:

- Any cohomology theory $H: (\mathbf{T}^{\aleph_1})^{op} \rightarrow \mathbf{Ab}$ is represented by a not necessarily \aleph_1 -compact object E in \mathbf{T} , $H = \text{Hom}_{\mathbf{T}}(-, E)|_{\mathbf{T}^{\aleph_1}}$.

This is the only available non-trivial result on α -Adams representability for $\alpha > \aleph_0$.

Adams representability for $\alpha = \aleph_1$

Under the continuum hypothesis, the following theorem can be applied to $\mathbf{D}(\text{sheaves}/M)$.

Theorem (\aleph_1 -Adams representability, M-Raventós'09)

Let \mathbf{T} be an \aleph_1 -generated triangulated category with $\text{card } \mathbf{T}^{\aleph_1} \leq \aleph_1$.

Then:

- Any cohomology theory $H: (\mathbf{T}^{\aleph_1})^{op} \rightarrow \mathbf{Ab}$ is represented by a not necessarily \aleph_1 -compact object E in \mathbf{T} , $H = \text{Hom}_{\mathbf{T}}(-, E)|_{\mathbf{T}^{\aleph_1}}$.

This is the only available non-trivial result on α -Adams representability for $\alpha > \aleph_0$.

Adams representability for $\alpha = \aleph_1$

Under the continuum hypothesis, the following theorem can be applied to $\mathbf{D}(\text{sheaves}/M)$.

Theorem (\aleph_1 -Adams representability, M-Raventós'09)

Let \mathbf{T} be an \aleph_1 -generated triangulated category with $\text{card } \mathbf{T}^{\aleph_1} \leq \aleph_1$.

Then:

- Any cohomology theory $H: (\mathbf{T}^{\aleph_1})^{op} \rightarrow \mathbf{Ab}$ is represented by a not necessarily \aleph_1 -compact object E in \mathbf{T} , $H = \text{Hom}_{\mathbf{T}}(-, E)|_{\mathbf{T}^{\aleph_1}}$.

This is the only available non-trivial result on α -Adams representability for $\alpha > \aleph_0$.

Representability of cohomology theories

Fernando Muro

Universidad de Sevilla
Departamento de Álgebra

Joint Mathematical Conference CSASC 2010
Prague, January 2010

Cofiber sequence of complexes

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \vdots & & \vdots \\ \downarrow & & \downarrow \\ X_{n+1} & \xrightarrow{f} & Y_{n+1} \\ \downarrow d_X & & \downarrow d_Y \\ X_n & \xrightarrow{f} & Y_n \\ \downarrow d_X & & \downarrow d_Y \\ X_{n-1} & \xrightarrow{f} & Y_{n-1} \\ \vdots & & \vdots \end{array}$$

Cofiber sequence of complexes

$$\begin{array}{ccc}
 X & \xrightarrow{f} & Y \\
 \vdots & & \vdots \\
 \downarrow & & \downarrow \\
 X_{n+1} & \xrightarrow{f} & Y_{n+1} \\
 \downarrow d_X & & \downarrow d_Y \\
 X_n & \xrightarrow{f} & Y_n \\
 \downarrow d_X & & \downarrow d_Y \\
 X_{n-1} & \xrightarrow{f} & Y_{n-1} \\
 \vdots & & \vdots
 \end{array}
 \qquad
 \begin{array}{c}
 C_f \\
 \text{mapping cone} \\
 \vdots \\
 \downarrow \\
 Y_{n+1} \oplus X_n \\
 \downarrow \begin{pmatrix} d_Y & f \\ 0 & -d_X \end{pmatrix} = d_{C_f} \\
 Y_n \oplus X_{n-1} \\
 \downarrow \begin{pmatrix} d_Y & f \\ 0 & -d_X \end{pmatrix} = d_{C_f} \\
 Y_{n-1} \oplus X_{n-2} \\
 \vdots
 \end{array}$$

Cofiber sequence of complexes

$$\begin{array}{ccccc}
 X & \xrightarrow{f} & Y & \xrightarrow{i} & C_f \\
 & & & & \text{mapping cone} \\
 \vdots & & \vdots & & \vdots \\
 \downarrow & & \downarrow & & \downarrow \\
 X_{n+1} & \xrightarrow{f} & Y_{n+1} & \xrightarrow{\text{inclusion}} & Y_{n+1} \oplus X_n \\
 \downarrow d_X & & \downarrow d_Y & & \downarrow \begin{pmatrix} d_Y & f \\ 0 & -d_X \end{pmatrix} = d_{C_f} \\
 X_n & \xrightarrow{f} & Y_n & \xrightarrow{\text{inclusion}} & Y_n \oplus X_{n-1} \\
 \downarrow d_X & & \downarrow d_Y & & \downarrow \begin{pmatrix} d_Y & f \\ 0 & -d_X \end{pmatrix} = d_{C_f} \\
 X_{n-1} & \xrightarrow{f} & Y_{n-1} & \xrightarrow{\text{inclusion}} & Y_{n-1} \oplus X_{n-2} \\
 \vdots & & \vdots & & \vdots
 \end{array}$$

Cofiber sequence of complexes

$$\begin{array}{ccccccc}
 X & \xrightarrow{f} & Y & \xrightarrow{i} & C_f & & \Sigma X \text{ suspension} \\
 & & & & \text{mapping cone} & & \\
 \vdots & & \vdots & & \vdots & & \vdots \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 X_{n+1} & \xrightarrow{f} & Y_{n+1} & \xrightarrow{\text{inclusion}} & Y_{n+1} \oplus X_n & & X_n \\
 \downarrow d_X & & \downarrow d_Y & & \downarrow \begin{pmatrix} d_Y & f \\ 0 & -d_X \end{pmatrix} = d_{C_f} & & \downarrow -d_X \\
 X_n & \xrightarrow{f} & Y_n & \xrightarrow{\text{inclusion}} & Y_n \oplus X_{n-1} & & X_{n-1} \\
 \downarrow d_X & & \downarrow d_Y & & \downarrow \begin{pmatrix} d_Y & f \\ 0 & -d_X \end{pmatrix} = d_{C_f} & & \downarrow -d_X \\
 X_{n-1} & \xrightarrow{f} & Y_{n-1} & \xrightarrow{\text{inclusion}} & Y_{n-1} \oplus X_{n-2} & & X_{n-2} \\
 \vdots & & \vdots & & \vdots & & \vdots
 \end{array}$$

Cofiber sequence of complexes

$$\begin{array}{ccccccc}
 X & \xrightarrow{f} & Y & \xrightarrow{i} & C_f & \xrightarrow{q} & \Sigma X \text{ suspension} \\
 & & & & \text{mapping cone} & & \\
 \vdots & & \vdots & & \vdots & & \vdots \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 X_{n+1} & \xrightarrow{f} & Y_{n+1} & \xrightarrow{\text{inclusion}} & Y_{n+1} \oplus X_n & \xrightarrow{\text{projection}} & X_n \\
 \downarrow d_X & & \downarrow d_Y & & \downarrow \begin{pmatrix} d_Y & f \\ 0 & -d_X \end{pmatrix} = d_{C_f} & & \downarrow -d_X \\
 X_n & \xrightarrow{f} & Y_n & \xrightarrow{\text{inclusion}} & Y_n \oplus X_{n-1} & \xrightarrow{\text{projection}} & X_{n-1} \\
 \downarrow d_X & & \downarrow d_Y & & \downarrow \begin{pmatrix} d_Y & f \\ 0 & -d_X \end{pmatrix} = d_{C_f} & & \downarrow -d_X \\
 X_{n-1} & \xrightarrow{f} & Y_{n-1} & \xrightarrow{\text{inclusion}} & Y_{n-1} \oplus X_{n-2} & \xrightarrow{\text{projection}} & X_{n-2} \\
 \vdots & & \vdots & & \vdots & & \vdots
 \end{array}$$

Crash course on singular cohomology

Compute the cohomology of spheres,

$$S^n = \{(x_0, \dots, x_n) \in \mathbb{R}^{n+1} \mid x_0^2 + \dots + x_n^2 = 1\};$$

$$S^0 = \{\pm 1\} = \bullet \quad \bullet, \quad H^0(S^0, \mathbb{Z}) = \text{pointed maps } S^0 \rightarrow \mathbb{Z} \cong \mathbb{Z};$$

$$\Sigma S^0 = \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \end{array} \cong \bigcirc = S^1,$$

$$H^1(S^1, \mathbb{Z}) \cong H^0(S^0, \mathbb{Z}) \cong \mathbb{Z}, \quad H^n(S^1, \mathbb{Z}) = 0 \text{ for } n \neq 1;$$

Crash course on singular cohomology

Compute the cohomology of spheres,

$$S^n = \{(x_0, \dots, x_n) \in \mathbb{R}^{n+1} \mid x_0^2 + \dots + x_n^2 = 1\};$$

$$S^0 = \{\pm 1\} = \bullet \quad \bullet, \quad H^0(S^0, \mathbb{Z}) = \text{pointed maps } S^0 \rightarrow \mathbb{Z} \cong \mathbb{Z};$$

$$\Sigma S^0 = \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \end{array} \cong \bigcirc = S^1,$$

$$H^1(S^1, \mathbb{Z}) \cong H^0(S^0, \mathbb{Z}) \cong \mathbb{Z}, \quad H^n(S^1, \mathbb{Z}) = 0 \text{ for } n \neq 1;$$

Crash course on singular cohomology

Compute the cohomology of spheres,

$$S^n = \{(x_0, \dots, x_n) \in \mathbb{R}^{n+1} \mid x_0^2 + \dots + x_n^2 = 1\};$$

$$S^0 = \{\pm 1\} = \bullet \quad \bullet, \quad H^0(S^0, \mathbb{Z}) = \text{pointed maps } S^0 \rightarrow \mathbb{Z} \cong \mathbb{Z};$$

$$\Sigma S^0 = \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \end{array} \cong \bigcirc = S^1,$$

$$H^1(S^1, \mathbb{Z}) \cong H^0(S^0, \mathbb{Z}) \cong \mathbb{Z}, \quad H^n(S^1, \mathbb{Z}) = 0 \text{ for } n \neq 1;$$

Crash course on singular cohomology

Compute the cohomology of spheres,

$$S^n = \{(x_0, \dots, x_n) \in \mathbb{R}^{n+1} \mid x_0^2 + \dots + x_n^2 = 1\};$$

$$S^0 = \{\pm 1\} = \bullet \quad \bullet, \quad H^0(S^0, \mathbb{Z}) = \text{pointed maps } S^0 \rightarrow \mathbb{Z} \cong \mathbb{Z};$$

$$\Sigma S^0 = \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \end{array} \cong \bigcirc = S^1,$$

$$H^1(S^1, \mathbb{Z}) \cong H^0(S^0, \mathbb{Z}) \cong \mathbb{Z}, \quad H^n(S^1, \mathbb{Z}) = 0 \text{ for } n \neq 1;$$

Crash course on singular cohomology

Compute the cohomology of spheres,

$$S^n = \{(x_0, \dots, x_n) \in \mathbb{R}^{n+1} \mid x_0^2 + \dots + x_n^2 = 1\};$$

$$S^0 = \{\pm 1\} = \bullet \quad \bullet, \quad H^0(S^0, \mathbb{Z}) = \text{pointed maps } S^0 \rightarrow \mathbb{Z} \cong \mathbb{Z};$$

$$\Sigma S^0 = \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \end{array} \cong \bigcirc = S^1,$$

$$H^1(S^1, \mathbb{Z}) \cong H^0(S^0, \mathbb{Z}) \cong \mathbb{Z}, \quad H^n(S^1, \mathbb{Z}) = 0 \text{ for } n \neq 1;$$

Crash course on singular cohomology

Compute the cohomology of spheres,

$$S^n = \{(x_0, \dots, x_n) \in \mathbb{R}^{n+1} \mid x_0^2 + \dots + x_n^2 = 1\};$$

$$S^0 = \{\pm 1\} = \bullet \quad \bullet, \quad H^0(S^0, \mathbb{Z}) = \text{pointed maps } S^0 \rightarrow \mathbb{Z} \cong \mathbb{Z};$$

$$\Sigma S^0 = \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \end{array} \cong \bigcirc = S^1,$$

$$H^1(S^1, \mathbb{Z}) \cong H^0(S^0, \mathbb{Z}) \cong \mathbb{Z}, \quad H^n(S^1, \mathbb{Z}) = 0 \text{ for } n \neq 1;$$

Crash course on singular cohomology

Compute the cohomology of spheres,

$$S^n = \{(x_0, \dots, x_n) \in \mathbb{R}^{n+1} \mid x_0^2 + \dots + x_n^2 = 1\};$$

$$S^0 = \{\pm 1\} = \bullet \quad \bullet, \quad H^0(S^0, \mathbb{Z}) = \text{pointed maps } S^0 \rightarrow \mathbb{Z} \cong \mathbb{Z};$$

$$\Sigma S^0 = \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \bullet \quad \bullet \\ \diagdown \quad \diagup \\ \bullet \end{array} \cong \bigcirc = S^1,$$

$$H^1(S^1, \mathbb{Z}) \cong H^0(S^0, \mathbb{Z}) \cong \mathbb{Z}, \quad H^n(S^1, \mathbb{Z}) = 0 \text{ for } n \neq 1;$$

Crash course on singular cohomology

Compute the cohomology of spheres,

$$S^n = \{(x_0, \dots, x_n) \in \mathbb{R}^{n+1} \mid x_0^2 + \dots + x_n^2 = 1\};$$

$$\Sigma S^1 = \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \text{---} \text{---} \text{---} \\ \diagdown \quad \diagup \\ \bullet \end{array} \cong \begin{array}{c} \text{---} \text{---} \text{---} \\ \text{---} \text{---} \text{---} \end{array} = S^2,$$

$$H^2(S^2, \mathbb{Z}) \cong H^1(S^1, \mathbb{Z}) \cong \mathbb{Z}, \quad H^n(S^2, \mathbb{Z}) = 0 \text{ for } n \neq 2;$$

and inductively,

$$\Sigma S^{n-1} = S^n, \quad \begin{aligned} H^n(S^n, \mathbb{Z}) &\cong H^{n-1}(S^{n-1}, \mathbb{Z}) \cong \mathbb{Z}, \\ H^m(S^n, \mathbb{Z}) &= 0 \text{ for } m \neq n. \end{aligned}$$

Crash course on singular cohomology

Compute the cohomology of spheres,

$$S^n = \{(x_0, \dots, x_n) \in \mathbb{R}^{n+1} \mid x_0^2 + \dots + x_n^2 = 1\};$$

$$\Sigma S^1 = \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \text{---} \text{---} \text{---} \text{---} \\ \diagdown \quad \diagup \\ \bullet \end{array} \cong \begin{array}{c} \text{---} \text{---} \text{---} \text{---} \\ \diagup \quad \diagdown \\ \text{---} \text{---} \text{---} \text{---} \end{array} = S^2,$$

$$H^2(S^2, \mathbb{Z}) \cong H^1(S^1, \mathbb{Z}) \cong \mathbb{Z}, \quad H^n(S^2, \mathbb{Z}) = 0 \text{ for } n \neq 2;$$

and inductively,

$$\Sigma S^{n-1} = S^n, \quad \begin{aligned} H^n(S^n, \mathbb{Z}) &\cong H^{n-1}(S^{n-1}, \mathbb{Z}) \cong \mathbb{Z}, \\ H^m(S^n, \mathbb{Z}) &= 0 \text{ for } m \neq n. \end{aligned}$$

Crash course on singular cohomology

Compute the cohomology of spheres,

$$S^n = \{(x_0, \dots, x_n) \in \mathbb{R}^{n+1} \mid x_0^2 + \dots + x_n^2 = 1\};$$

$$\Sigma S^1 = \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \text{---} \text{---} \text{---} \text{---} \\ \diagdown \quad \diagup \\ \bullet \end{array} \cong \begin{array}{c} \text{---} \text{---} \text{---} \text{---} \\ \diagup \quad \diagdown \\ \text{---} \text{---} \text{---} \text{---} \\ \diagdown \quad \diagup \end{array} = S^2,$$

$$H^2(S^2, \mathbb{Z}) \cong H^1(S^1, \mathbb{Z}) \cong \mathbb{Z}, \quad H^n(S^2, \mathbb{Z}) = 0 \text{ for } n \neq 2;$$

and inductively,

$$\Sigma S^{n-1} = S^n, \quad \begin{aligned} H^n(S^n, \mathbb{Z}) &\cong H^{n-1}(S^{n-1}, \mathbb{Z}) \cong \mathbb{Z}, \\ H^m(S^n, \mathbb{Z}) &= 0 \text{ for } m \neq n. \end{aligned}$$

Crash course on singular cohomology

Compute the cohomology of spheres,

$$S^n = \{(x_0, \dots, x_n) \in \mathbb{R}^{n+1} \mid x_0^2 + \dots + x_n^2 = 1\};$$

$$\Sigma S^1 = \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \text{---} \text{---} \text{---} \\ \diagdown \quad \diagup \\ \bullet \end{array} \cong \begin{array}{c} \text{---} \text{---} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \text{---} \text{---} \end{array} = S^2,$$

$$H^2(S^2, \mathbb{Z}) \cong H^1(S^1, \mathbb{Z}) \cong \mathbb{Z}, \quad H^n(S^2, \mathbb{Z}) = 0 \text{ for } n \neq 2;$$

and inductively,

$$\Sigma S^{n-1} = S^n, \quad \begin{aligned} H^n(S^n, \mathbb{Z}) &\cong H^{n-1}(S^{n-1}, \mathbb{Z}) \cong \mathbb{Z}, \\ H^m(S^n, \mathbb{Z}) &= 0 \text{ for } m \neq n. \end{aligned}$$

Crash course on singular cohomology

Compute the cohomology of spheres,

$$S^n = \{(x_0, \dots, x_n) \in \mathbb{R}^{n+1} \mid x_0^2 + \dots + x_n^2 = 1\};$$

$$\Sigma S^1 = \begin{array}{c} \bullet \\ \diagup \quad \diagdown \\ \text{---} \text{---} \text{---} \text{---} \\ \diagdown \quad \diagup \\ \bullet \end{array} \cong \begin{array}{c} \text{---} \text{---} \text{---} \text{---} \\ \diagup \quad \diagdown \\ \text{---} \text{---} \text{---} \text{---} \end{array} = S^2,$$


$$H^2(S^2, \mathbb{Z}) \cong H^1(S^1, \mathbb{Z}) \cong \mathbb{Z}, \quad H^n(S^2, \mathbb{Z}) = 0 \text{ for } n \neq 2;$$

and inductively,

$$\Sigma S^{n-1} = S^n, \quad \begin{aligned} H^n(S^n, \mathbb{Z}) &\cong H^{n-1}(S^{n-1}, \mathbb{Z}) \cong \mathbb{Z}, \\ H^m(S^n, \mathbb{Z}) &= 0 \text{ for } m \neq n. \end{aligned}$$


Crash course on singular cohomology

Let C_f be a simplicial complex,




Crash course on singular cohomology

Let C_f be a simplicial complex,




Any simplex is a cone over its boundary, therefore C_f can be obtained from Y ,




Crash course on singular cohomology

Let C_f be a simplicial complex,




as the mapping cone of the **inclusion** f ,

Any simplex is a cone over its boundary, therefore C_f can be obtained from Y ,




Crash course on singular cohomology

Let C_f be a simplicial complex,



Any simplex is a cone over its boundary, therefore C_f can be obtained from Y ,

inclusion
 i



as the mapping cone of the inclusion f ,

$$S^1 = X \xrightarrow[\text{inclusion}]{f} Y \xrightarrow[\text{inclusion}]{i} C_f \xrightarrow{q} \Sigma X = S^2.$$

$X =$



f inclusion

Crash course on singular cohomology

Given a cofiber sequence

$$S^n \xrightarrow{f} Y \xrightarrow{i} C_f \xrightarrow{q} S^{n+1};$$

$$H^m(Y, \mathbb{Z}) = H^m(C_f, \mathbb{Z}), \text{ if } m \neq n, n+1;$$

$$0 \leftarrow H^{n+1}(C_f, \mathbb{Z}) \xleftarrow{\mathbb{Z} \xleftarrow{H^n(f, \mathbb{Z})}} H^n(Y, \mathbb{Z}) \xleftarrow{} H^n(C_f, \mathbb{Z}) \leftarrow 0,$$

$$H^n(C_f, \mathbb{Z}) = \text{Ker } H^n(f, \mathbb{Z}),$$

$$H^{n+1}(C_f, \mathbb{Z}) = \text{Coker } H^n(f, \mathbb{Z}).$$

◀ back

Crash course on singular cohomology

Given a cofiber sequence

$$S^n \xrightarrow{f} Y \xrightarrow{i} C_f \xrightarrow{q} S^{n+1};$$

$$H^m(Y, \mathbb{Z}) = H^m(C_f, \mathbb{Z}), \text{ if } m \neq n, n+1;$$

$$0 \leftarrow H^{n+1}(C_f, \mathbb{Z}) \xleftarrow{\mathbb{Z} \xleftarrow{H^n(f, \mathbb{Z})}} H^n(Y, \mathbb{Z}) \xleftarrow{} H^n(C_f, \mathbb{Z}) \leftarrow 0,$$

$$H^n(C_f, \mathbb{Z}) = \text{Ker } H^n(f, \mathbb{Z}),$$

$$H^{n+1}(C_f, \mathbb{Z}) = \text{Coker } H^n(f, \mathbb{Z}).$$

◀ back

Crash course on singular cohomology

Given a cofiber sequence

$$S^n \xrightarrow{f} Y \xrightarrow{i} C_f \xrightarrow{q} S^{n+1};$$

$$H^m(Y, \mathbb{Z}) = H^m(C_f, \mathbb{Z}), \text{ if } m \neq n, n+1;$$

$$0 \leftarrow H^{n+1}(C_f, \mathbb{Z}) \xleftarrow{\mathbb{Z} \xleftarrow{H^n(f, \mathbb{Z})}} H^n(Y, \mathbb{Z}) \xleftarrow{H^n(C_f, \mathbb{Z})} 0,$$

$$H^n(C_f, \mathbb{Z}) = \text{Ker } H^n(f, \mathbb{Z}),$$

$$H^{n+1}(C_f, \mathbb{Z}) = \text{Coker } H^n(f, \mathbb{Z}).$$

◀ back

Crash course on singular cohomology

Given a cofiber sequence

$$S^n \xrightarrow{f} Y \xrightarrow{i} C_f \xrightarrow{q} S^{n+1};$$

$$H^m(Y, \mathbb{Z}) = H^m(C_f, \mathbb{Z}), \text{ if } m \neq n, n+1;$$

$$0 \leftarrow H^{n+1}(C_f, \mathbb{Z}) \xleftarrow{\mathbb{Z} \xleftarrow{H^n(f, \mathbb{Z})}} H^n(Y, \mathbb{Z}) \xleftarrow{} H^n(C_f, \mathbb{Z}) \leftarrow 0,$$

$$H^n(C_f, \mathbb{Z}) = \text{Ker } H^n(f, \mathbb{Z}),$$

$$H^{n+1}(C_f, \mathbb{Z}) = \text{Coker } H^n(f, \mathbb{Z}).$$

◀ back