Supplementary Material

Thermal time dominates satellite predictors for within-season wheat biomass estimation under spatial cross-validation

Francisco Zambrano[a], Ignacio Fuentes[a], Abel Herrera[b], Mauricio Molina-Rocco[c]

[a] *Facultad de Medicina Veterinaria y Agronomía, Universidad de Las Américas, Santiago, Chile,*

[b] *Hemera Centro de Observación de la Tierra, Facultad de Ciencias, Ingeniería y Tecnología; Universidad Mayor, Santiago, Chile,*

[c] *Laboratorio de Suelos y Sostenibilidad. Departamento de Acuicultura y Recursos Agroalimentarios, Universidad de Los Lagos, Osorno, 5311157, Chile,*

---

1. Table S1: Additional vegetation indices

Table 3 of the main text lists the 21 vegetation indices (VIs), derived from Sentinel-2 (S2) and PlanetScope (PS) spectral bands, that were retained as predictors after eliminating highly correlated indices (Section 2.2.5 of the main text). Table S1 below lists the remaining 32 VIs that were part of the same 53-index base candidate set (Section 2.2.5) but were not retained in Table 3.

As described in the main text, a subset of these S2-derived indices were additionally recalculated using the Sentinel-2 Band 8A (narrow near-infrared) instead of Band 8 (broad near-infrared), following the same formula with the near-infrared (NIR) term replaced by the Band 8A reflectance; this is denoted with an asterisk (*) next to the S2 dataset code, following the same convention as Table 3 of the main text. Together, the 53 base indices (21 in Table 3 and 32 in Table S1) and their 36 Band-8A-recalculated counterparts make up the full set of 89 candidate VIs referenced in Section 2.2.5 of the main text.

Band and index notation follows Table 3 of the main text: Red, Green, Blue = visible bands; RE1, RE2, RE3 = S2 red-edge bands 5, 6, and 7; NIR = near-infrared band (S2 Band 8 / PS Band 8); SWIR1, SWIR2 = S2 shortwave-infrared bands 11 and 12; Water vapour = S2 Band 9. Formula definitions follow standard vegetation-index literature, including the references cited in Section 2.2.5 of the main text for VI selection (Peng et al., 2023; Segarra et al., 2022; Uribeetxebarria et al., 2023; Wang et al., 2022; Wu et al., 2008; Zheng et al., 2017); the exact implementation of every index listed here and in Table 3 is available in the project code repository (script `02_calcular_vis.R`).

*Corresponding author

Table S1. Additional vegetation indices (covariates) derived from Sentinel-2 (S2) and PlanetScope (PS) spectral bands, computed as part of the 53-index base candidate set (Section 2.2.5 of the main text) but not retained in Table 3. Asterisk (*) next to the S2 dataset code indicates that the index was also recalculated using Sentinel-2 Band 8A (narrow NIR) instead of Band 8 (broad NIR).

N°	Dataset used	Name	Abbreviation	Formula
1	S2	NDVI Variant (Red-edge 3)	$NDVI_2$	$(RE_3 - Red)/(RE_3+Red)$
2	S2	Red-edge NDVI (705 nm)	$NDVI_{705}$	$(RE_2 - RE_1)/(RE_2 + RE_1)$
3	S2*-PS	Modified Simple Ratio	MSR	$(NIR/Red - 1)/\sqrt{(NIR/Red + 1)}$
4	S2	Red-edge Modified Simple Ratio	MSR_{705}	$(RE_2/RE_1 - 1)/\sqrt{(RE_2/RE_1 + 1)}$
5	S2*-PS	Soil-Adjusted Vegetation Index	SAVI	$\frac{1.5\cdot(NIR-Red)}{(NIR+Red+0.5)}$
6	S2	SAVI Variant (Red-edge 3)	$SAVI_2$	$\frac{1.5\cdot(RE_3-Red)}{(RE_3+Red+0.5)}$
7	S2*-PS	Green SAVI	$SAVI_{green}$	$\frac{1.5\cdot(NIR-Green)}{(NIR+Green+0.5)}$
8	S2*-PS	Green OSAVI	$OSAVI_{green}$	$\frac{1.16\cdot(NIR-Green)}{(NIR+Green+0.16)}$
9	S2*-PS	Red-edge OSAVI	$OSAVI_{rededge}$	$\frac{1.16\cdot(NIR-RE_1)}{(NIR+RE_1+0.16)}$
10	S2	-	$TCARI/OSAVI_2$	$TCARI/OSAVI_2$

N°	Dataset used	Name	Abbreviation	Formula
11	S2*	SWIR 11 Related OSAVI	$\text{OSAVI}_{\text{SWIR11}}$	$\frac{1.16\cdot(NIR-SWIR_1)}{(NIR+SWIR_1+0.16)}$
12	S2	SWIR 12 Related TCARI	$\text{TCARI}_{\text{SWIR12}}$	$3\cdot((RE_1-SWIR_2)-0.2\cdot(RE_1-Green)\cdot(RE_1/SWIR_2))$
13	S2*	SWIR 12 Related OSAVI	$\text{OSAVI}_{\text{SWIR12}}$	$\frac{1.16\cdot(NIR-SWIR_2)}{(NIR+SWIR_2+0.16)}$
14	S2*-PS	Chlorophyll Index - Green	CI_{green}	$(NIR/Green)-1$
15	S2*-PS	Chlorophyll Index - Red-edge	$\text{CI}_{\text{rededge}}$	$(NIR/RE_1)-1$
16	S2*-PS	OSAVI x Chlorophyll Index Red-edge	OSAVI x $\text{CI}_{\text{rededge}}$	$\left[\frac{1.16\cdot(NIR-Red)}{(NIR+Red+0.16)}\right]\cdot[(NIR/RE_1)-1]$
17	S2*-PS	Difference Vegetation Index	DVI	$NIR-Red$
18	S2*-PS	Green DVI	$\text{DVI}_{\text{green}}$	$NIR-Green$
19	S2*-PS	Red-edge DVI	$\text{DVI}_{\text{rededge}}$	$NIR-RE_1$
20	S2	Two-Band Enhanced Vegetation Index	EVI2	$\frac{2.5\cdot(RE_3-Red)}{(1+RE_3+2.4\cdot Red)}$
21	S2*-PS	Red-edge Vegetation Index 1	REVI1	$NIR-RE_1$

N°	Dataset used	Name	Abbreviation	Formula
22	S2*	Red-edge Vegetation Index 2	REVI2	$NIR - RE_2$
23	S2*-PS	Weighted Difference Vegetation Index	WDVI	$NIR - 0.5 \cdot Red$
24	S2-PS	Green-Red Vegetation Index	GRVI	$(Green - Red)/(Green + Red)$
25	S2*-PS	Green Ratio Vegetation Index	GRVI2	$NIR/Green$
26	S2*-PS	Green Atmospherically Resistant Index	GARI	$\frac{NIR-(Green-1.7\cdot(Blue-Red))}{NIR+(Green-1.7\cdot(Blue-Red))}$
27	S2*-PS	Leaf Chlorophyll Index	LCI	$(NIR - RE_1)/(NIR - Red)$
28	S2	MERIS Terrestrial Chlorophyll Index	MTCI	$(RE_2 - RE_1)/(RE_1 - Red)$
29	S2*	Normalized Difference Red-edge Index 2	NDRE2	$(NIR - RE_2)/(NIR + RE_2)$

N°	Dataset used	Name	Abbreviation	Formula
30	S2	Normalized Difference Red-edge Index (RE3/Green)	NDREI	$(RE_3 - Green)/(RE_3 + Green)$
31	S2*	Normalized Difference Moisture Index	NDMI	$(NIR - SWIR_1)/(NIR + SWIR_1)$
32	S2*-PS	Kernel NDVI	KNDVI	$\frac{1 - \exp\left(-\frac{(NIR - Red)^2}{2 \cdot 0.15^2}\right)}{1 + \exp\left(-\frac{(NIR - Red)^2}{2 \cdot 0.15^2}\right)}$

2. Note S1: Predictors removed by the correlation filter in each fold

The correlation filter (Pearson r > 0.9) was estimated on the training data of each LOSSO-CV fold, after imputation and normalization, so the retained predictors differ among folds. Table S3 lists the predictors removed from the set with all 54 candidate predictors (All) in each fold. Most removed predictors are daily Sentinel-2 indices that are highly correlated with one another. $\sum$GDD was removed in the folds where Villa Baviera and La Cancha were held out, because in those training sets it was highly correlated with cumulative vegetation indices. The removals can be reproduced with `script/31_loso_diseno_limpio.R`.

Table S3. Predictors removed by the within-fold correlation filter from the All set, by held-out field. The suffix 8A marks Sentinel-2 indices computed with band B8A, and cumsum marks cumulative sums from the start of the season.

3. Equations S1-S3: Performance metrics

Section 2.3.4 of the main text reports model performance using the mean absolute error (MAE), root mean square error (RMSE), and coefficient of determination (R^2), defined in Equations S1-S3 below and computed separately for each held-out site-season, where y_i and $\hat{y}_i$ are the observed and out-of-fold predicted AGB values for sample i, $\bar{y}$ is the mean observed AGB of the held-out site-season, and n is the number of samples in that site-season. R^2 computed in this way is negative when the predictions are less accurate than the mean of the held-out site-season.

Held-out field	n	Removed predictors
Hidango North	17	S2: MCARI/OSAVI2, TCARI/OSAVI2, NDRE2, GNDVI, MCARI/OSAVI 8A, SIPI 8A, $MCARI_{SW11}$, OSAVI, OSAVI2, NDRE1, SR, TCARI/OSAVI, CI_{red}, EVI, CI_{red} 8A cumsum; PS: B8, SR
Hidango South	15	S2: TCARI/OSAVI2, $TCARI_{SW11}$, OSAVI2, NDRE1, EVI, GNDVI, MCARI/OSAVI 8A, NDRE3 8A cumsum, MCARI/OSAVI, OSAVI, SR, TCARI/OSAVI, CI_{red}, SIPI; PS: CI_{red}
La Cancha	16	$\sum$GDD; S2: TCARI/OSAVI2, OSAVI, OSAVI2, NDRE1, EVI, GNDVI, CI_{red}, MCARI/OSAVI 8A, MCARI/OSAVI, $MCARI_{SW11}$, TCARI/OSAVI, SR, SIPI; PS: OSAVI, SR
Villa Baviera	20	$\sum$GDD; S2: TCARI/OSAVI2, GNDVI, CI_{red}, MCARI/OSAVI 8A, MCARI/OSAVI, OSAVI, OSAVI2, NDRE2, NDRE1, TCARI/OSAVI, SR, SIPI, EVI; PS: OSAVI, EVI, SIPI, CI_{red}, EVI cumsum, CI_{red} cumsum

$$MAE = \frac{1}{n}\sum_{i=1}^{n}|y_i - \hat{y}_i| \tag{S1}$$

$$RMSE = \sqrt{\frac{1}{n}\sum_{i=1}^{n}(y_i - \hat{y}_i)^2} \tag{S2}$$

$$R^2 = 1 - \frac{\sum_{i=1}^{n}(y_i - \hat{y}_i)^2}{\sum_{i=1}^{n}(y_i - \bar{y})^2} \tag{S3}$$

4. Table S2: Performance per held-out site-season

Table S2 reports the RMSE and R^2 of every predictor set and model for each of the four held-out site-seasons. Mean and standard deviation across site-seasons are given in Table 5 of the main text. The values are produced by `script/31_loso_diseno_limpio.R` and stored in `output/tables/rev2_loso_limpio_por_fold.csv`.

Table S2. RMSE (t/ha) and, in parentheses, R^2 (1 − SSE/SST) for each held-out site-season under leave-one-site-season-out cross-validation (LOSSO-CV). Each value is obtained with a model trained on the other three site-seasons.

References

Peng, J., Zeiner, N., Parsons, D., Féret, J.B., Söderström, M., Morel, J., 2023. Forage biomass estimation using sentinel-2 imagery at high latitudes. Remote Sens. 15, 2350. doi:10.3390/rs15092350.

Predictor set	Model	Hidango North	Hidango South	La Cancha	Villa Baviera
DAS	LM	5.57 (0.74)	8.81 (−0.66)	9.95 (0.44)	7.04 (0.25)
GDD	LM	6.17 (0.68)	6.69 (0.04)	8.77 (0.57)	3.31 (0.83)
W	LM	7.24 (0.55)	7.69 (−0.27)	9.15 (0.53)	3.97 (0.76)
W	RF	7.69 (0.50)	7.64 (−0.25)	7.83 (0.66)	6.84 (0.29)
W	XGBoost	7.00 (0.58)	8.34 (−0.49)	8.31 (0.61)	6.57 (0.35)
S1	RF	12.48 (−0.33)	10.56 (−1.39)	12.67 (0.10)	7.99 (0.04)
S1	XGBoost	12.74 (−0.38)	11.24 (−1.71)	14.49 (−0.18)	10.29 (−0.60)
S2	RF	9.01 (0.31)	5.77 (0.29)	12.86 (0.07)	4.68 (0.67)
S2	XGBoost	9.46 (0.24)	5.61 (0.33)	12.30 (0.15)	4.33 (0.72)
PS	RF	6.61 (0.63)	4.91 (0.48)	8.45 (0.60)	14.62 (−2.23)
PS	XGBoost	7.68 (0.50)	5.36 (0.39)	9.26 (0.52)	15.57 (−2.66)
S1+GDD	RF	5.35 (0.76)	6.57 (0.08)	7.95 (0.65)	3.34 (0.83)
S1+GDD	XGBoost	5.69 (0.72)	7.79 (−0.30)	8.51 (0.59)	3.69 (0.79)
S2+GDD	RF	7.70 (0.49)	6.70 (0.04)	12.86 (0.07)	3.78 (0.78)
S2+GDD	XGBoost	8.45 (0.39)	6.02 (0.22)	12.30 (0.15)	3.91 (0.77)
PS+GDD	RF	5.62 (0.73)	5.42 (0.37)	7.94 (0.65)	9.41 (−0.34)
PS+GDD	XGBoost	6.85 (0.60)	5.58 (0.33)	8.62 (0.58)	7.35 (0.18)
S1+W	RF	6.69 (0.62)	6.85 (−0.01)	6.17 (0.79)	4.49 (0.70)
S1+W	XGBoost	5.78 (0.71)	7.65 (−0.26)	7.64 (0.67)	4.52 (0.69)
S2+W	RF	8.10 (0.44)	6.95 (−0.03)	12.65 (0.10)	2.98 (0.87)
S2+W	XGBoost	9.12 (0.29)	6.65 (0.05)	12.03 (0.19)	2.26 (0.92)
PS+W	RF	6.66 (0.62)	5.59 (0.33)	7.35 (0.70)	14.66 (−2.25)
PS+W	XGBoost	7.44 (0.53)	4.89 (0.49)	7.85 (0.65)	12.89 (−1.51)
S2+W+S1	RF	8.16 (0.43)	6.50 (0.09)	12.36 (0.14)	2.65 (0.89)
S2+W+S1	XGBoost	8.09 (0.44)	6.18 (0.18)	12.60 (0.11)	2.57 (0.90)
All	RF	7.00 (0.58)	5.43 (0.37)	10.67 (0.36)	5.05 (0.61)
All	XGBoost	8.04 (0.45)	5.47 (0.36)	10.14 (0.42)	3.58 (0.81)

Segarra, J., Araus, J., Kefauver, S., 2022. Farming and earth observation: Sentinel-2 data to estimate within-field wheat grain yield. Int. J. Appl. Earth Obs. Geoinformation 107, 102697. doi:10.1016/j.jag.2022.102697.

Uribeetxebarria, A., Castellón, A., Aizpurua, A., 2023. Optimizing wheat yield prediction integrating data from sentinel-1 and sentinel-2 with catboost algorithm. Remote Sens. 15, 1640. doi:10.3390/rs15061640.

Wang, F., Yang, M., Ma, L., Zhang, T., Qin, W., Li, W., Zhang, Y., Sun, Z., Wang, Z., Li, F., Yu, K., 2022. Estimation of above-ground biomass of winter wheat based on consumer-grade multi-spectral uav. Remote Sens. 14, 1251. doi:10.3390/rs14051251.

Wu, C., Niu, Z., Tang, Q., Huang, W., 2008. Estimating chlorophyll content from hyperspectral vegetation indices: Modeling and validation. Agric. For. Meteorol. 148, 1230–1241. doi:10.1016/j.agrformet.2008.03.005.

Zheng, Y., Wu, B., Zhang, M., 2017. Estimating the above ground biomass of winter wheat using the sentinel-2 data. Natl. Remote Sens. Bull. 21, 318–328. doi:10.11834/jrs.20176269.