


Poly-substance use, treatment completion, and contact with the justice system: a multistate analysis of treatments for substance use disorders between 2010-2019 in Chile



A. González-Santa Cruz^{1, 2, }, J. Ruiz-Tagle Maturana^{1, 3, }, M. Mateo Piñones^{1, 4, }, A. Castillo-Carniglia^{5, 5, }



¹ Young Researcher, Millennium Nucleus for the evaluation and analysis of Drug Policies

² Ph.D. student, School of Public Health, Universidad de Chile

³ Ph.D. student, Programa de Doctorado en Políticas Públicas, Universidad Mayor, Santiago, Chile.

⁴ Ph.D. student, Griffith University, Australia

Background

Substance use disorders (SUD) often co-occur with criminality, including violence, arrests, and incarceration^[1;2;3;4;5]. People with polysubstance use (PSU) are considered a high-risk population, as they are associated with mortality, relapse, and contact with the criminal justice system (CJS)^[6;7;8;9;10]. Although completing SUD treatment is linked with better outcomes, including preventing contact with CJS, the role of treatment completion in the link between PSU and contact with CJS is unclear^[11;12;13;14]. Studies have found mixed evidence regarding the association between PSU and treatment completion rates^[15;16;17;18;19]. Thus, it is crucial to determine the role of treatment completion in order to improve outcomes in people with PSU. However, analyzing the role of treatment outcomes in people with PSU is challenging, as there is limited research on this population in Latin America, and high-risk populations have often been overlooked^[20;21;22;23;24].

Objectives

Estimate the mediating effects of completing SUD treatment on the relationship between PSU at admission and contact with CJS among adult patients admitted to SUD treatment programs in Chile during 2010-2019. Specific: (1) To describe the prevalence of PSU, treatment completion, and contact with CJS in the sample, (2) to compare the risk of contact with CJS between people with poly and single-substance use, and (3) to estimate the proportion of the effect of PSU and treatment outcome on the contact with CJS.

Methods

This research design is a retrospective cohort based on the administrative data's record linkage. The study will use data from Chilean SUTs programs and Prosecutor's Office through a deterministic linkage process. We will request an amendment to an existing ethical approval from a study using the same data.

Variables

Exposure: baseline PSU (using more than one main substance among alcohol and illicit drugs at admission to SUD treatment, whether sequential or concurrent); **Mediator:** SUD treatment outcome (complete vs. dropout or spelled by misconduct); **Outcome:** contact with CJS (offense that led to a condemnatory sentence).

Analytical Plan

The study controlled for various confounding variables related to substance use, demographics, and social factors through weights generated through the inverse probability of PSU. Patients were weighted by the inverse probability of

polysubstance use based on several predictors; Weights were truncated at the 1st and 99th percentiles^[25].




Figure 1: Covariate balance

We used the illness-death multistate model to simultaneously estimate transitions between admission and treatment outcome, treatment outcome and contact with CJS, and admission and contact with CJS (without completing treatment).




Figure 2: Multistate scheme

We calculated the Aalen-Johansen estimator for transition probabilities at 6 months, 1 and 3 years using the `multistate` Stata command^[26]. Secondary analyses focused on mediation, estimating the effects of PSU given treatment outcome at 6 months, 1 and 3 years using a standard time-to-first-event approach. Proportions mediated were estimated using the bootstrap method or m-estimation of standard errors on standardized survival curves and restricted mean survival times (RMST) through the `stpm2` and `stipw` Stata commands^[27;28;29;30]. We also plan to run separate analyses on patients admitted to different treatment settings. Preliminary markdowns are available on: https://fondecytacc.github.io/ndP/an_ser_2023_step_0.html.

Preliminary Results

-

References

- [1] A. A. Duke, K. M. Z. Smith, L. M. S. Oberleitner, et al. "Alcohol, drugs, and violence: A meta-meta-analysis." In: *Psychology of Violence* 8.2 (mar.. 2018), pp. 238–249. ISSN: 2152-081X. DOI: 10.1037/vio0000106.
- [2] N. F. Sgale and K. Turney. "Beyond Incarceration: Criminal Justice Contact and Mental Health". In: *American Sociological Review* 82.4 (ago.. 2017), pp. 719–743. ISSN: 0003-1224. DOI: 10.1177/0003122417131386.
- [3] E. G. Thomas, M. J. Spittal, F. S. Taxman, et al. "Association between contact with mental health and substance use services and reincarceration after release from prison". In: *PLOS ONE* 17.9 (sept.. 2022), p. e0272870. ISSN: 1932-6203. DOI: 10.1371/journal.pone.0272870.
- [4] Y. Liu, V. Williamson, B. Sellow, et al. "The importance of considering polysubstance use: lessons from cocaine research". In: *Drug and Alcohol Dependence* 192 (nov.. 2018), pp. 16–28. ISSN: 03768716. DOI: 10.1016/j.drugalcdep.2018.07.022.
- [5] J. P. Connor, M. J. Gullio, A. White, et al. "Polysubstance use". In: *Current Opinion in Psychiatry* 27.4 (jul.. 2014), pp. 269–275. ISSN: 0951-7367. DOI: 10.1097/YCO.0000000000000069.
- [6] L. Giering and A. L. Brettevill-Jensen. "Patterns of substance use and mortality risk in a cohort of 'hard-to-reach' polysubstance users". In: *Addiction* 113.4 (abr.. 2018), pp. 729–739. ISSN: 09592140. DOI: 10.1111/add.14053.
- [7] A. N. Hassan and B. Le Foll. "Polydrug use disorders in individuals with opioid use disorder". In: *Drug and Alcohol Dependence* 198 (may.. 2019), pp. 28–33. ISSN: 03768716. DOI: 10.1016/j.drugalcdep.2019.01.031.
- [8] L. Wang, J. E. Min, E. Krebs, et al. "Polydrug use and its association with drug treatment outcomes among primary heroin, methamphetamine, and cocaine users". In: *International Journal of Drug Policy* 49 (nov.. 2017), pp. 32–40. ISSN: 09593959. DOI: 10.1016/j.drugpo.2017.07.009.
- [9] L. Quenk, G. C. K. Chan, A. White, et al. "Concurrent and Simultaneous Polydrug Use: Latent Class Analysis of an Australian Nationally Representative Sample of Young Adults". In: *Frontiers in Public Health* 1 (2013). ISSN: 2296-2665. DOI: 10.3389/fpubh.2013.00061.
- [10] J. A. Ford, K. Ortiz, T. S. Schepps, et al. "Types of criminal legal system exposure and polysubstance use: Prevalence and correlates among U.S. adults in the National Survey on Drug Use and Health, 2015a-e 2019". In: *Drug and Alcohol Dependence* 237 (ago.. 2022), p. 109511. ISSN: 03768716. DOI: 10.1016/j.drugalcdep.2022.109511. https://www.naaddc.org/assets/24/14/Written%20recoveryremission_from_substance_abuse_disorders.pdf (visited on 03/14/2023).
- [12] H. W. Andersson, M. Wenaas, and T. Nordfjærn. "Relapse after inpatient substance use treatment: A prospective cohort study among users of illicit substances". In: *Addictive Behaviors* 90 (mar.. 2019), pp. 222–228. ISSN: 03064603. DOI: 10.1016/j.addbeh.2018.11.008.
- [13] K. P. Rezaei-Zadeh, R. N. Engstrom, A. Sharma, et al. "Generational trends and patterns in readmission within a statewide cohort of clients receiving heroin use disorder treatment in Maryland, 2007a-e 2013". In: *Journal of Substance Abuse Treatment* 96 (2019), pp. 82–91. ISSN: 0740-5472. DOI: <https://doi.org/10.1016/j.jsat.2018.10.010>. URL: <https://www.sciencedirect.com/science/article/pii/S0740547218301399>.
- [14] C. Timko, A. Nash, M. D. Owens, et al. "Systematic Review of Criminal and Legal Involvement After Substance Use and Mental Health Treatment Among Veterans: Building Toward Needed Research". In: *Substance Abuse: Research and Treatment* 14 (ene.. 2020), p. 117822181950128. ISSN: 1178-2218. DOI: 10.1177/1178221819501281.
- [15] J. Levola, A. Aranko, and T. Pitkähnen. "Psychosocial difficulties and treatment retention in inpatient detoxification programmes". In: *Nordic Studies on Alcohol and Drugs* 38.5 (oct.. 2021), pp. 434–449. ISSN: 1455-0725. DOI: 10.1177/14550725211021263.
- [16] N. G. Choi and D. M. DiNitto. "Older-Adult Marijuana Users in Substance Use Treatment: Characteristics Associated with Treatment Completion". In: *Journal of Psychoactive Drugs* 52.3 (may.. 2020), pp. 218–227. ISSN: 0279-1072. DOI: 10.1080/02791072.2020.1745966.
- [17] H. W. Andersson, A. D. F. Laursen, and T. Nordfjærn. "Emerging Adults in Inpatient Substance Use Treatment: A Prospective Cohort Study of Patient Characteristics and Treatment Outcomes". In: *European addiction research* 27.3 (2021), pp. 206–215. ISSN: 1421-9891. DOI: 10.1159/000512156.
- [18] H. W. Andersson, A. Steinsbekk, E. Walderhaug, et al. "Predictors of Dropout From Inpatient Substance Use Treatment: A Prospective Cohort Study". In: *Substance Abuse: Research and Treatment* 12 (ene.. 2018), p. 117822181876055. ISSN: 1178-2218. DOI: 10.1177/1178221818760551.
- [19] D. Basu, A. Ghosh, S. Sarkar, et al. "Initial treatment dropout in patients with substance use disorders attending a tertiary care de-addiction centre in north India". In: *Indian Journal of Medical Research* 146.6 (2017), p. 77. ISSN: 0971-5916. DOI: 10.4103/ijmr.IJMR_1309_15.
- [20] K. Lalwani, P. Whitehorn-Smith, G. Walcott, et al. "Prevalence and sociodemographic factors associated with polysubstance use: analysis of a population-based survey in Jamaica". In: *BMC Psychiatry* 22.1 (dic.. 2022), p. 513. ISSN: 1471-244X. DOI: 10.1186/s12958-022-04160-2.
- [21] J. C. Reyes, C. M. Perez, H. M. Colon, et al. "Prevalence and Patterns of Polydrug Use in Latin America: Analysis of Population-based Surveys in Six Countries". In: *Review of European Studies* 5.1 (feb.. 2013). ISSN: 1918-7181. DOI: 10.5539/res.v5n1p10.
- [22] R. Santis, B. C. G. Hidalgo, C. V. Hayden, et al. "Consumo de sustancias y conductas de riesgo en consumidores de pasta base de cocaína no consultantes a servicios de rehabilitación". In: *Revista médica de Chile* 135.1 (ene.. 2007). ISSN: 0034-9887. DOI: 10.4067/S0034-98872007000100007.
- [23] C. F. Olivari, J. Gaele, N. Rodriguez, et al. "Polydrug Use and Co-occurring Substance Use Disorders in a Respondent Driven Sampling of Cocaine Base Paste Users in Santiago, Chile". In: *Journal of Psychoactive Drugs* 54.4 (ago.. 2022), pp. 348–357. ISSN: 0279-1072. DOI: 10.1080/02791072.2021.1976886.
- [24] F. Vilgrón, T. Molina G., M. E. Gras-Pérez, et al. "Precocidad de inicio del consumo de sustancias psicoactivas y su relación con otros comportamientos de riesgo para la salud en adolescentes chilenos". In: *Revista médica de Chile* 150.5 (may.. 2022), pp. 584–596. ISSN: 0034-9887. DOI: 10.4067/S0034-98872022000500584.
- [25] S. R. Cole and M. A. Hernan. "Constructing Inverse Probability Weights for Marginal Structural Models". In: *American Journal of Epidemiology* 168.6 (jul.. 2008), pp. 656–664. ISSN: 0002-9262. DOI: 10.1093/aje/kwn164.
- [26] M. J. Crowther and P. Lambert. MULTISTATE: Stata module to perform multi-state survival analysis. ene.. 2023. URL: <https://econpapers.repec.org/RePEc:boc:bocode:s458207>.
- [27] L. A. Stefanski and D. D. Boos. "The Calculus of M-Estimation". In: *The American Statistician* 56.1 (feb.. 2002), pp. 29–38. ISSN: 0003-1305. DOI: 10.1198/000313002753631330.
- [28] T. J. VanderWeele. "Causal Mediation Analysis With Survival Data". In: *Epidemiology* 22.4 (jul.. 2011), pp. 582–585. ISSN: 1044-3983. DOI: 10.1097/EDE.0b013e31821db37e. <https://ideas.repec.org/c/boc/bocode/s457128.html>.
- [29] P. Lambert. STPM2: Stata module to estimate flexible parametric survival models. Statistical Software Components, Boston College Department of Economics. feb.. 2010. URL: <https://ideas.repec.org/c/boc/bocode/s457128.html>.
- [30] M. Hill. "Development and application of methods in parametric survival models: interval censoring, inverse probability weighting and multistate survival models". University of Leicester, 2022. URL: <https://doi.org/10.26392/leicester.data.2153514.v1>.

Funding sources

- This work was funded by ANID - Millennium Science Initiative Program - N° NCS2021_003 (Castillo-Carniglia) and N° NCS2021_013 (Calvo); The authors have no conflict of interest to declare
- **Correspondence to:** Andrés González-Santa Cruz, gonzalez.santacruz.andres@gmail.com