


Assessing the impact of substance use treatment for preventing criminal justice system contact in Chile



M. Mateo Piñones^{1,2, }, A. González-Santa Cruz^{1,3, }, A. Castillo-Carniglia^{4,5, }

¹ Young Researcher, Millennium Nucleus for the evaluation and analysis of Drug Policies

² Ph.D. student, School of Public Health, Universidad de Chile

³ Ph.D. student, Griffith University, Australia

⁴ Director, Millennium Nucleus for the evaluation and analysis of Drug Policies

⁵ Associate Professor, Society & Health Research Center



Background

Research has shown that reducing SUDs through effective treatment leads to a reduction in criminal activity^[1]. However, most evidence comes from developed countries, and results from the Latin American context are largely unknown^[2]. The social, cultural, economic, epidemiological context and substance use treatment (SUT) policy response are different in this region, making the question about SUT effectiveness through locally based data relevant^[3]. **We analyzed Chile as a case study and examined the impact of SUT on the prevention of contact with the criminal justice system (CJS) in the middle (3 years) and long term (5 years).**

Methods

- **Design:** a population-based record-linkage retrospective cohort design.
- **Participants:** patients enrolled in publicly funded SUT programs in Chile, 2010-2019.
- **Exposure:** Treatment completion, Late (≥ 3 months of treatment) & Early dropout (< 3 months); **Outcome:** contact with the CJS (offenses ending with a condemnatory sentence and offenses that ended with imprisonment after baseline treatment outcome)

Analysis plan

- We calculated cumulative incidence rates and incidence rate ratios (IRR)
- We used Royston-Parmar survival models and adjust for the effects of other factors, and predicted standardized survival probabilities and restricted mean survival times (RMST)^[4].
- We imputed missing data under MAR assumption^[5].
- We calculated e-values of the strength of confounding needed to take away the associations between treatment outcome and contact with CJS
- Codes are available at https://fondecytacc.github.io/nDP/index_prop_grant22_23.html.

Covariates are listed below:

- | | |
|---|---|
| <ul style="list-style-type: none">• Treatment setting• Substance use onset age• Primary substance at admission• Occupational status• Number of children (binary)• Macrozone• Number of previous offenses (acquisitive)• Number of previous offenses (other)• Substance use severity (dependence status) (ICD-10)• Percentage of poverty of the municipality of residence• Birth year• Physical comorbidity | <ul style="list-style-type: none">• Sex• Educational attainment• Primary substance at admission usage frequency• Poly-substance use• Tenure status of households• Number of previous offenses (violent)• Number of previous offenses (substance-related)• Psychiatric comorbidity (ICD-10)• Urban/rural municipality of residence• Initial substance• Cohabitation status• Admission Age |
|---|---|

Preliminary Results

Of the 109,756 (patients = 85,048) SENDA records of admissions, 70,863 (83%) were eligible to be matched with the Prosecutor's Office database (discarded ongoing treatments or treatments that ended in referrals).

- **Condemnatory sentence.** 22,287 (31%).
- **Imprisonment.** 5,144 (7%).

- **Early dropout vs. Treatment completion:** Patients completing treatment took longer to an offense leading to condemnatory sentence (IRR = 2.18 95% CI 2.09, 2.27; aHR [adjusted hazard ratio]: 1.74 95% CI 1.66, 1.83) condemnatory sentence and imprisonment (IRR = 2.90 95% CI 2.64, 3.18; aHR = 1.99 95% CI 1.79, 2.22).
- **Late dropout vs. Treatment completion:** Patients completing treatment took longer to condemnatory sentence (IRR = 1.73 95% CI 1.67, 1.80; aHR = 1.58 95% CI 1.52, 1.65) and to imprisonment (IRR = 1.93 95% CI 1.77, 2.10; aHR = 1.65 95% CI 1.51, 1.81).
- **Late vs. Early dropout:** Patients completing treatment took longer to condemnatory sentence (IRR = 1.26 95% CI 1.22, 1.30) and imprisonment (IRR = 1.50 95% CI 1.41, 1.61).

E-values

- **Condemnatory Sentence:** E-value of at least 2.19 for Early and 2.01 for Late dropout vs. treatment completion at baseline ($t=0$).
- **Imprisonment:** E-value of at least 2.99 for Early and 2.38 for Late dropout vs. treatment completion at baseline ($t=0$).

The following figure depicts the predicted differences in survival probabilities and RMTLs for committing an offense that results in a condemnatory sentence and imprisonment.




Figure 1: Differences in survival probabilities (left) and RMSTs (right) for time-to-condemnatory sentence (up) & imprisonment (bottom)

Discussion

- SUT can be effective in preventing contact with the criminal justice system (CJS) in Chile.
- More research is needed to understand the effects of SUT on contact with the CJS.
- Completing SUT can help to reduce the harms of substance use disorders and criminality.

References

- [1] M. Prendergast, D. Podus, E. Chang, et al. "Erratum to The effectiveness of drug abuse treatment: a meta-analysis of comparison group studies". In: *Drug and Alcohol Dependence* - DRUG ALCOHOL DEPENDENCE 84 (sept.. 2006), pp. 133-133. DOI: 10.1016/j.drugaicdep.2006.02.002.
- [2] H. Klingemann. "Successes and Failures in Treatment of Substance Abuse: Treatment System Perspectives and Lessons from the European Continent". In: *Nordisk Alkohol- and Narkotikatidskrift* 37.4 (2020), pp. 323-37.
- [3] M. Mateo Pinones, A. González-Santa Cruz, R. Portilla Huidobro, et al. "Evidence-based policymaking: Lessons from the Chilean Substance Use Treatment Policy". En. In: *Int. J. Drug Policy* 109.103860 (nov.. 2022), p. 103860.
- [4] P. Lambert. *STPM2: Stata module to estimate flexible parametric survival models*. Statistical Software Components, Boston College Department of Economics. feb.. 2010. URL: <https://ideas.repec.org/c/boc/bocode/s457128.html>.
- [5] M. Mayer. "missRanger: Fast Imputation of Missing Values". (2023). R package version 2.2.0. URL: <https://github.com/mayer79/missRanger>.

Funding sources

- This work was funded by ANID - Millennium Science Initiative Program - N° NCS2021_003 (Castillo-Carniglia) and N° NCS2021_013 (Calvo); Authors have no conflict of interest to declare