

# High mortality in SUT Patients, particularly among women, and regional variations in mortality risks

## Mortality following substance use disorder treatment: population-based record-linkage retrospective cohort design

A. González-Santa Cruz<sup>1,2</sup>,   
gonzalez.santacruz.andres@gmail.com

A. Castillo-Carniglia<sup>1,3</sup>,   
alvacasti@gmail.com

<sup>1</sup> Millennium Nucleus for the evaluation and analysis of Drug Policies

<sup>2</sup> Ph.D. student, School of Public Health, Universidad de Chile

<sup>3</sup> Department of Public Health, Facultad de Medicina y Ciencia, Universidad San Sebastián, Chile

## Background

- Chile has one of the highest alcohol consumption in the continent<sup>[1]</sup> and one of the highest rates of high school cocaine, coca paste, and marijuana use in the Americas<sup>[2]</sup>
- Had over 170% growth in drug-related mortality, 2000 vs 2019<sup>[3]</sup>
- In 2020, 6% of Chileans (12-64) had a problematic substance (alcohol & drug) use<sup>[4]</sup>
- Chilean government funds substance use treatments (SUT) for all with public health insurance (~81% of the population)<sup>[5]</sup>
- There is limited information regarding short-term, medium-term, and long-term mortality risks
- Analyzing mortality among administrative regions may reveal disparities in post-treatment outcomes
- Regional data can inform policy and resource allocation

## Objectives

To describe the standardized mortality ratios (SMRs) for all adult patients in publicly funded SUT within 2010-2022, along with regional specificities

## Methods

**Design:** a population-based retrospective cohort of adults enrolled in Chilean SUT programs (18-65) with national mortality data (2010-2020).

### Analysis plan:

- Calculated SMRs by comparing observed vs. expected deaths within regions, age groups, periods, and sexes
- Stratum-specific population estimates obtained from National Institute of Statistics projections<sup>[6]</sup>
- Regional data on substance use from 2020 Chile National Drug Study by National Drug and Alcohol Prevention and Rehabilitation Service<sup>[4]</sup>
- Calculated 95% confidence intervals using Vandenbroucke's method<sup>[7]</sup>

## Preliminary Results




Figure 1: Characteristics of SUT patients




Figure 2: Standardized mortality ratios for people in SUT, by age group and sex




Figure 3: Regional data in quantiles

- The SMR in the SUT population was 15.5 (95%CI 15.0, 16.1) times higher than expected based on the general population.
- Women aged between 30-44 had a mortality risk up to 29.6 times (95%CI 26.0, 33.5) higher than expected.
- Regions with high SMRs for men also have high SMRs for women.
- Ñuble showed some intervals with zero values due to its establishment in 2018.
- There were regional variations in mortality risks post-SUT.
- The regional prevalence of problematic substance use does not appear to be related to SMRs.

## Discussion

- High cumulative mortality risk in SUT patients vs. general population
- Higher risk particularly among women
- Largest sex gap at younger ages<sup>[8]</sup>
- Further exploration of substance use's contribution to elevated risk
- Mortality risks were higher in areas influenced by ethnic and cultural barriers, discrimination, or geographical isolation, which hinder access to treatment<sup>[9;10]</sup>
- Future studies should address the causes of these disparities beyond aggregated problematic substance use

## References

- B. Vicente, S. Saldivia, and R. Pihán. "Prevalencias y brechas hoy: salud mental mañana". In: *Acta bioethica* 22 (2016), pp. 51–61. ISSN: 1726-569X. DOI: [10.4067/S1726-569X2016000100006](https://doi.org/10.4067/S1726-569X2016000100006).
- Inter-American Drug Abuse Control Commission [CICAD]. Report on Drug Use in the Americas 2019: Executive Summary. Tech. rep. Washington, DC, 2019. URL: <http://www.cicad.oas.org/cicaddocs/Document.aspx?id=4976>.
- OECD and World Bank. Health at a Glance: Latin America and the Caribbean 2023. 2023. DOI: <https://doi.org/10.1787/532b0e2d-en>. URL: <https://www.oecd-ilibrary.org/content/publication/532b0e2d-en>.
- Servicio Nacional para la Prevención y Rehabilitación del Consumo de Drogas y Alcohol [SENDA]. Décimo Cuarto Estudio Nacional de Drogas en Población General de Chile, 2020. Tech. rep. Santiago, Chile: Ministerio del Interior y Seguridad Pública, 2022. URL: <https://www.senda.gob.cl/wp-content/uploads/2022/01/Estudio-P62020.pdf>.
- M. Mateo-Pinones, A. González-Santa Cruz, R. Portilla Huidobro, et al. "Evidence-based policymaking: Lessons from the Chilean Substance Use Treatment Policy". In: *International Journal of Drug Policy* 109 (nov. 2022), p. 103860. ISSN: 0955-3959. DOI: [10.1016/j.drugpo.2022.103860](https://doi.org/10.1016/j.drugpo.2022.103860). URL: <https://www.sciencedirect.com/science/article/pii/S0955395922002766>.
- Instituto Nacional de Estadísticas [INE]. Tablas de Mortalidad de Chile 1992-2050: Documento Metodológico. ene. 2023. URL: [https://www.ine.gob.cl/docs/default-source/proyecciones-de-poblacion/metodologia/proyecci%C3%B3n-base-2017/tablas-de-mortalidad-de-chile-1992-2050-metodologia.pdf?sfvrsn=ff4b0f8a\\_5](https://www.ine.gob.cl/docs/default-source/proyecciones-de-poblacion/metodologia/proyecci%C3%B3n-base-2017/tablas-de-mortalidad-de-chile-1992-2050-metodologia.pdf?sfvrsn=ff4b0f8a_5).
- M. Szklo and F. J. Nieto. *Epidemiology : beyond the basics*. Fourth edition. Burlington, Mass: Jones & Bartlett Learning, 2019. ISBN: 978-1-4496-0469-1.
- Á. Castillo-Carniglia, J. S. Kaufman, and P. Pino. "Geographical distribution of alcohol-attributable mortality in Chile: A Bayesian spatial analysis". In: *Addictive Behaviors* 42 (mar. 2015), p. 207â€"215. ISSN: 0306-4603. DOI: [10.1016/j.addbeh.2014.11.025](https://doi.org/10.1016/j.addbeh.2014.11.025). URL: <http://dx.doi.org/10.1016/j.addbeh.2014.11.025>.
- F. Mascayano, T. Tapia, S. Schilling, et al. "Stigma toward mental illness in Latin America and the Caribbean: a systematic review". In: *Revista Brasileira de Psiquiatria* 38.1 (mar. 2016), p. 73â€"85. ISSN: 1516-4446. DOI: [10.1590/1516-4446-2015-1652](https://doi.org/10.1590/1516-4446-2015-1652). URL: <http://dx.doi.org/10.1590/1516-4446-2015-1652>.
- J. C. Sapag, B. F. Sena, I. V. Bustamante, et al. "Stigma towards mental illness and substance use issues in primary health care: Challenges and opportunities for Latin America". In: *Global Public Health* 13.10 (ago. 2017), p. 1468â€"1480. ISSN: 1744-1706. DOI: [10.1080/17441692.2017.1356347](https://doi.org/10.1080/17441692.2017.1356347). URL: <http://dx.doi.org/10.1080/17441692.2017.1356347>.

## Funding sources

- This work was funded by ANID - Millennium Science Initiative Program - N° NCS2021\_003 (Castillo-Carniglia) and ANID - Doctoral Scholarship National Doctorate/2023-21230172 (González-Santa Cruz)

Correspondence to: Andrés González-Santa Cruz,  
gonzalez.santacruz.andres@gmail.com

