

R Textbook Companion for
Introduction to Probability and Statistics for
Engineers and Scientists
by Sheldon M. Ross¹

Created by
Bhushan Manjarekar
B.E.
Electronics Engineering
Mumbai University
Cross-Checked by
R TBC Team

May 22, 2020

¹Funded by a grant from the National Mission on Education through ICT
- <http://spoken-tutorial.org/NMEICT-Intro>. This Textbook Companion and R
codes written in it can be downloaded from the "Textbook Companion Project"
section at the website - <https://r.fossee.in>.

Book Description

Title: Introduction to Probability and Statistics for Engineers and Scientists

Author: Sheldon M. Ross

Publisher: Elsevier Academic Press, USA

Edition: 3

Year: 2004

ISBN: 0-12-598057-4

R numbering policy used in this document and the relation to the above book.

Exa Example (Solved example)

Eqn Equation (Particular equation of the above book)

For example, Exa 3.51 means solved example 3.51 of this book. Sec 2.3 means an R code whose theory is explained in Section 2.3 of the book.

Contents

List of R Codes	4
2 Descriptive Statistics	5
3 Elements Of Probability	12
4 Random Variables And Expectation	19
5 Special Random Variables	37
6 Distribution of Sampling Statistics	46
7 Parameter Estimation	49
8 Hypothesis Testing	62
9 Regression	73
10 Analysis of Variance	91
11 Goodness of Fit Tests and Categorical Data Analysis	97
12 Non parametric Hypothesis Tests	106
13 Quality Control	115
14 Life Testing	123

List of R Codes

Exa 2.2.a	Relative Frequency	5
Exa 2.2.b	Pie Chart	5
Exa 2.3.a	Sample mean	6
Exa 2.3.b	Sample mean of age	6
Exa 2.3.c	Sample Median	6
Exa 2.3.d	Mean and Median	7
Exa 2.3.e	Mean Median and Mode	7
Exa 2.3.f	sample variance	8
Exa 2.3.g	sample variance of accidents	8
Exa 2.3.h	Percentile	8
Exa 2.3.i	Quartiles	9
Exa 2.4.a	Chebyshev Inequality	9
Exa 2.5.a	Empirical Rule	10
Exa 2.6.a	Sample Correlation Coefficient	10
Exa 2.6.b	Sample Correlation Coefficient	11
Exa 3.4.a	Union	12
Exa 3.5.a	Basic Principle of Counting	12
Exa 3.5.b	Basic Principle of Counting	13
Exa 3.5.c	Basic Principle of Counting	13
Exa 3.5.d	Committee Probability	13
Exa 3.5.f	Pairing Probability	14
Exa 3.6.a	Acceptable Transistor	15
Exa 3.6.b	Both Boys	15
Exa 3.6.c	Branch Manager	15
Exa 3.7.a	Accident Probability	16
Exa 3.7.b	Accident within a year	16
Exa 3.7.c	Multiple Choice Test	16
Exa 3.7.d	blood test	16

Exa 3.7.e	Criminal Investigation	17
Exa 3.7.f	Missing Plane	17
Exa 3.8.a	Independent Events	18
Exa 4.1.a	sum of two fair dice	19
Exa 4.1.c	X exceeds 1	20
Exa 4.2.a	sum of pmf	21
Exa 4.2.b	pdf	21
Exa 4.3.a	Joint distribution of batteries	22
Exa 4.3.b	Joint distribution of boys and girls	23
Exa 4.3.c	Joint Density Function	24
Exa 4.3.d	Density of Independent Random Variables	26
Exa 4.3.e	Density of Independent Random Variables	26
Exa 4.3.f	Conditional Probability Mass Function	27
Exa 4.3.g	Conditional Probability Mass Function	28
Exa 4.4.a	Expectation of a fair die	29
Exa 4.4.d	Expectation of the message time	29
Exa 4.5.a	Expectation	30
Exa 4.5.b	Expected cost of breakdown	31
Exa 4.5.c	Expectation	31
Exa 4.5.d	Expectation	32
Exa 4.5.e	Expected profit	32
Exa 4.5.g	Different types of coupons	33
Exa 4.6.a	Variance of a fair die	34
Exa 4.7.a	Variance of 10 rolls of a fair die	34
Exa 4.7.b	Variance of 10 tosses of a coin	35
Exa 4.9.a	Inequalities	36
Exa 5.1.a	Returning of disks	37
Exa 5.1.b	Colour of Eyes	37
Exa 5.1.e	Binomial Random Variable	38
Exa 5.1.f	Binomial Distribution	38
Exa 5.2.a	Probability of accident	39
Exa 5.2.b	Defective Items	39
Exa 5.2.c	Number of Alpha particles	39
Exa 5.2.d	Claims handled by an insurance company	39
Exa 5.2.f	Defective stereos	40
Exa 5.3.a	Functional system	40
Exa 5.3.b	Determining Population Size	40
Exa 5.3.c	Conditional Probability	40

Exa 5.4.a	Probabilities of a uniform random variable	41
Exa 5.4.b	Bus Timings	41
Exa 5.4.c	Current in a diode	42
Exa 5.5.a	Normal Random Variable	42
Exa 5.5.b	Noise in Binary Message	43
Exa 5.5.c	Power dissipation	43
Exa 5.5.d	Yearly precipitation	43
Exa 5.6.a	Wearing of Battery	43
Exa 5.6.b	Working Machines	44
Exa 5.6.c	Series System	44
Exa 5.8.a	Chi square random variable	44
Exa 5.8.b	Chi square random variable	44
Exa 5.8.c	Locating a Target	45
Exa 5.8.d	Locating a Target in 2D space	45
Exa 5.8.e	T distribution	45
Exa 5.8.f	F Distribution	45
Exa 6.3.a	Claims handled by an insurance company	46
Exa 6.3.c	Class strength	46
Exa 6.3.d	Weights of workers	46
Exa 6.3.e	Distance of a start	47
Exa 6.5.a	Processing time	47
Exa 6.6.a	Candidate winning an election	47
Exa 6.6.b	Pork consumption	48
Exa 7.2.a	Maximum likelihood estimator of a bernoulli parameter	49
Exa 7.2.b	Errors in a manuscript	49
Exa 7.2.c	Maximum likelihood estimator of a poisson parameter	49
Exa 7.2.d	Number of traffic accidents	50
Exa 7.2.e	Maximum likelihood estimator in a normal population	50
Exa 7.2.f	Kolmogorovs law of fragmentation	50
Exa 7.2.g	Estimating Mean of a Uniform Distribution	51
Exa 7.3.a	Error in a signal	51
Exa 7.3.b	Confidence interval	52
Exa 7.3.c	Confidence interval	52
Exa 7.3.d	Weight of a salmon	53
Exa 7.3.e	Error in a signal	53
Exa 7.3.f	Average resting pulse	54
Exa 7.3.g	Evaluating integrals	54
Exa 7.3.h	Thickness of washers	55

Exa 7.4.a	Cable insulation	56
Exa 7.4.b	Battery production	56
Exa 7.5.a	Transistors	57
Exa 7.5.b	Survey	58
Exa 7.5.c	Acceptable chips	58
Exa 7.6.a	Life of a product	59
Exa 7.7.a	Point estimator	59
Exa 7.7.b	Point estimator	59
Exa 7.7.c	Point estimator of a uniform distribution	60
Exa 7.8.a	Bayes estimator	60
Exa 7.8.b	Bayes estimator of a normal population	60
Exa 7.8.d	estimator of the signal value	61
Exa 8.3.a	Noise in a Signal	62
Exa 8.3.b	Error in a signal	62
Exa 8.3.c	Error in a signal	63
Exa 8.3.d	Number of signals to be sent	63
Exa 8.3.e	Number of signals to be sent	64
Exa 8.3.f	Nicotine content in a cigarette	64
Exa 8.3.g	Blood cholesterol level	65
Exa 8.3.h	Water usage	65
Exa 8.3.i	Life of a tire	66
Exa 8.3.j	Service Time	66
Exa 8.4.a	Tire lives	67
Exa 8.4.b	Medicine for cold	67
Exa 8.4.c	Unknown population variance	68
Exa 8.4.d	effectiveness of safety program	69
Exa 8.5.a	effectiveness of machine	69
Exa 8.5.b	Catalyst	69
Exa 8.6.a	Computer chip manufacturing	70
Exa 8.6.b	Finding p value	70
Exa 8.6.c	Change in manufacturing pattern	71
Exa 8.7.a	Mean number of defective chips	71
Exa 8.7.b	Safety Conditions in a plant	71
Exa 8.7.c	Better proof reader	72
Exa 9.1.a	Scatter Diagram	73
Exa 9.2.a	Relative humidity and moisture content	73
Exa 9.3.a	Moisture against Density	74
Exa 9.4.a	Effect of speed on mileage	75

Exa 9.4.b	Confidence interval estimate	76
Exa 9.4.c	Regression to the mean	77
Exa 9.4.d	Motor vehicle deaths	78
Exa 9.4.e	Confidence interval for height	79
Exa 9.4.f	Confidence interval for height	80
Exa 9.5.a	Height of son and father	81
Exa 9.7.a	Percentage of chemical used	82
Exa 9.8.b	Distance vs Travel Time	83
Exa 9.9.a	Polynomial Fitting	84
Exa 9.10.a	Multiple Linear Regression	85
Exa 9.10.b	Estimate of variance	86
Exa 9.10.c	Diameter of a tree	87
Exa 9.10.d	Estimating hardness	89
Exa 9.11.a	Animal sick falling	90
Exa 10.3.a	Dependence of mileage on gas used	91
Exa 10.3.b	Dependence of mileage on gas used	92
Exa 10.3.c	Difference in GPA	93
Exa 10.4.b	Estimating Parameters	94
Exa 10.5.a	Species collected	95
Exa 11.2.a	Relation between death date and birth date	97
Exa 11.2.b	Quality of bulbs	98
Exa 11.2.d	Six outcomes	98
Exa 11.3.a	Weekly accidents	99
Exa 11.4.a	Political affiliation and Gender	100
Exa 11.4.b	Machine Breakdown and shift	101
Exa 11.5.a	Lung cancer and smoking	102
Exa 11.5.b	Females reporting abuse	104
Exa 11.6.a	Testing distribution of a population	105
Exa 12.2.a	testing the median	106
Exa 12.2.b	testing the median	106
Exa 12.3.b	Signed Rank Test	107
Exa 12.3.c	Determining Population Distribution	107
Exa 12.4.a	Treatments against corrosion	108
Exa 12.4.b	Determining P	109
Exa 12.4.c	Finding p value	110
Exa 12.4.d	Comparing production methods	111
Exa 12.4.e	Determining p value	112
Exa 12.5.a	Testing randomness	112

Exa 12.5.c Determining p value	114
Exa 13.2.a Steel shaft diameter	115
Exa 13.2.b unknown mean and variance	116
Exa 13.3.a determining control limits	116
Exa 13.4.a Defectives Screws	117
Exa 13.5.a Control during production of cars	118
Exa 13.6.b Service Time	119
Exa 13.6.c Exponentially weighted moving average control	120
Exa 13.6.d Finding control limit	121
Exa 14.3.a Lifetime of a transistor	123
Exa 14.3.b Lifetime of Battery	123
Exa 14.3.c One at a time sequential test	124
Exa 14.3.d Lifetime of semiconductors	124
Exa 14.3.e Bayes estimator	125
Exa 14.4.a Lifetime of items produced by two plants	125

Chapter 2

Descriptive Statistics

R code Exa 2.2.a Relative Frequency

```
1 starting_salary<-c(47, 48, 49 ,50 ,51, 52, 53, 54,
  56, 57, 60)
2 frequency <-c(4, 1 ,3 ,5 ,8 ,10, 0 ,5 ,2, 3, 1)
3 total = sum(frequency)
4 relative_frequency = frequency/total
5 cat("The relative frequencies are ",relative_
  frequency)
```

R code Exa 2.2.b Pie Chart

```
1 slices<-c(42, 50, 32, 55, 9 ,12)
2 lbls<-c("Lung ", "Breast ", "Colon ", "Prostate ", "
  Melanoma ", "Bladder ")
3 pct <- round(slices/sum(slices)*100)
4 lbls <- paste(lbls, pct) # add percents to labels
5 lbls <- paste(lbls,"%",sep="") # ad % to labels
6 pie(slices,labels = lbls, col=rainbow(length(lbls)),
  main="Pie Chart")
```

R code Exa 2.3.a Sample mean

```
1 scores<-c(284, 280, 277, 282, 279, 285, 281, 283,  
            278, 277)  
2 new_scores=(scores - 280)  
3 final_mean = mean(new_scores)+ 280  
4 cat("Final Mean = ", final_mean)
```

R code Exa 2.3.b Sample mean of age

```
1 age<-c(15, 16, 17, 18, 19, 20)  
2 frequencies<-c(2, 5, 11, 9 ,14, 13)  
3 product  = age*frequencies  
4 total_people = sum(frequencies)  
5 mean_age = sum(product)/total_people  
6 cat("The sample mean of the ages is",mean_age)
```

R code Exa 2.3.c Sample Median

```
1 age<-c(15, 16, 17, 18, 19, 20)  
2 frequencies<-c(2, 5, 11, 9 ,14 ,13)  
3 final_age = matrix(0,1,length(age))  
4 i=1  
5 for (j in 1:6){  
6     for (k in 1:frequencies[j]){  
7         final_age[i] = age[j]  
8         i = i +1  
9     }  
10 }
```

```
11 final_median = median(final_age)
12 final_median
```

R code Exa 2.3.d Mean and Median

```
1 germ_free_mice<-c(158, 192, 193, 194, 195, 202, 212,
  215, 229, 230, 237, 240, 244, 247, 259, 301,
  301, 321, 337, 415 ,434 ,444 ,485 ,496 ,529 ,537,
  624, 707 ,800)
2 conventional_mice<-c(159, 189, 191, 198, 235 ,245,
  250, 256, 261, 265, 266 ,280 ,343 ,356, 383, 403,
  414 ,428 ,432)
3 cat("Sample mean for germ-free mice is ",mean(germ_
  free_mice))
4 cat("Sample median for germ-free mice is ",median(
  germ_free_mice))
5 cat("Sample mean for conventional mice is ",mean(
  conventional_mice))
6 cat("Sample median for conventional mice is ",median(
  conventional_mice))
```

R code Exa 2.3.e Mean Median and Mode

```
1 value<-c(1, 2, 3, 4, 5, 6)
2 frequencies<-c(9, 8 ,5 ,5, 6 ,7)
3 final_value = matrix(0,1,length(value))
4 i=1
5 for (j in 1:6){
6   for (k in 1:frequencies[j]){
7     final_value[i] = value[j]
8     i = i +1
9   }
10 }
```

```

11 product  = value*frequencies;
12 product
13 sum(product)
14
15 total_value = sum(frequencies);
16 mean_value = sum(product)/total_value #the answer in
    the textbook is incorrect
17 which(frequencies==max(frequencies))
18 m1= max(frequencies);
19 n= which(frequencies==max(frequencies))
20
21 cat("The sample mean is",mean_value)
22 cat("The median is",median(final_value))
23 cat("The mode is",value[n])

```

R code Exa 2.3.f sample variance

```

1 A <-c(3, 4, 6, 7, 10)
2 B <-c(-20, 5 ,15 ,24)
3 cat("The sample variance of A is",var(A))
4 cat("The sample variance of B is",var(B))

```

R code Exa 2.3.g sample variance of accidents

```

1 accidents<-c(22, 22, 26, 28, 27, 25 ,30, 29, 24)
2 new_accidents = accidents - 22
3 cat("The variance of the number of accidents is",var
    (new_accidents))

```

R code Exa 2.3.h Percentile

```

1 population<-c(7333253, 3448613, 2731743, 1702086,
  1524249, 1151977, 1048949 ,1022830 ,998905,
  992038, 816884, 752279, 734676, 702979, 665070,
  635913, 617044 ,614289 ,579307, 567094, 547727
  ,520947 ,514013, 504505, 493559)
2 cat("The sample 10 percentile is",quantile(
  population, .10))
3 cat("The sample 50 percentile is",quantile(
  population, .50))
4 cat("The sample 80 percentile is",quantile(
  population, .80))
5 cat("The median is",median(population))

```

R code Exa 2.3.i Quartiles

```

1 noise<-c(82, 89, 94, 110, 74, 122, 112, 95, 100,
  78, 65, 60, 90 ,83 ,87 ,75 ,114 ,85 ,69 ,94 ,124
  ,115 ,107 ,88 ,97 ,74 ,72 ,68 ,83 ,91 ,90 ,102,
  77, 125, 108 ,65)
2 cat("The quartiles are",quantile(noise))

```

R code Exa 2.4.a Chebyshev Inequality

```

1 cars<-c(448162, 404192, 368327, 318308, 272122,
  260486, 249128, 234936, 218540, 207977)
2 interval1 = mean(cars) - (1.5*sd(cars));
3 interval2 = mean(cars) + (1.5*sd(cars));
4 data = 100*5/9;
5 cat("Atleast 55.55% of the data lies in the interval
  ",interval1 ,"to",interval2)

```

R code Exa 2.5.a Empirical Rule

```
1 data<-c(90, 91 ,94 ,83 ,85 ,85 ,87 ,88 ,72, 74, 74,
          75, 77, 77, 78, 60, 62 ,63, 64, 66, 66, 52 ,55
          ,55 ,56 ,58 ,43, 46)
2 cat("According to the empirical rule 68% of the data
      lies between",mean(data)-sd(data),"and",mean(
      data)+sd(data))
3 cat("95% of the data lies between",mean(data)-(2*sd(
      data)),"and",mean(data)+(2*sd(data)))
4 cat("99.7% of the data lies between", mean(data)-(3*
      sd(data)),"and",mean(data)+(3*sd(data)))
```

R code Exa 2.6.a Sample Correlation Coefficient

```
1 temp<-c(24.2, 22.7 ,30.5, 28.6, 25.5, 32.0, 28.6,
          26.5, 25.3, 26.0, 24.4 ,24.8 ,20.6, 25.1, 21.4,
          23.7, 23.9, 25.2, 27.4, 28.3 ,28.8 ,26.6)
2 defects<-c(25, 31 ,36 ,33 ,19 ,24 ,27 ,25 ,16 ,14
            ,22 ,23 ,20 ,25 ,25 ,23 ,27 ,30 ,33 ,32 ,35, 24)
3 temp_new = temp- mean(temp)
4 defects_new = defects - mean(defects)
5 num=0
6 s1 =0;
7 s2=0;
8 for (i in 1:22){
9   num = num + (temp_new[i]*defects_new[i])
10   s1 = s1 + (temp_new[i]*temp_new[i])
11   s2 = s2 + (defects_new[i]*defects_new[i])
12 }
13 coefficient = num/sqrt(s1*s2)
14 coefficient
```

R code Exa 2.6.b Sample Correlation Coefficient

```
1 year<-c(12, 16, 13, 18, 19 ,12, 18, 19, 12, 14)
2 pulserate<-c(73, 67, 74, 63 ,73 ,84 ,60 ,62, 76, 71)
3 year_new = year- mean(year)
4 pulserate_new = pulserate - mean(pulserate)
5 num=0
6 s1 =0
7 s2=0
8 for (i in 1:10){
9   num = num + (year_new[i]*pulserate_new[i])
10  s1 = s1 + (year_new[i]*year_new[i])
11  s2 = s2 + (pulserate_new[i]*pulserate_new[i])
12 }
13 coefficient = num/sqrt(s1*s2)
14 coefficient
```

Chapter 3

Elements Of Probability

R code Exa 3.4.a Union

```
1 cigarette = 0.28;
2 cigar = 0.07;
3 cigar_and_cigarette = 0.05 ;
4 cigar_or_cigarette = cigarette + cigar - cigar_and_
  cigarette;
5 cat((1-cigar_or_cigarette)*100 ,"% of the males
  smoke neither cigar nor cigarette")
```

R code Exa 3.5.a Basic Principle of Counting

```
1 white_balls= 6;
2 black_balls = 5;
3 total = white_balls + black_balls;
4 probability_whiteandblack = white_balls*black_balls/
  (total*(total-1));
5 probability_blackandwhite = white_balls*black_balls/
  (total*(total-1));
6 reqd_probability = probability_whiteandblack +
  probability_blackandwhite;
```

```
7 cat("Thus, the required probability is",reqd_  
    probability)
```

R code Exa 3.5.b Basic Principle of Counting

```
1 maths = 4;  
2 chemistry = 3;  
3 history = 2;  
4 language = 1;  
5 total_arrangements = factorial(4)*factorial(maths)*  
    factorial(chemistry)*factorial(history)*factorial  
    (language);  
6 cat("The total number of possible arrangements is ",  
    total_arrangements)
```

R code Exa 3.5.c Basic Principle of Counting

```
1 men = 6;  
2 women = 4;  
3 cat("No of different rankings possible is",factorial  
    (men+women))  
4 women_top4 = factorial(women)*factorial(men);  
5 prob = women_top4/factorial(men+women);  
6 cat("Probability that women receive the top 4 scores  
    is",prob)
```

R code Exa 3.5.d Committee Probability

```
1 men = 6;  
2 women = 9;
```

```

3 reqd_size =5;
4 total =factorial(men+women)/(factorial(reqd_size)*
    factorial(men+women-reqd_size));
5 given_committee = factorial(men)*factorial(women)/(
    factorial(3)*factorial(2)*factorial(men-3)*
    factorial(women-2));
6 prob = given_committee/total;
7 cat("Probability that the committee consists of 3
    men and 2 women is",prob)

```

R code Exa 3.5.f Pairing Probability

```

1 black_p = 6;
2 white_p = 6;
3 pair = 2;
4 total_p = black_p + white_p;
5
6 total_pairs = 1;
7 while(total_p >0){
8     total_pairs = total_pairs*factorial(total_p)/(
        factorial(pair) * factorial(total_p - pair) )
9     total_p = total_p -2
10 }
11 total_pairs= total_pairs/factorial(6);
12 black_pairs = 1;
13 while(black_p >0){
14     black_pairs = black_pairs*factorial(black_p)/((
        factorial(pair) * factorial(black_p - pair) ))
15     black_p = black_p -2
16 }
17 black_pairs= black_pairs/factorial(3);
18 white_pairs = black_pairs;
19 allowed_pairs = black_pairs * white_pairs;
20 probb = allowed_pairs/ total_pairs;
21 cat(" Probability that a random pairing will not

```

```
result in any of the white and black players  
rooming together is ",probb)
```

R code Exa 3.6.a Acceptable Transistor

```
1 defective =5;  
2 partially_defective = 10;  
3 acceptable = 25;  
4 cat("The required probability is",acceptable/(  
    acceptable+partially_defective))
```

R code Exa 3.6.b Both Boys

```
1 prob_bb = 0.25;  
2 prob_bg = 0.25;  
3 prob_gb = 0.25;  
4 prob_gg = 0.25;  
5 cat("Probability that both are boys is",prob_bb/(  
    prob_bg+prob_gb+prob_bb))
```

R code Exa 3.6.c Branch Manager

```
1 prob_phoenix = 0.3;  
2 prob_manager = 0.6;  
3 cat("Probability that Perez will be a Phoenix branch  
    office manager is",prob_phoenix*prob_manager)
```

R code Exa 3.7.a Accident Probability

```
1 accident_prone= 0.4;
2 nonaccident_prone= 0.2;
3 pop_accident = 0.3;
4 prob = pop_accident*accident_prone + (1-pop_accident
    )*nonaccident_prone;
5 cat("The required probability is ",prob)
```

R code Exa 3.7.b Accident within a year

```
1 accident_prone= 0.4;
2 nonaccident_prone= 0.2;
3 pop_accident = 0.3;
4 prob_of_accident = pop_accident*accident_prone + (1-
    pop_accident)*nonaccident_prone;
5 prob = pop_accident * accident_prone /prob_of_
    accident;
6 cat("The required probability is",prob)
```

R code Exa 3.7.c Multiple Choice Test

```
1 m = 5;
2 p =1/2;
3 cat("The required probability is", (m*p)/(1+((m-1)*p
    )))
```

R code Exa 3.7.d blood test

```
1 detect_present = 0.99;
```

```

2 detect_notpresent = 0.01;
3 pop_disease = 0.005;
4 prob = detect_present*pop_disease/((detect_present*
    pop_disease) +(detect_notpresent*(1-pop_disease))
    ) ;
5 cat("The required probability is",prob)

```

R code Exa 3.7.e Criminal Investigation

```

1 criminal_char = 0.9
2 convinced= 0.6;
3 pop_char = 0.2;
4 prob = (convinced*criminal_char) /((convinced*
    criminal_char) + (pop_char*(1-convinced)));
5 cat("The required probability is",prob)

```

R code Exa 3.7.f Missing Plane

```

1 alpha1 = 0.4;
2 plane_in_region1 = 1/3;
3 plane_in_region2 = 1/3;
4 plane_in_region3 = 1/3;
5 prob1 = (alpha1*plane_in_region1)/((alpha1*plane_in_
    region1)+ 1*plane_in_region2 + 1*plane_in_region3
    );
6 prob2 = (1*plane_in_region2)/((alpha1*plane_in_
    region1)+ 1*plane_in_region2 + 1*plane_in_region3
    );
7 cat("The probability that the planes is in region 1
    given that the search of region 1 did not uncover
    it ",prob1)

```

```
8 cat("The probability that the planes is in region 2/  
3 given that the search of region 1 did not  
uncover it ",prob2)
```

R code Exa 3.8.a Independent Events

```
1 prob_A = 4/52;  
2 prob_H = 13/52;  
3 cat("P(AH) is",prob_A*prob_H)
```

Chapter 4

Random Variables And Expectation

R code Exa 4.1.a sum of two fair dice

```
1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)
4 #read(IPSUR)
5 library(prob)
6 #read(prob)
7
8 #First we set up the sample space with the rolldie
  function
9 S <- rolldie(2, makespace = TRUE)
10 #Next, we add random variables U and V with the
    addrv function.
11 S <- addrv(S, FUN = max, invars = c("X1", "X2"),
    name = "U")
12 S <- addrv(S, FUN = sum, invars = c("X1", "X2"),
    name = "V")
13 head(S)
14 UV <- marginal(S, vars = c("U", "V"))
15 xtabs(round(probs, 3) ~ U + V, data = UV)
```

```

16 marginal(UV, vars = "U")
17 marginal(UV, vars = "V")
18
19 temp <- xtabs(probs ~ U + V, data = UV)
20 rowSums(temp)
21 colSums(temp)
22 colSums(temp)
23
24 '
25 data = sample(1:6,10000000,replace=TRUE)+sample
      (1:6,10000000,replace=TRUE)
26 table(data)/length(data)
27 sum(table(data)/length(data))
28
29 table(sample(1:6,10000000,replace=TRUE))/length(
      sample(1:6,10000000,replace=TRUE))
30 '

```

R code Exa 4.1.c X exceeds 1

```

1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)
4 #read(IPSUR)
5 library(prob)
6 #read(prob)
7
8 "
9  $F(x) = 0 \text{ --- } \{x \leq 0\}$ 
10  $F(x) = 1 - \exp(-x^2) \text{ --- } \{x > 0\}$ 
11
12  $P\{X > 1\} = ?$ 
13 "
14 P = 1 - (1 - exp(-(1^2)))
15 P

```

R code Exa 4.2.a sum of pmf

```
1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)
4 #read(IPSUR)
5 library(prob)
6 #read(prob)
7
8 P1 = 1/2
9 P2 = 1/3
10 P3 = 1 - P1 - P2
11 P3
```

R code Exa 4.2.b pdf

```
1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)
4 #read(IPSUR)
5 library(prob)
6 #read(prob)
7
8
9 integrand <- function(x) {(4*(x) - 2*(x^2))}
10 R = integrate(integrand, lower = 0, upper = 2)
11 C = 1/R$value
12 C
13
14 # P {X>1}
15 integrand <- function(x) {C*(4*(x) - 2*(x^2))}
```

```
16 P = integrate(integrand, lower = 1, upper = 2)
17 P
```

R code Exa 4.3.a Joint distribution of batteries

```
1 #install.packages("IP SUR")
2 #install.packages("prob")
3 library(IP SUR)
4 #read(IP SUR)
5 library(prob)
6 #read(prob)
7 #install.packages("combinat")
8 library(combinat)
9
10
11 p_0_0 = (length(combn(5, 3))/3)/(length(combn(12, 3))
12      )/3)
13 p_0_1 = (length(combn(4, 1))/1)*(length(combn(5, 2))
14      /2)/(length(combn(12, 3))/3)
15 p_0_1 = (length(combn(4, 2))/2)*(length(combn(5, 1))
16      /1)/(length(combn(12, 3))/3)
17 p_0_3 = (length(combn(4, 3))/3)/(length(combn(12, 3))
18      )/3)
19 p_1_0 = (length(combn(3, 1))/1)*(length(combn(5, 2))
20      /2)/(length(combn(12, 3))/3)
21 p_1_1 = (length(combn(3, 1))/1)*(length(combn(4, 1))
22      /1)*(length(combn(5, 1))/1)/(length(combn(12, 3))
23      /3)
24 p_1_2 = (length(combn(3, 1))/1)*(length(combn(4, 2))
25      /2)/(length(combn(12, 3))/3)
26 p_2_0 = (length(combn(3, 2))/2)*(length(combn(5, 1))
27      /1)/(length(combn(12, 3))/3)
28 p_2_1 = (length(combn(3, 2))/2)*(length(combn(4, 1))
29      /1)/(length(combn(12, 3))/3)
30 p_3_0 = (length(combn(3, 3))/3)/(length(combn(12, 3))
```

) / 3)

R code Exa 4.3.b Joint distribution of boys and girls

```
1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)
4 #read(IPSUR)
5 library(prob)
6 #read(prob)
7 #install.packages("combinat")
8 library(combinat)
9 library(discreteRV)
10 '
11 library(discreteRV)
12 d<-RV(c("Boy","Girl"),odds=c(0.5,0.5))
13 e<-RV(c("0","1","2","3"),odds = c
      (0.15,0.20,0.35,0.3))
14 f<-joint(d,e)
15 probs(f)
16 '
17
18 #P{B=0,G=0} = P{no children}
19 P = .15
20 P
21 #P{B=0,G=1} = P{1 Girl and total of 1 child} = P{1
      child} P{1 girl | 1 child}
22 P = .20 * (1/2)
23 P
24 #P{B=0,G=2} = P{2 Girls and total of 2 children} =
      P{2 children} P{2 girls | 2 children}
25 P = .35*(1/2)^2
26 P
27 #P{B=0,G=3} = P{3 Girls and total of 3 children} =
      P{3 children} P{3 girls | 3 children}
```

```

28 P = .3*(1/2)^3
29 P
30 #P{B=1,G=0} = P{1 Boy and total of 1 child} = P{1
    child} P{1 boy | 1 child}
31 P = .2*(1/2)
32 P
33 #P{B=1,G=1} = P{1 Boy, 1 Girl and total of 2 children
    } = P{2 children} P{1 boy | 2 children}
34 P = .35*.5
35 #P = .35*((1/2)^2+(1/2)^2)
36 P
37 #P{B=1,G=2} = P{1 Boy, 2 Girls and total of 3
    children} = P{2 children} P{2 Girls | 2 children
    + 1 boy|1 child}
38 P = .3*((1/2)^3+(1/2)^2)
39 P

```

R code Exa 4.3.c Joint Density Function

```

1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)
4 #read(IPSUR)
5 library(prob)
6 #read(prob)
7 #install.packages("combinat")
8 library(combinat)
9 library(discreteRV)
10 require(pracma)
11 library(cubature)
12 #1
13 fun0 = function(x,y,z) { exp(-x)*exp(-2*y)*z }
14 fun01 <- function(z) {
15     integrate(function(y) {
16         sapply(y, function(y) {

```

```

17         integrate(function(x) fun0(x,y,z), 1, Inf)$
           value
18     })
19 }, 0, 1)$value
20 }
21
22 fun01(2)
23 #(1-exp(-2))*exp(-1)
24
25 #2
26 fun0 = function(x,y,z) { exp(-x)*exp(-2*y)*z }
27 fun01 <- function(z) {
28     integrate(function(y) {
29         sapply(y, function(y) {
30             integrate(function(x) fun0(x,y,z), 0, y)$value
31         })
32     }, 0, Inf)$value
33 }
34
35 fun01(2)
36
37
38 ,
39 library(cubature)
40 f <- function(x) 2*exp(-x[1])*exp(-2*x[2])
41 adaptIntegrate(f, lowerLimit = c(1,0), upperLimit = c(
42     Inf,1))
43
44 require(pracma)
45 f <- function(x, y) 2*exp(-x)*exp(-2*y)
46 quad2d(f, 0, 1, 1, Inf)
47
48 http://stackoverflow.com/questions/23901682/double-
   integration-in-r-with-additional-argument

```

R code Exa 4.3.d Density of Independent Random Variables

```
1 #install.packages("IPSUR")
2 #install.packages("prob")
3 #install.packages("discreteRV")
4 #install.packages("pracma")
5 #install.packages("cubature")
6 library(IPSUR)
7 #read(IPSUR)
8 library(prob)
9 #read(prob)
10 #install.packages("combinat")
11 library(combinat)
12 library(discreteRV)
13 require(pracma)
14 library(cubature)
15
16 fun0 = function(x,y,a) { exp(-x)*exp(-y) }
17 fun01 <- function(z) {
18   integrate(function(y) {
19     sapply(y, function(y) {
20       integrate(function(x) fun0(x,y,z), 0, a*y)$
21         value
22     }, 0, Inf)$value
23   })
24
25 fun01(1)
26 fun01(2)
```

R code Exa 4.3.e Density of Independent Random Variables


```

1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)
4 #read(IPSUR)
5 library(prob)
6 #read(prob)
7 #install.packages("combinat")
8 library(combinat)
9 library(discreteRV)
10 require(pracma)
11 library(cubature)
12
13 #P{X1 = 1, X2 = 2, X3 = 0}
14 P = 0.20*0.10*0.30

```

R code Exa 4.3.f Conditional Probability Mass Function

```

1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)
4 #read(IPSUR)
5 library(prob)
6 #read(prob)
7 #install.packages("combinat")
8 library(combinat)
9 library(discreteRV)
10 require(pracma)
11 library(cubature)
12
13 #P{G=1} = 0.3875
14 C = 0.3875
15 #P{B=0|G=1}=(P{B=0,G=1})/(P{G=1})
16 P = 0.1/C
17 P
18 #P{B=1|G=1}

```

```

19 P = .175/C
20 P
21 #P{B=2|G=1}
22 P = .1125/C
23 P
24 #P{B=3|G=1}
25 P = 0
26 P

```

R code Exa 4.3.g Conditional Probability Mass Function

```

1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)
4 #read(IPSUR)
5 library(prob)
6 #read(prob)
7 #install.packages("combinat")
8 library(combinat)
9 library(discreteRV)
10 require(pracma)
11 library(cubature)
12
13 #P(0,0)
14 P_0_0 = .4
15 #P(0,1)
16 P_0_1 = .2
17 #P(1,0)
18 P_1_0 = .1
19 #P(1,1)
20 P_1_1 = .3
21
22 #P{Y = 1} = P(0,1) + #P(1,1)
23 P_Y = P_0_1 + P_1_1
24

```

```

25 #P{X = 0|Y=1} = P(0,1)/P{Y=1}
26 P1 = P_0_1 / P_Y
27 P1
28
29 #P{X = 1|Y=1} = P(1,1)/P{Y=1}
30 P2 = P_1_1 / P_Y
31 P2

```

R code Exa 4.4.a Expectation of a fair die

```

1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)
4 #read(IPSUR)
5 library(prob)
6 #read(prob)
7 #install.packages("combinat")
8 library(combinat)
9 library(discreteRV)
10 require(pracma)
11 library(cubature)
12
13
14 S <- rolldie(1, makespace = TRUE)
15 Ex = sum(S$X1*S$probs)
16 Ex

```

R code Exa 4.4.d Expectation of the message time

```

1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)
4 #read(IPSUR)

```

```

5 library(prob)
6 #read(prob)
7 #install.packages("combinat")
8 library(combinat)
9 library(discreteRV)
10 require(pracma)
11 library(cubature)
12
13
14 integrand <- function(x) {x / 1.5}
15 R = integrate(integrand, lower = 0, upper = 1.5)
16 R

```

R code Exa 4.5.a Expectation

```

1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)
4 #read(IPSUR)
5 library(prob)
6 #read(prob)
7 #install.packages("combinat")
8 library(combinat)
9 library(discreteRV)
10 require(pracma)
11 library(cubature)
12
13
14 # P {X}
15 PY0 = .2
16 PY1 = .5
17 PY2 = .3
18 # E[X^2] = E[Y] = Y0 * PY0 + Y1 * PY1 + Y2 * PY0
19 E = 0 * PY0 + 1 * PY1 + 4 * PY2
20 E

```

R code Exa 4.5.b Expected cost of breakdown

```
1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)
4 #read(IPSUR)
5 library(prob)
6 #read(prob)
7 #install.packages("combinat")
8 library(combinat)
9 library(discreteRV)
10 require(pracma)
11 library(cubature)
12
13
14 integrand <- function(x) {x*(1/3)*(x^(-2/3))}
15 R = integrate(integrand, lower = 0, upper = 1)
```

R code Exa 4.5.c Expectation

```
1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)
4 #read(IPSUR)
5 library(prob)
6 #read(prob)
7 #install.packages("combinat")
8 library(combinat)
9 library(discreteRV)
10 require(pracma)
11 library(cubature)
```

```

12
13
14 # P {X}
15 P0 = .2
16 P1 = .5
17 P2 = .3
18 # E[X^2] = X0^2*P0 + X1^2*P1 + X2^2*P2
19 E = (0^2)*P0 + (1^2)*P1 + (2^2)*P2
20 E

```

R code Exa 4.5.d Expectation

```

1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)
4 #read(IPSUR)
5 library(prob)
6 #read(prob)
7 #install.packages("combinat")
8 library(combinat)
9 library(discreteRV)
10 require(pracma)
11 library(cubature)
12
13
14 integrand <- function(x) {x^(3)}
15 R = integrate(integrand, lower = 0, upper = 1)

```

R code Exa 4.5.e Expected profit

```

1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)

```

```

4 #read(IPSUR)
5 library(prob)
6 #read(prob)
7 #install.packages("combinat")
8 library(combinat)
9 library(discreteRV)
10 require(pracma)
11 library(cubature)
12
13
14 profits <- c(10,20,40)
15 probs <- c(.2,.8,.3)
16 E1 = profits[1]*probs[1]+0*(1-probs[1])
17 E2 = profits[2]*probs[2]+0*(1-probs[2])
18 E3 = profits[3]*probs[3]+0*(1-probs[3])
19 E = E1 + E2 + E3
20 E

```

R code Exa 4.5.g Different types of coupons

```

1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)
4 #read(IPSUR)
5 library(prob)
6 #read(prob)
7 #install.packages("combinat")
8 library(combinat)
9 library(discreteRV)
10 require(pracma)
11 library(cubature)
12
13
14 EX = 20 * (1 - (19/20)^10)
15 EX

```

R code Exa 4.6.a Variance of a fair die

```
1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)
4 #read(IPSUR)
5 library(prob)
6 #read(prob)
7 #install.packages("combinat")
8 library(combinat)
9 library(discreteRV)
10 require(pracma)
11 library(cubature)
12
13
14 S <- rolldie(1, makespace = TRUE)
15 Ex = sum(S$X1*S$probs)
16 Ex2 = sum((S$X1)^2*S$probs)
17 Ex2
18 Varx = Ex2 - (Ex)^2
19 Varx
```

R code Exa 4.7.a Variance of 10 rolls of a fair die

```
1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)
4 #read(IPSUR)
5 library(prob)
6 #read(prob)
7 #install.packages("combinat")
```



```

8 library(combinat)
9 library(discreteRV)
10 require(pracma)
11 library(cubature)
12
13 S <- rolldie(1, makespace = TRUE)
14 Ex = sum(S$X1*S$probs)
15 Ex2 = sum((S$X1)^2*S$probs)
16 Ex2
17 Varx = Ex2 - (Ex)^2
18 Varx
19 Varx10 = 10 * Varx
20 Varx10

```

R code Exa 4.7.b Variance of 10 tosses of a coin

```

1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)
4 #read(IPSUR)
5 library(prob)
6 #read(prob)
7 #install.packages("combinat")
8 library(combinat)
9 library(discreteRV)
10 require(pracma)
11 library(cubature)
12
13 Var = .5 * (1 - .5)
14 Var
15
16 Var10 = 10 * Var
17 Var10

```

R code Exa 4.9.a Inequalities

```
1 #install.packages("IPSUR")
2 #install.packages("prob")
3 library(IPSUR)
4 #read(IPSUR)
5 library(prob)
6 #read(prob)
7 #install.packages("combinat")
8 library(combinat)
9 library(discreteRV)
10 require(pracma)
11 library(cubature)
12
13
14 # Markov
15 # P {X > 75}
16 meanX = 50
17 P = meanX / 75
18 P
19
20 #Chebyshev
21 # P {|X - 50| >= 10 } <= (SigmaX^2 / 10^2)
22 VarX = 25
23 P = 1 - (VarX / 10^2)
24 P
```

Chapter 5

Special Random Variables

R code Exa 5.1.a Returning of disks

```
1 #Probability that the package will have to be
  replaced
2 library(IPSUR)
3 P = 1 - choose(10,0)*(0.01)^0*(0.99)^10-choose(10,1)
  *(0.01)^1*(0.99)^9
4 P
5 #Probability that exactly one of the three packages
  will be returned
6 PP = choose(3,1)*P*(1-P)^2
7 PP
```

R code Exa 5.1.b Colour of Eyes

```
1 library(IPSUR)
2 #The probability that a child recieves the blue eyed
  gene from both parents
3 Pblue = (1/2)*(1/2)
4 Pblue
```

```

5 #the probability that exactly two of them have blue
  eye color
6 P = choose(4,2)*(1/4)^2*(3/4)^2
7 P

```

R code Exa 5.1.e Binomial Random Variable

```

1 library(IPSUR)
2 install.packages("binom")
3 library(binom)
4 pbinom(q = 0, size = 6, prob = 0.4)
5 pbinom(q = 1, size = 6, prob = 0.4) - pbinom(q = 0,
  size = 6, prob = 0.4)
6 pbinom(q = 2, size = 6, prob = 0.4) - pbinom(q = 1,
  size = 6, prob = 0.4)
7 pbinom(q = 3, size = 6, prob = 0.4) - pbinom(q = 2,
  size = 6, prob = 0.4)
8 pbinom(q = 4, size = 6, prob = 0.4) - pbinom(q = 3,
  size = 6, prob = 0.4)
9 pbinom(q = 5, size = 6, prob = 0.4) - pbinom(q = 4,
  size = 6, prob = 0.4)
10 pbinom(q = 6, size = 6, prob = 0.4) - pbinom(q = 5,
  size = 6, prob = 0.4)

```

R code Exa 5.1.f Binomial Distribution

```

1 pbinom(q = 70, size = 100, prob = 0.75) - pbinom(q =
  69, size = 100, prob = 0.75)
2 pbinom(q = 70, size = 100, prob = 0.75)

```

R code Exa 5.2.a Probability of accident

```
1 #P{X>=1} = 1 - P{X = 0}
2 1-ppois(0,3)
3 #Or
4 1-dpois(0,3)
```

R code Exa 5.2.b Defective Items

```
1 ppois(1,1)
```

R code Exa 5.2.c Number of Alpha particles

```
1 ppois(2,3.2)
```

R code Exa 5.2.d Claims handled by an insurance company

```
1 #Proportion of days have less than 3 claims
2 ppois(2,5)
3
4 dpois(4,lambda = 5)
5
6 #the probability that 3 of the next 5 days will have
   claims is
7 library(IPSUR)
8 choose(5,3) *(dpois(4,lambda = 5))^3 *(1-dpois(4,
   lambda = 5))^2
```

R code Exa 5.2.f Defective stereos

```
1 i<-seq(0, 3, 1)
2 sum(dpois(i,8))
```

R code Exa 5.3.a Functional system

```
1 dhyper(4, m = 15, n = 5, k = 6)+dhyper(5, m = 15, n
  = 5, k = 6)+dhyper(6, m = 15, n = 5, k = 6)
```

R code Exa 5.3.b Determining Population Size

```
1 hyper <- function(N, M, n, i) {
2   factorial(N)*factorial(M)*factorial(n)*factorial(N
   +M-n)/(factorial(i)*factorial(N-i)*factorial(n-
   i)*factorial(M-n+i)*factorial(N+M))
3 }
4
5 r= 50
6 n=100
7 X=25
8 cat("Estimate of the number of animals in the region
   is",r*n/X)
```

R code Exa 5.3.c Conditional Probability

```
1 bino <- function(n, k, p) {
2   factorial(n)*(p^k)*((1-p)^(n-k))/(factorial(k)*
   factorial(n-k))
3 }
```

```

4
5 condprob <- function(n,k,p,i) {
6   bino(n,i,p)*bino(m,k-i,p)/bino(n+m,k, p);
7 }
8
9 #The function condprob will give  $P\{X=i | X+Y=k\}$ 

```

R code Exa 5.4.a Probabilities of a uniform random variable

```

1 beta = 10
2 alpha = 0
3 #a
4 a = 2
5 b = 9
6 (b-a)/(beta-alpha)
7 #b
8 a = 1
9 b = 4
10 (b-a)/(beta-alpha)
11 #c
12 a = 0
13 b = 5
14 (b-a)/(beta-alpha)
15 #d
16 a = 6
17 b = 10
18 (b-a)/(beta-alpha)

```

R code Exa 5.4.b Bus Timings

```

1 beta = 30
2 alpha = 0
3 #a

```

```

4 a1 = 10
5 b1 = 15
6 a2 = 25
7 b2 = 30
8 (b1-a1)/(beta-alpha)+(b2-a2)/(beta-alpha)
9
10 #b
11 a1 = 0
12 b1 = 3
13 a2 = 15
14 b2 = 18
15 (b1-a1)/(beta-alpha)+(b2-a2)/(beta-alpha)

```

R code Exa 5.4.c Current in a diode

```

1 I0 = 10^-6
2 a = 5
3 integrand <- function(x) {0.5*exp(a*x)}
4 R = integrate(integrand, lower = 1, upper = 3)
5 (I0*R$value)-I0

```

R code Exa 5.5.a Normal Random Variable

```

1 #a
2 pnorm(11,3,sqrt(16))
3 #b
4 1-pnorm(-1,3,sqrt(16))
5 #c
6 pnorm(7,3,sqrt(16))-pnorm(2,3,sqrt(16))

```

R code Exa 5.5.b Noise in Binary Message

```
1 #a
2 pnorm(-1.5)
3 #b
4 1-pnorm(2.5)
```

R code Exa 5.5.c Power dissipation

```
1 mu = 6
2 sigma = 1
3 #a
4 E = 3*(mu^2+sigma^2) #ref page 169
5 #b
6 1-pnorm(sqrt(40),mu,sigma)
```

R code Exa 5.5.d Yearly precipitation

```
1 #a
2 1-pnorm(25,24.16,sqrt(19.22))
3 #b
4 1-pnorm(3,0,sqrt(19.22))
```

R code Exa 5.6.a Wearing of Battery

```
1 1-pexp(5000, rate = 10000)
```

R code Exa 5.6.b Working Machines

```
1 #When C is put to use , one other machine(either A or
   B ) will still be working .
2 #The probability of this machine or C failing is
   equal due to
3 #the memoryless propoerty of exponential random
   variables .
4
5 cat("The probability that machine which is still
   operable is machine C is ",1/2)
```

R code Exa 5.6.c Series System

```
1 new <- function(lamda,n, t ) {
2   newsum = 0;
3   for (i in 1:n){
4     newsum= newsum + lamda[i]
5     result=(exp(1))^( -1*newsum*t)
6   }
7 }
```

R code Exa 5.8.a Chi square random variable

```
1 pchisq(30,df =26)
```

R code Exa 5.8.b Chi square random variable

```
1 qchisq((1-0.05) ,df =15)
```

R code Exa 5.8.c Locating a Target

```
1 1-pchisq(9/4,df =3)
```

R code Exa 5.8.d Locating a Target in 2D space

```
1 1-pchisq(9/4,df =2)  
2 #or  
3 1-pexp(9/4,1/2)
```

R code Exa 5.8.e T distribution

```
1 #a  
2 pt(1.4,12)  
3 #b  
4 -qt(0.025,9)
```

R code Exa 5.8.f F Distribution

```
1 pf(1.5,6,14)
```

Chapter 6

Distribution of Sampling Statistics

R code Exa 6.3.a Claims handled by an insurance company

```
1 1-pnorm(8300000,320*25000,540*sqrt(25000))
```

R code Exa 6.3.c Class strength

```
1 1-pnorm((150.5-(450*0.3))/sqrt(450*.3*(1-.3)))
```

R code Exa 6.3.d Weights of workers

```
1 #a
2 2*pnorm(.8889)-1
3 #b
4 2*pnorm(1.7778)-1
```

R code Exa 6.3.e Distance of a start

```
1 qnorm(((1+.95)/2),0,1)
2 n = (1.96*4)^2
3 n
```

R code Exa 6.5.a Processing time

```
1 n = 15
2 sigmasq = 9
3 Ssq = 12
4 1-pchisq(((n-1)*Ssq/sigmasq),n-1)
```

R code Exa 6.6.a Candidate winning an election

```
1 favour = 0.45;
2 samplesize = 200;
3 expec= favour*samplesize;
4 sd = sqrt(samplesize*favour*(1-favour));
5 cat("The expected value is",expec)
6 cat("The standard deviation is",sd)
7
8 #b
9 prob = 1-pbinom(q = samplesize/2, size = samplesize,
  prob = favour)
10 cat("Probability that more than half the members of
  the sample favour the candidate",prob)
```

R code Exa 6.6.b Pork consumption

```
1 meaneach = 147;
2 sdeach   = 62;
3 samplesize = 25;
4 lim =150;
5 samplemean = meaneach;
6 samplesd= sdeach/sqrt(samplesize)
7 prob = 1 - pnorm(q = lim,mean = samplemean,sd =
    samplesd)
8 prob
```

Chapter 7

Parameter Estimation

R code Exa 7.2.a Maximum likelihood estimator of a bernoulli parameter

```
1 samplesize = 1000;
2 acceptable = 921;
3 cat("The maximum likelihood estimate of p is",
    acceptable/samplesize)
```

R code Exa 7.2.b Errors in a manuscript

```
1 totalerror<-function(n1, n2, n12){
2   totalerror = n1*n2/n12
3   return(totalerror)
4 }
5 totalerror(n1 = 33000, n2 = 33000, n12 = 17000)
```

R code Exa 7.2.c Maximum likelihood estimator of a poisson parameter

```
1 total_people = 857;
```

```

2 days= 20;
3 cat("The maximum likelihood estimate of lambda",
    total_people/days)

```

R code Exa 7.2.d Number of traffic accidents

```

1 accidents<-c(4,0,6,5,2,1,2,0,4,3)
2 lambda= mean(accidents)
3 cat("lambda:",lambda)
4 prob = ppois(q = 2, lambda = lambda)
5 cat("Proportion of non-rainy days that had 2 or
    fewer accidents that year:",prob)

```

R code Exa 7.2.e Maximum likelihood estimator in a normal population

```

1 normal <-function(X, Xmean, n){
2   u= Xmean;
3   newsum = 0;
4   for (i in 1:n){
5     newsum= newsum + (X(i)-Xmean)^2
6   }
7   sigmasquared = sqrt((newsum/n));
8   return(u,sigmasquared)
9 }

```

R code Exa 7.2.f Kolmogorovs law of fragmentation

```

1 X<-c(2.2,3.4,1.6,0.8,2.7,3.3,1.6,2.8,2.5,1.9)
2 upperlimX = 3
3 lowerlimX = 2

```



```

4 upperlimlogX= log(upperlimX)
5 lowerlimlogX = log(lowerlimX)
6
7 logX = log(X)
8 samplemean= mean(logX)
9 samplesd= sqrt(var(logX))
10 samplemean
11 samplesd
12 prob = pnorm(upperlimlogX, samplemean, samplesd) -
        pnorm(lowerlimlogX, samplemean, samplesd)
13 prob

```

R code Exa 7.2.g Estimating Mean of a Uniform Distribution

```

1 unif <- function(X, n) {
2   max(X)/2
3 }

```

R code Exa 7.3.a Error in a signal

```

1 avg = 0;
2 var = 4;
3 num = 9;
4 X<-c(5,8.5,12,15,7,9,7.5,6.5,10.5)
5 samplemean= mean(X);
6 lowerlim = samplemean - (1.96*sqrt(var/num))
7 upperlim = samplemean + (1.96*sqrt(var/num))
8 cat("The 95% confidence interval is ",lowerlim,"to "
    ,upperlim)

```

R code Exa 7.3.b Confidence interval

```
1 avg = 0;
2 var = 4;
3 num = 9;
4 X<-c(5,8.5,12,15,7,9,7.5,6.5,10.5)
5 samplemean= mean(X);
6 lowerlim = samplemean - (1.645*sqrt(var/num))
7 upperlim = samplemean + (1.645*sqrt(var/num))
8
9 cat("The 95% upper confidence interval is ",lowerlim
    ," to infinity")
10 cat("The 95% upper confidence interval is minus
    infinity to ",upperlim)
```

R code Exa 7.3.c Confidence interval

```
1 var = 4;
2 num = 9;
3 X<-c(5,8.5,12,15,7,9,7.5,6.5,10.5)
4 samplemean= mean(X);
5 samplesd = sd(X)
6 alpha= 0.005;
7 zalpha = qnorm(p = alpha,mean = 0,sd = 1,lower.tail
    = FALSE)
8 zalpha
9 lowerlim = samplemean - (zalpha*sqrt(var/num))
10 upperlim = samplemean + (zalpha*sqrt(var/num))
11 cat("The 95% confidence interval is ",lowerlim, "to
    ",upperlim)
12
13 alpha= 0.01;
14 zalpha = qnorm(p = alpha,mean = 0,sd = 1,lower.tail
    = FALSE)
15 lowerlim = samplemean - (zalpha*sqrt(var/num))
```

```

16 upperlim = samplemean + (zalpha*sqrt(var/num))
17 cat("The 95% upper confidence interval is ",lowerlim
    , " to infinity")
18 cat("The 95% upper confidence interval is minus
    infinity to ",upperlim)

```

R code Exa 7.3.d Weight of a salmon

```

1 sd= 0.3;
2 lim = 0.1;
3 num = (1.96*sd/lim)^2;
4 cat("Sample size should be greater than",round(num))

```

R code Exa 7.3.e Error in a signal

```

1 X<-c(5, 8.5, 12, 15, 7, 9, 7.5, 6.5, 10.5)
2 num = 9;
3 meanX= mean(X);
4 X2 = X^2;
5 s2= (sum(X2)- (num*(meanX^2)))/(num-1);
6 s= sqrt(s2);
7 tval = qt(.975,8)
8 upperlim = meanX + (tval*s)/sqrt(num);
9 lowerlim = meanX - (tval*s)/sqrt(num);
10 cat("The 95% confidence interval is ", lowerlim, "
    to ",upperlim )
11
12 #7.3 a
13 s2= 4
14 s= sqrt(s2);
15 tval = qt(.975,8)
16 upperlim = meanX + (tval*s)/sqrt(num);
17 lowerlim = meanX - (tval*s)/sqrt(num);

```

```
18 cat("The 95% confidence interval is ", lowerlim, "
    to ",upperlim )
```

R code Exa 7.3.f Average resting pulse

```
1 X<-c(54, 63, 58, 72, 49, 92, 70, 73, 69, 104, 48,
      66, 80, 64, 77)
2 num = 15;
3 meanX= mean(X);
4 X2 = X^2;
5 s2= (sum(X2)- (num*(meanX^2)))/(num-1);
6 s= sqrt(s2);
7 tval = qt(.975,num-1)
8 upperlim = meanX + (tval*s)/sqrt(num);
9 lowerlim = meanX - (tval*s)/sqrt(num);
10 cat("The 95% confidence interval is ",lowerlim,"to "
     ,upperlim)
11 alpha = 0.05;
12 tval = qt(1-alpha,num-1)
13 lim = meanX + (tval*s)/sqrt(num);
14 cat("The 95% lower confidence interval is from minus
     infinity to ",lim)
```

R code Exa 7.3.g Evaluating integrals

```
1 #Method 1: Monte Carlo Integration
2 #install.packages("functional")
3 library(functional)
4 funct<-function(sampleXVals) {
5   oned = sqrt(1-(sampleXVals*sampleXVals))
6   return(oned)
7 }
8 MCIntegrate <- function(funct, a,b,N) {
```

```

9   sampleXVals = matrix(0,N)
10  set.seed(148)
11  sampleXVals = a + (b-a)*runif(N)
12  Integral = (b-a)* mean(funcnt(sampleXVals))
13  return(Integral)
14 }
15 MCIntegrate(funcnt,0,1,50000000)
16
17
18 #Method 2
19 meanX = 0.786;
20 s= 0.03;
21 num = 100;
22 alpha = 0.05;
23 tval = qt(1-alpha,num-1)
24 upperlim = meanX + (tval*s)/sqrt(num);
25 lowerlim = meanX - (tval*s)/sqrt(num);
26 cat("The 95% confidence interval is ",lowerlim,"to "
    ,upperlim)

```

R code Exa 7.3.h Thickness of washers

```

1  num=10;
2  X<-c(0.123, 0.133, 0.124, 0.126, 0.120, 0.130,
      0.125, 0.128, 0.124, 0.126)
3  var(X)
4  s2 = var(X)
5  chi1 = qchisq(1-.95,num-1)
6  chi2 = qchisq(.95,num-1)
7  lowerlim = (num-1)*s2/chi2;
8  lowerlim
9  upperlim = (num-1)*s2/chi1;
10 upperlim
11 cat("The 90% confidence interval is ",sqrt(lowerlim)
    , "to ",sqrt(upperlim))

```

R code Exa 7.4.a Cable insulation

```
1 A<-c(36, 44, 41, 53, 38, 36, 34, 54, 52, 37, 51, 44,
      35, 44)
2 B<-c(52, 64, 38, 68, 66, 52, 60, 44, 48, 46, 70, 62)
3 sigmaA= 40;
4 sigmaB= 100;
5 alpha = 1-0.95;
6 beta= alpha/2;
7 meanA = mean(A);
8 meanB= mean(B);
9 zbeta = qnorm(1-beta)
10
11 lowerlim= mean(A) - mean(B) - (zbeta*sqrt((sigmaA/
      length(A)) + (sigmaB/length(B)))) ;
12 upperlim= mean(A) - mean(B) + (zbeta*sqrt((sigmaA/
      length(A)) + (sigmaB/length(B)))) ;
13 cat("The 95% confidence interval is ",lowerlim, "to
      ",upperlim)
14
15 beta=alpha;
16 zbeta = qnorm(1-beta)
17
18 upperlim= mean(A) - mean(B) + (zbeta*sqrt((sigmaA/
      length(A)) + (sigmaB/length(B)))) ;
19 cat("A value that exceed the difference of the means
      with 95% confidence is",upperlim)
```

R code Exa 7.4.b Battery production

```
1 tech1<-c(140, 136, 138, 150, 152, 144, 132, 142,
          150, 154, 136, 142)
```

```

2 tech2<-c(144, 132, 136, 140, 128, 150, 130, 134,
           130, 146, 128, 131, 137, 135)
3 num1= 12;
4 num2= 14;
5 mean1= mean(tech1);
6 mean2= mean(tech2);
7 alpha = 0.9;
8 S1 = var(tech1)
9 S2 = var(tech2)
10 Sp = (((num1-1)*S1) + ((num2-1)*S2))/(num1+ num2 -2)
      ;
11 Sp= sqrt(Sp);
12 num= (1/num1)+(1/num2);
13 betaa = (1-alpha)/2;
14 tval = qt(1-betaa,num1+num2-2)
15 upperlim = mean1-mean2 + (tval*Sp)*sqrt(num);
16 lowerlim = mean1-mean2 - (tval*Sp)*sqrt(num);
17 cat("The 90% confidence interval is ",lowerlim, "to
      ", upperlim)
18 alpha = 0.95
19 betaaa = 1-alpha;
20 tval = qt(1-betaaa,num1+num2-2)
21 lowerlim = mean1-mean2 - (tval*Sp)*sqrt(num);
22 cat("the upper confidence interval is",lowerlim," to
      infinity")

```

R code Exa 7.5.a Transistors

```

1 phat = 0.8;
2 zalpha = 1.96;
3
4 samplesize = 100;
5 lowerlim = phat - (zalpha*sqrt(phat*(1-phat)/
      samplesize));
6 upperlim = phat + (zalpha*sqrt(phat*(1-phat)/

```

```
    samplesize));  
7 cat("The 95% confidence interval is ",lowerlim, "to  
    ",upperlim)
```

R code Exa 7.5.b Survey

```
1 phat = 0.52;  
2 error = 0.04;  
3 zalpha = 1.96;  
4 samplesize = (error/zalpha)^2/(phat*(1-phat));  
5 1/samplesize
```

R code Exa 7.5.c Acceptable chips

```
1 initialsample = 30;  
2 acceptable= 26;  
3 phat = acceptable/initialsample;  
4 error = 0.05/2;  
5 zalpha = 2.58;  
6  
7 samplesize = (error/zalpha)^2/(phat*(1-phat));  
8 finalsize = round(1/samplesize);  
9 acceptablenew= 1040 + acceptable;  
10 phat = acceptablenew/finalsize;  
11 lowerlim = phat - (zalpha*sqrt(phat*(1-phat)/  
    finalsize));  
12 upperlim = phat + (zalpha*sqrt(phat*(1-phat)/  
    finalsize));  
13 cat("The 99% confidence interval is ",lowerlim, "to  
    ",upperlim)
```

R code Exa 7.6.a Life of a product

```
1 sum_lives = 1740;
2 num = 10;
3 alpha = (1-0.95)/2;
4 chi1= qchisq(1-alpha,2*num)
5 chi2 = qchisq(alpha,2*num)
6 lowerlim = 2*sum_lives/chi1;
7 upperlim = 2*sum_lives/chi2;
8 cat("The 95% confidence interval is ",lowerlim, "to
    ",upperlim)
9
10 '
11 Textbook confidence interval is from 101.847 to
    360.211
12 whereas in above solution is 101.84489 to 362.8485
    because of the difference in the value of chi-
    square(0.975, 20).
13 The textbook says the value is 9.661 whereas R
    calculates its value as 9.59
14 '
```

R code Exa 7.7.a Point estimator

```
1 estimator1 <- function(x) {
2   X[1]
3 }
4 estimator2 <- function(variables) {
5   mean(X)
6 }
```

R code Exa 7.7.b Point estimator

```

1 estimate <- function(d, sigma) {
2   sigmainv = 1/sigma
3   new = d/sigma
4   result1 = sum(new)/sum(sigmainv)
5 }
6
7 merror <- function(sigma) {
8   sigmainv = 1/sigma
9   result1 = 1/sum(sigmainv)
10 }

```

R code Exa 7.7.c Point estimator of a uniform distribution

```

1 unbiasedestimator <- function(X, n) {
2   c=(n+2)/(n+1)
3   result = c*max(X)
4 }

```

R code Exa 7.8.a Bayes estimator

```

1 estimator <- function(X, n) {
2   (sum(X) +1)/(n+2)
3 }

```

R code Exa 7.8.b Bayes estimator of a normal population

```

1 meanestimator <- function(sigma0 , u, sigma, n, X) {
2   meanX= mean(X)
3   result = (n*meanX/sigma0)/((n/sigma0)+(1/sigma)) +
             (u/sigma)/((n/sigma0)+(1/sigma))

```

```

4 }
5
6 varestimator <- function(sigma0 , sigma, n) {
7   (sigma0*sigma)/((n*sigma)+sigma0)
8 }

```

R code Exa 7.8.d estimator of the signal value

```

1 meanestimator <- function(sigma0 , u, sigma, n, X) {
2   meanX= mean(X);
3   result = (n*meanX/sigma0)/((n/sigma0)+(1/sigma)) +
4             (u/sigma)/((n/sigma0)+(1/sigma));
5   return(result)
6 }
7
8 varestimator <- function(sigma0 , sigma, n) {
9   result = (sigma0*sigma)/((n*sigma)+sigma0);
10  return(result)
11 }
12
13 u = 50;
14 sigma= 100;
15 sigma0 = 60;
16 n =1;
17 X =40;
18 expec = meanestimator(sigma0 , u, sigma, n, X);
19 var = varestimator (sigma0, sigma,n);
20
21 zalpha = 1.645
22 lowerlim = -1*sqrt(var)*zalpha+expec;
23 upperlim = sqrt(var)*zalpha+expec;
24 cat("With probability 0.9, the sent signal lies
25     between ",lowerlim, "to ", upperlim )

```

Chapter 8

Hypothesis Testing

R code Exa 8.3.a Noise in a Signal

```
1 noise_var = 4;
2 noise_mean = 0;
3 num = 5;
4 Xbar = 9.5;
5 u = 8;
6 statistic = sqrt(num/noise_var)*(Xbar - u);
7 compare = qnorm(0.975,0, 1);
8 if(statistic<compare){
9   cat("Hypothesis is accepted");
10 } else{
11   cat("Hypothesis is not accepted")
12 }
```

R code Exa 8.3.b Error in a signal

```
1 noise_var = 4;
2 noise_mean = 0;
3 num = 5;
```

```

4 Xbar = 8.5;
5 u = 8;
6 statistic = sqrt(num/noise_var)*(Xbar - u);
7
8 prob = 2*pnorm(-1*statistic , 0,1 );
9 cat("P-value is",prob)
10
11 Xbar = 11.5;
12 statistic = sqrt(num/noise_var)*(Xbar - u);
13 prob = pnorm(-1*statistic , 0,1 );
14 cat("P-value is",prob)

```

R code Exa 8.3.c Error in a signal

```

1 noise_var = 4;
2 num = 5;
3 Xbar = 10;
4 u = 8;
5 statistic = sqrt(num/noise_var)*(Xbar - u);
6 compare = qnorm(0.975,0, 1);
7 lim1 = statistic + compare;
8 lim2 = statistic - compare;
9 prob = pnorm(lim1 , 0,1 ) - pnorm(lim2 , 0,1 );
10 prob

```

R code Exa 8.3.d Number of signals to be sent

```

1 alpha = 0.025;
2 betaa = 0.25;
3 u1 = 9.2;
4 uo = 8;
5 var =4;
6 zalpha = qnorm(1-alpha,0, 1);

```

```

7 zbeta = qnorm(1-betaa,0, 1);
8 n = ((zalpha + zbeta)/(u1-u0))^2 *var;
9 cat("Required number of samples is",ceiling(n))
10 statistic = sqrt(ceiling(n)/var)*(u1 - u0);
11 lim1 = -1*statistic + zalpha;
12 lim2 = -1*statistic - zalpha;
13 prob = pnorm(lim1 , 0,1 )- pnorm(lim2 , 0,1 );
14 cat("Thus, if the message is sent the reqd number of
      times is , then the probability that the null
      hypothesis will be rejected is",1-prob)

```

R code Exa 8.3.e Number of signals to be sent

```

1 n =5;
2 Xbar = 9.5;
3 u = 8;
4 var = 4;
5 statistic = sqrt(n/var)*(Xbar - u);
6 p = 1 - pnorm(statistic, 0, 1);
7 cat("The test would call for rejection at all
      significance levels greater than or equal to ",p)

```

R code Exa 8.3.f Nicotine content in a cigarette

```

1 n =20;
2 Xbar = 1.54;
3 uo = 1.6;
4 sd = 0.8;
5 statistic = sqrt(n)*(Xbar - uo)/sd;
6 cat("Test statistic is",statistic)
7 p = pnorm(statistic, 0, 1);
8 cat("P-value is",p)

```

R code Exa 8.3.g Blood cholesterol level

```
1 n = 50;
2 Xbar = 14.8;
3 S = 6.4;
4 T = sqrt(n)*Xbar/S;
5 cat("The T value is",T)
```

R code Exa 8.3.h Water usage

```
1 X<-c(340, 356, 332, 362, 318, 344, 386, 402, 322,
      360, 362, 354, 340, 372, 338, 375, 364, 355, 324,
      370)
2 uo = 350;
3 Xbar = mean(X);
4 var = var(X);
5 S = sqrt(var)
6 n = length(X)
7 T = sqrt(n)*(Xbar - uo)/S;
8 Tvalue = qt(0.95,n-1)
9 cat("The T value is ",T)
10 if(T<Tvalue){
11   cat("Null hypothesis is accepted at 10% level of
      significance")
12 }else{
13   cat("Null hypothesis is not accepted at 10% level
      of significance")
14 }
15 cat("P value of the test data",2*pnorm(-1*T,0,1))
```

R code Exa 8.3.i Life of a tire

```
1 X<-c(36.1, 40.2, 33.8, 38.5, 42, 35.8, 37, 41, 36.8,
      37.2, 33, 36)
2 n = length(X);
3 uo = 40;
4 Xbar=mean(X);
5 sd = sqrt(var(X));
6 T = sqrt(n)*(Xbar - uo)/sd;
7 Tvalue = qt(0.05,n-1);
8 cat("The T value is ",T)
9 if(T<Tvalue){
10   cat("Null hypothesis is rejected at 5% level of
      significance")
11 }else{
12   cat("Null hypothesis is accepted at 5% level of
      significance")
13 }
14
15 cat("P value of the test data",pnorm(T,0,1))
16 '
17 There is a mismatch between book answer for the P
   value of the test data.
18 '
```

R code Exa 8.3.j Service Time

```
1 X<-c(8.6, 9.4, 5.0, 4.4, 3.7, 11.4, 10.0, 7.6, 14.4,
      12.2, 11.0, 14.4, 9.3, 10.5, 10.3, 7.7, 8.3,
      6.4, 9.2, 5.7, 7.9, 9.4, 9.0, 13.3, 11.6, 10.0,
      9.5, 6.6)
2 n = length(X);
3 uo = 8;
4 Xbar=mean(X);
5 sd = sqrt(var(X));
```



```

6 T = sqrt(n)*(Xbar - uo)/sd;
7 cat("The test statistic is ",T)
8 p = 1- pt(T, n-1);
9 cat("P-value is",p)
10 cat("A small p value indicates that the mean service
      time exceeds 8 minutes")

```

R code Exa 8.4.a Tire lives

```

1 A<-c(61.1, 58.2, 62.3, 64, 59.7, 66.2, 57.8, 61.4,
      62.2, 63.6)
2 B<-c(62.2, 56.6, 66.4, 56.2, 57.4, 58.4, 57.6, 65.4)
3 uA = mean(A);
4 uB = mean(B);
5 varA = 40^2;
6 varB =60^2;
7 n= length(A);
8 m =length(B);
9 den = sqrt((varA/n)+ (varB/m));
10 statistic = (uA -uB)/den;
11 cat("The test statistic is",statistic);
12 cat("A small value of the test statistic indicates
      that the null hypothesis is accepted")

```

R code Exa 8.4.b Medicine for cold

```

1 X<-c(5.5, 6.0, 7.0, 6.0, 7.5, 6.0, 7.5, 5.5, 7.0,
      6.5)
2 Y<-c(6.5, 6.0, 8.5, 7.0, 6.5, 8.0, 7.5, 6.5, 7.5,
      6.0, 8.5, 7.0)
3 n = length(X);
4 m= length(Y);
5 Xbar= mean(X);

```

```

6 Ybar = mean(Y);
7 Sx = var(X);
8 Sy = var(Y);
9 Sp = ((n-1)*Sx/(n+m-2)) + ((m-1)*Sy/(n+m-2));
10 den = sqrt(Sp*((1/n)+(1/m)));
11 TS = (Xbar - Ybar)/den;
12 cat("The test statistic is",TS);
13 tvalue = qt( 0.95, m+n-2)
14 if(TS<tvalue){
15     cat("Null hypothesis is rejected at 5% level of
        significance")
16 }else{
17     cat("Null hypothesis is accepted at 5% level of
        significance")
18 }
19 cat("The Pvalue is",pt(TS,m+n-2))

```

R code Exa 8.4.c Unknown population variance

```

1 A<-c(61.1, 58.2, 62.3, 64, 59.7, 66.2, 57.8, 61.4,
      62.2, 63.6)
2 B<-c(62.2, 56.6, 66.4, 56.2, 57.4, 58.4, 57.6, 65.4)
3 uA = mean(A);
4 uB = mean(B);
5 n= length(A);
6 m =length(B);
7 Sx = var(A);
8 Sy = var(B);
9 Sp = ((n-1)*Sx/(n+m-2)) + ((m-1)*Sy/(n+m-2));
10 den = sqrt(Sp*((1/n)+(1/m)));
11 TS = (uA-uB)/den;
12 cat("The test statistic is",TS);
13 pvalue = 2*(1- pt(TS,m+n-2));
14 cat("Null hypothesis is accepted at any significance
      level less than",pvalue)

```

R code Exa 8.4.d effectiveness of safety program

```
1 A<-c(30.5, 18.5, 24.5, 32, 16, 15, 23.5, 25.5, 28,
      18)
2 B<-c(23, 21, 22, 28.5, 14.5, 15.5, 24.5, 21, 23.5,
      16.5)
3 n= length(A);
4 W = B-A;
5 Wbar = mean(W);
6 S = sqrt(var(W));
7 T = sqrt(n)*Wbar/S;
8 cat("The test statistic is",T);
9 pvalue = pt(T, n-1);
10 cat("The p value is",pvalue)
```

R code Exa 8.5.a effectiveness of machine

```
1 n =20;
2 S2= 0.025;
3 chk = 0.15;
4 compare = (n-1)*S2/(chk^2);
5 pvalue = 1- pchisq(compare, n-1);
6 cat("The p-value is",pvalue)
7 cat("Thus , the null hypothesis is accepted")
```

R code Exa 8.5.b Catalyst

```
1 S1 = 0.14;
2 S2 = 0.28;
```

```

3 n= 10;
4 m= 12;
5 ratio = S1/S2;
6 prob1 = pf(ratio, n-1, m-1);
7 prob2 = 1-prob1;
8 prob = min(prob1,prob2);
9 pvalue = 2*prob;
10 cat("The p value is",pvalue)
11 cat("So the hypothesis of equal variance cannot be
    rejected")

```

R code Exa 8.6.a Computer chip manufacturing

```

1 samplesize = 300;
2 p =0.02;
3 defective=9;
4 val = pbinom(defective, samplesize, p, 1-p);
5 cat("P0.02{X>10} = ",val);
6 cat("Manufacturers claim cannot be rejected at the
    5% level of significance")

```

R code Exa 8.6.b Finding p value

```

1 samplesize = 300;
2 p =0.02;
3 defective=9;
4 compare = 10;
5 npo = samplesize*p;
6 sd = sqrt(npo*(1-p));
7 tol = 0.5;
8 pvalue = 1- pnorm(compare-tol, npo,sd );
9 cat("The pvalue is",pvalue)

```

R code Exa 8.6.c Change in manufacturing pattern

```
1 samplesize = 500;
2 p =0.04;
3 defective=16;
4 prob1 = 1-pbinom(defective, samplesize, p, 1-p)
5 prob2 = pbinom(defective-1, samplesize, p, 1-p);
6 pvalue = 2*min(prob1,prob2);
7 cat("The pvalue is",pvalue)
```

R code Exa 8.7.a Mean number of defective chips

```
1 x<-c(28, 34, 32, 38, 22)
2 claim = 25;
3 total = sum(x);
4 pval = 1 - ppois(total-1, (claim*length(x)));
5 cat("The pvalue is",pval)
```

R code Exa 8.7.b Safety Conditions in a plant

```
1 plant1<-c(16, 18, 9, 22, 17, 19, 24, 8)
2 plant2<-c(22, 18, 26, 30, 25, 28)
3 X1= sum(plant1);
4 X2 = sum(plant2);
5 n =length(X1);
6 m= length(X2);
7 prob1 = 1 - pbinom(X1, X1+X2,(4/7),(3/7));
8 prob2 = pbinom(X1-1 ,X1+X2, 4/7, 3/7 );
9 cat(prob1, prob2)
```

```
10 pvalue = 2*min(prob1,prob2);
11 cat("The pvalue is",pvalue)
```

R code Exa 8.7.c Better proof reader

```
1 Aerror =28;
2 Berror = 18;
3 common =10;
4 N2 = Aerror - common;
5 N3 =Berror- common;
6 pval = pbinom(N2-1, N2 + N3, 0.5, 0.5);
7 cat("P-value is",pval)
```

Chapter 9

Regression

R code Exa 9.1.a Scatter Diagram

```
1 X<-c(100, 110, 120, 130, 140, 150, 160, 170, 180,
      190);
2 Y<-c(45, 52, 54, 63, 62, 68, 75, 76, 92, 88)
3 dev.new(width=5, height=4)
4 plot(X, Y)
5 cat("A linear regression model seems appropriate")
```

R code Exa 9.2.a Relative humidity and moisture content

```
1 A<-c(46, 53, 29, 61, 36, 39, 47, 49, 52, 38, 55, 32,
      57, 54, 44)
2 B<-c(12, 15, 7, 17, 10, 11, 11, 12, 14, 9, 16, 8,
      18, 14, 12)
3 plot(A, B,xlab ="Relative humidity",ylab = "Moisture
      content",main = "Estimated regression line")
4 X = lm(B~A)
5 Y = X$coefficients[1]
6 X = X$coefficients[2]
```

```

7 p = seq(0,65,0.1)
8 q = p*X + Y
9 lines(p, q)

```

R code Exa 9.3.a Moisture against Density

```

1 x<-c(5, 6, 7, 10, 12, 15, 18, 20)
2 y<-c(7.4, 9.3, 10.6, 15.4, 18.1, 22.2, 24.1, 24.8)
3 plot(x,y);
4
5 xbar = mean(x);
6 ybar= mean(y);
7 n= 8;
8 SxY = 0;
9 for (i in (1:n)){
10   SxY = SxY + (x[i]*y[i]) - (xbar*ybar)
11 }
12
13 Sxx = 0;
14 for (i in 1:n){
15   Sxx= Sxx + (x[i]*x[i]) - (xbar*xbar);
16 }
17
18 SYy = 0;
19 for (i in 1:n){
20   SYy = SYy + (y[i]*y[i]) - (ybar*ybar);
21 }
22
23 B = SxY/Sxx;
24 A = ybar - (B*xbar);
25 cat("A is",A);
26 cat("B is",B);
27 p= seq(0,20,0.1)
28 q= A + B*p;
29 lines(p,q)

```



```

30
31 SSR = ((Sxx*SYy)-(SxY*SxY))/Sxx ;
32 cat("The SSR is",SSR)

```

R code Exa 9.4.a Effect of speed on mileage

```

1 x<-c(45, 50, 55, 60, 65, 70, 75)
2 y<-c(24.2, 25.0, 23.3, 22.0, 21.5, 20.6, 19.8)
3 xbar = mean(x);
4 ybar= mean(y);
5 n= 7;
6 SxY = 0;
7 for (i in 1:n){
8   SxY = SxY + (x[i]*y[i]) - (xbar*ybar)
9 }
10
11 Sxx = 0;
12 for (i in 1:n){
13   Sxx= Sxx + (x[i]*x[i]) - (xbar*xbar);
14 }
15 SYy = 0;
16 for (i in 1:n){
17   SYy = SYy + (y[i]*y[i]) - (ybar*ybar);
18 }
19 B = SxY/Sxx;
20 A = ybar - (B*xbar);
21 cat("A is",A);
22 cat("B is",B);
23
24 SSR = ((Sxx*SYy)-(SxY*SxY))/Sxx ;
25 cat("The SSR is",SSR)
26
27 ts = sqrt(((n-2)*Sxx)/SSR)*abs(B);
28 cat("the test statistic is",ts);
29 tvalue= qt(0.995,5);

```

```

30 cat("tvalue is",tvalue);
31
32 if(tvalue < ts){
33     cat("Hypothesis beta= 0 is rejected at 1% level of
        significance")
34 } else{
35     cat("Hypothesis beta= 0 is accepted at 1% level of
        significance")
36 }

```

R code Exa 9.4.b Confidence interval estimate

```

1 x<-c(45, 50, 55, 60, 65, 70, 75)
2 y<-c(24.2, 25.0, 23.3, 22.0, 21.5, 20.6, 19.8)
3 xbar = mean(x);
4 ybar= mean(y);
5 n= 7;
6 SxY = 0;
7 for (i in 1:n){
8     SxY = SxY + (x[i]*y[i]) - (xbar*ybar)
9 }
10
11 Sxx = 0;
12 for (i in 1:n){
13     Sxx= Sxx + (x[i]*x[i]) - (xbar*xbar);
14 }
15 SY Y = 0;
16 for (i in 1:n){
17     SY Y = SY Y + (y[i]*y[i]) - (ybar*ybar);
18 }
19 B = SxY/Sxx;
20 A = ybar - (B*xbar);
21 cat("A is",A);
22 cat("B is",B);
23

```

```

24 SSR = ((Sxx*SYy) - (SxY*SxY))/Sxx ;
25 cat("The SSR is",SSR)
26
27 tvalue= qt(0.975, 5);
28 cat("tvalue is",tvalue);
29
30 k = sqrt(SSR/((n-2)*Sxx))*tvalue;
31 int1 = B + k;
32 int2= B-k;
33 cat("The 95% confidence interval is ",int2, "to ",
      int1)

```

R code Exa 9.4.c Regression to the mean

```

1 x<-c(60, 62, 64, 65, 66, 67, 68, 70, 72, 74)
2 y<-c(63.6, 65.2, 66, 65.5, 66.9, 67.1, 67.4, 68.3,
      70.1, 70)
3 plot(x,y,xlab ="Father 's Height",ylab = "Son 's
      Height")
4 xbar = mean(x)
5 ybar= mean(y)
6 n= 10;
7 SxY = 0
8 for (i in 1:n){
9   SxY = SxY + (x[i]*y[i]) - (xbar*ybar);
10 }
11
12 Sxx = 0;
13 for (i in 1:n){
14   Sxx= Sxx + (x[i]*x[i]) - (xbar*xbar);
15 }
16
17 SYy = 0;
18 for (i in 1:n){
19   SYy = SYy + (y[i]*y[i]) - (ybar*ybar);

```

```

20 }
21
22 B = SxY/Sxx;
23 A = ybar - (B*xbar);
24 cat("A is",A);
25 cat("B is",B);
26 p= seq(60,72,0.1)
27 q= A + B*p;
28 lines(p,q)
29 SSR = ((Sxx*SYy) - (SxY*SxY))/Sxx ;
30 ts = sqrt(((n-2)*Sxx)/SSR)*(B-1)
31 ts
32 tvalue= qt(0.99,n-2)
33 tvalue
34 if(ts<(-1*tvalue)){
35   cat("Null hypothesis is rejected at 1% level of
      significance")
36 } else{
37   cat("Null hypothesis is accepted at 1% level of
      significance")
38 }

```

R code Exa 9.4.d Motor vehicle deaths

```

1 x<-c(121, 96, 85, 113, 102, 118, 90, 84, 107, 112,
      95, 101)
2 y<-c(104, 91, 101, 110, 117, 108, 96, 102, 114, 96,
      88, 106)
3
4 plot(x,y,xlab = "Deaths in 1988",ylab = "Deaths in
      1989");
5
6 xbar = mean(x);
7 ybar= mean(y);
8 n= 12;

```

```

9
10 SxY = 0
11 for (i in 1:n){
12   SxY = SxY + (x[i]*y[i]) - (xbar*ybar);
13 }
14
15 Sxx = 0;
16 for (i in 1:n){
17   Sxx= Sxx + (x[i]*x[i]) - (xbar*xbar);
18 }
19
20 SYy = 0;
21 for (i in 1:n){
22   SYy = SYy + (y[i]*y[i]) - (ybar*ybar);
23 }
24
25 B = SxY/Sxx;
26 A = ybar - (B*xbar);
27 cat("A is",A);
28 cat("B is",B)

```

R code Exa 9.4.e Confidence interval for height

```

1 x<-c(60, 62, 64, 65, 66, 67, 68, 70, 72, 74)
2 y<-c(63.6, 65.2, 66, 65.5, 66.9, 67.1, 67.4, 68.3,
      70.1, 70)
3 x0 = 68;
4 xbar = mean(x);
5 ybar= mean(y);
6 n= 10;
7 SxY = 0
8 for (i in 1:n){
9   SxY = SxY + (x[i]*y[i]) - (xbar*ybar);
10 }
11

```

```

12 Sxx = 0;
13 for (i in 1:n){
14     Sxx= Sxx + (x[i]*x[i]) - (xbar*xbar);
15 }
16
17 SYX = 0;
18 for (i in 1:n){
19     SYX = SYX + (y[i]*x[i]) - (ybar*xbar);
20 }
21 B = SYX/Sxx;
22 A = ybar - (B*xbar);
23 tvalue= qt(0.975, n-2);
24 SSR = ((Sxx*SYX)- (SYX*SYX))/Sxx ;
25 intvl = A + (B*x0);
26 change = sqrt((1/n)+(((x0-xbar)^2)/Sxx))* sqrt(SSR/(
      n-2))*tvalue;
27 intvl1 = intvl - change;
28 intvl2= intvl + change;
29 cat("The 95% confidence interval is ",intvl1, "to ",
      intvl2)

```

R code Exa 9.4.f Confidence interval for height

```

1 x<-c(60, 62, 64, 65, 66, 67, 68, 70, 72, 74)
2 y<-c(63.6, 65.2, 66, 65.5, 66.9, 67.1, 67.4, 68.3,
      70.1, 70)
3 x0 = 68;
4 xbar = mean(x);
5 ybar= mean(y);
6 n= 10;
7 SYX = 0
8 for (i in 1:n){
9     SYX = SYX + (x[i]*y[i]) - (xbar*ybar);
10 }
11

```

```

12 Sxx = 0;
13 for (i in 1:n){
14     Sxx= Sxx + (x[i]*x[i]) - (xbar*xbar);
15 }
16
17 SYX = 0;
18 for (i in 1:n){
19     SYX = SYX + (y[i]*x[i]) - (ybar*xbar);
20 }
21 B = SYX/Sxx;
22 A = ybar - (B*xbar);
23 tvalue= qt(0.975, n-2);
24 SSR = ((Sxx*SYX)- (SYX*SYX))/Sxx ;
25 intvl = A + (B*x0);
26 change = sqrt(((n+1)/n)+(((x0-xbar)^2)/Sxx))* sqrt(
        SSR/(n-2))*tvalue;
27 intvl1 = intvl - change;
28 intvl2= intvl + change;
29 cat("The 95% confidence interval is ",intvl1, "to ",
        intvl2)

```

R code Exa 9.5.a Height of son and father

```

1 x<-c(60, 62, 64, 65, 66, 67, 68, 70, 72, 74)
2 y<-c(63.6, 65.2, 66, 65.5, 66.9, 67.1, 67.4, 68.3,
        70.1, 70)
3
4 xbar = mean(x);
5 ybar= mean(y);
6 n= 10;
7
8 SYX = 0
9 for (i in 1:n){
10     SYX = SYX + (x[i]*y[i]) - (xbar*ybar);
11 }

```

```

12
13 Sxx = 0;
14 for (i in 1:n){
15     Sxx= Sxx + (x[i]*x[i]) - (xbar*xbar);
16 }
17
18 SYX = 0;
19 for (i in 1:n){
20     SYX = SYX + (y[i]*x[i]) - (ybar*xbar);
21 }
22
23 B = SYX/Sxx;
24 A = ybar - (B*xbar);
25
26 SSR = ((Sxx*SYX)- (SYX*SYX))/Sxx ;
27 R2 = 1 - (SSR/SYX);
28 cat("The coefficient of determination is",R2)

```

R code Exa 9.7.a Percentage of chemical used

```

1 x<-c(5, 10, 20, 30, 40, 50, 60, 80)
2 yold<-c(0.061, 0.113, 0.192, 0.259, 0.339, 0.401,
          0.461, 0.551)
3 plot(x, yold);
4 y = -1*log(1-yold);
5 plot(x, y);
6
7
8 xbar = mean(x);
9 ybar= mean(y);
10 n= 8;
11
12 SYX = 0
13 for (i in 1:n){
14     SYX = SYX + (x[i]*y[i]) - (xbar*ybar);

```



```

15 }
16
17 Sxx = 0;
18 for (i in 1:n){
19     Sxx= Sxx + (x[i]*x[i]) - (xbar*xbar);
20 }
21
22 SYX = 0;
23 for (i in 1:n){
24     SYX = SYX + (y[i]*x[i]) - (ybar*xbar);
25 }
26
27 B = SYX/Sxx;
28 A = ybar - (B*xbar);
29 SSR = ((Sxx*SYX) - (SYX*SYX))/Sxx ;
30 chat = exp(-1*A);
31 dhat = 1 - exp(-1*B);
32 phat = 1 - chat*(1-dhat)^x
33 pres = yold - phat
34 cat("chat is",chat);
35 cat("dhat is",dhat);
36 cat("phat is",phat);
37 cat("p - phat, i.e. the residuals are",pres)

```

R code Exa 9.8.b Distance vs Travel Time

```

1 x<-c(0.5, 1, 1.5, 2, 3, 4, 5, 6, 8, 10)
2 y<-c(15, 15.1, 16.5, 19.9, 27.7, 29.7, 26.7, 35.9,
      42, 49.4)
3 w<-c()
4 for (i in 1:10){
5     w[i] = 1/x[i]
6 }
7 n = 10;
8 p = matrix(0,2,2)

```

```

 9  q = matrix(0,2,1)
10  p[1, 1] = sum(w);
11  p[1,2] = n;
12  p[2,1] = n;
13  p[2,2] = sum(x);
14  new<-c()
15  for (i in 1:10){
16    new[i] = w[i]*y[i]
17  }
18
19  q[1,1]= 1*sum(new);
20  q[2,1] = 1*sum(y);
21  sol = solve(p,q);
22  A = sol[1,1];
23  B = sol[2,1];
24  cat("A is",A);
25  cat("B is",B);
26  plot(x,y)
27  l = seq(0,10,0.1)
28  k = A + B*l
29  lines(l,k)

```

R code Exa 9.9.a Polynomial Fitting

```

1  x<-seq(1,10,1);
2  y<-c(20.6, 30.8, 55, 71.4, 97.3, 131.8, 156.3,
      197.3, 238.7, 291.7)
3  plot(x, y,xlab = "X",ylab = "Y");
4  n = length(x)
5  xsquared = x^2;
6  xcube = x^3;
7  xfour = x^4;
8  xy = x*y;
9  x2y = xy*x;
10 p= matrix(0,3,3);

```

```

11 q = matrix(0,3,1);
12 p[1,1] = n;
13 p[1,2] = sum(x);
14 p[1,3]=sum(xsquared);
15 p[2,1] = sum(x);
16 p[2,2] = sum(xsquared);
17 p[2,3]=sum(xcube);
18 p[3,1] = sum(xsquared);
19 p[3,2] = sum(xcube);
20 p[3,3]=sum(xfour);
21 q[1,1]= 1*sum(y);
22 q[2,1] = 1*sum(xy);
23 q[3,1] = 1*sum(x2y);
24 B= solve(p, q);
25 cat("B0 is",B[1,1]);
26 cat("B1 is",B[2,1]);
27 cat("B2 is",B[3,1])

```

R code Exa 9.10.a Multiple Linear Regression

```

1 x1<-c(679, 1420, 1349, 296, 6975, 323, 4200, 633)
2 x2<-c(30.4, 34.1, 17.2, 26.8, 29.1, 18.7, 32.6,
      32.5)
3 y = matrix(1,8,1);
4 n = 8
5 y<-c(11.6,16.1, 9.3, 9.1, 8.4, 7.7, 11.3, 8.4)
6 x = matrix(1,8,3);
7 for (i in 1:8){
8   x[i,2]= x1[i];
9   x[i,3]= x2[i];
10 }
11
12 pro1 = t(x)
13 pro2= pro1*%x;
14 pro3 = solve(pro2);

```

```

15 pro4 = pro3%%pro1;
16 pro5 = pro4%%y;
17
18 B<-matrix(0,3,8)
19 matrix(1,3,1);
20 for (i in 1:3){
21   B[i,1]= 0;
22   for (k in 1:8){
23     B[i,1]=B[i,1]+(pro4[i, k]*y[k]);
24   }
25 }
26 cat(B);
27 SSR = t(y)
28 SSR= SSR%%y;
29 sub = t(B);
30 sub = sub%%t(x);
31 sub= sub%%y;
32 SSR =SSR[1,1] - sub[1,1];
33 cat("SSr is",SSR)

```

R code Exa 9.10.b Estimate of variance

```

1 x1<-c(679, 1420, 1349, 296, 6975, 323, 4200, 633)
2 x2<-c(30.4, 34.1, 17.2, 26.8, 29.1, 18.7, 32.6,
      32.5)
3 y = matrix(1,8,1);
4 n = 8
5 y<-c(11.6,16.1, 9.3, 9.1, 8.4, 7.7, 11.3, 8.4)
6 x = matrix(1,8,3);
7 for (i in 1:8){
8   x[i,2]= x1[i];
9   x[i,3]= x2[i];
10 }
11
12 pro1 = t(x)

```

```

13 pro2= pro1%*%x;
14 pro3 = solve(pro2);
15 pro4 = pro3%*%pro1;
16 pro5 = pro4%*%y;
17
18 B<-matrix(0,3,8)
19 matrix(1,3,1);
20 for (i in 1:3){
21   B[i,1]= 0;
22   for (k in 1:8){
23     B[i,1]=B[i,1]+(pro4[i, k]*y[k]);
24   }
25 }
26 cat(B);
27 SSR = t(y)
28 SSR= SSR%*%y;
29 sub = t(B);
30 sub = sub%*%t(x);
31 sub= sub%*%y;
32 SSR =SSR[1,1] - sub[1,1];
33 cat("SSr is",SSR)
34
35 k=2;
36 den = n-k-1;
37 sigma = SSR/den;
38 cat("The variance is",sigma)

```

R code Exa 9.10.c Diameter of a tree

```

1 x1<-c(44, 33, 33, 32, 34, 31, 33, 30, 34, 34, 33,
      36, 33, 34, 37)
2 x2<-c(1.3, 2.2, 2.2, 2.6, 2.0, 1.8, 2.2, 3.6, 1.6,
      1.5, 2.2, 1.7, 2.2, 1.3, 2.6)
3 x3<-c(250, 115, 75, 85, 100, 75, 85, 75, 225, 250,
      255, 175, 75, 85, 90)

```

```

4  x4<-c(0.63, 0.59, 0.56, 0.55, 0.54, 0.59, 0.56,
        0.46, 0.63, 0.60, 0.63, 0.58, 0.55, 0.57, 0.62)
5  y<-c(18.1, 19.6, 16.6, 16.4, 16.9, 17.0, 20.0, 16.6,
        16.2, 18.5, 18.7, 19.4, 17.6, 18.3, 18.8)
6  n =length(x1);
7  x= matrix(1,15, 5);
8  for (i in 1:15){
9    x[i,2]= x1[i]
10   x[i,3]= x2[i]
11   x[i,4]= x3[i]
12   x[i,5]= x4[i]
13 }
14
15 pro1 = t(x)
16 pro2= pro1**%x
17 pro3 = solve(pro2)
18 pro4 = pro3**%pro1
19 pro5 = pro4**%y
20 B<-matrix(0,5,15)
21 for (i in 1:5){
22   B[i,1]= 0
23   for (k in 1:15){
24     B[i,1]=B[i,1]+(pro4[i, k]*y[k])
25   }
26 }
27 SSR = t(y)
28 SSR= SSR**%y
29 sub = t(B)
30 sub = sub**%t(x)
31 sub= sub**%y
32 SSR =SSR[1,1] - sub[1,1]
33 SSR
34 xxinv = 0.379;
35 k= 4;
36 ts = sqrt((n-k-2)/SSR)*B[2]/0.616;
37 pvalue = 2*(1- pt(ts, n-k-2))
38 cat("The p-value is ",pvalue)
39 '

```

40 The SSR calculated by R is 19.34 whereas the
textbook gives the value as 19.26 , thus the
difference in the final answer.
41 ,

R code Exa 9.10.d Estimating hardness

```
1 y<-c(79.2, 64.0, 55.7, 56.3, 58.6, 84.3, 70.4, 61.3,
      51.3, 49.8)
2 x1<-c(0.02, 0.03, 0.03, 0.04, 0.10, 0.15, 0.15,
      0.09, 0.13, 0.09)
3 x2<-c(1.05, 1.20, 1.25, 1.30, 1.30, 1.00, 1.10,
      1.20, 1.40, 1.40)
4 tvalue= 2.365;
5 x = matrix(1,10,3);
6 for (i in 1:10){
7   x[i,2]= x1[i]
8   x[i,3]= x2[i]
9 }
10
11 pro1 = t(x)
12 pro2= pro1*%x
13 pro3 = solve(pro2)
14 pro4 = pro3*%pro1
15 pro5 = pro4*%y
16 B= matrix(1,3,1)
17
18 for (i in 1:3){
19   B[i,1]= 0
20   for ( k in 1:10){
21     B[i,1]= B[i,1]+(pro4[i, k]*y[k])
22   }
23 }
24
25 SSR = t(y)
```

```

26 SSR= SSR%%y
27 sub = t(B)
28 sub = sub%%t(x)
29 sub= sub%%y
30 SSR =SSR[1,1] - sub[1,1]
31 cat("SSr is",SSR)
32
33 smallx<-c(1, 0.15, 1.15)
34 product = smallx %% B
35 n = 10;
36 k=2;
37 val= sqrt(SSR/(n-k-1));
38
39 pro5 = smallx %% pro3
40 pro6 = pro5 %% (smallx)
41 pro7 = val*sqrt(pro6)*tvalue
42
43 up = product + pro7;
44 low = product - pro7;
45 cat(" 95% confidence interval is from ",low,"to",up)

```

R code Exa 9.11.a Animal sick falling

```

1 cancer = 84;
2 total = 111;
3 level = 250;
4 alpha= -1*log((total-cancer)/total)/level;
5 cat("Alpha is ",alpha)

```

Chapter 10

Analysis of Variance

R code Exa 10.3.a Dependence of mileage on gas used

```
1 Xij<-matrix(c(220, 251, 226, 246, 260, 244, 235,
               232, 242, 225, 252, 272, 250, 238, 256),nrow = 3,
               ncol = 5,byrow = TRUE)
2 Xi = matrix(0,3,1);
3 n= 5;
4 m=3;
5 for (i in 1:3){
6   for (j in 1:5){
7     Xi[i]= Xi[i] + Xij[i,j]
8   }
9 }
10 Xi
11 Xi = Xi/n;
12 SSW= 0;
13 for (i in 1:3){
14   for (j in 1:5){
15     SSW = SSW + ((Xij[i,j]-Xi[i])^2)
16   }
17 }
18 SSW
19 sigma1 = SSW/((n*m)-m);
```

```

20 Xdotdot = sum(Xi)/m;
21 new = (Xi - Xdotdot)^2;
22 SSb= n*sum(new);
23 sigma2 = SSb/(m-1);
24 TS = sigma2/sigma1;
25 cat(" Value of the test statistic is",TS)
26 pvalue = 1 - pf(TS,m-1, ((n*m)-m))
27 cat("The p-value is",pvalue)
28 if(pvalue>0.05){
29     cat( "Since the p-value is greater than .05, the
           null hypothesis that the mean mileage is the
           same for all 3 brands of gasoline cannot be
           rejected. ")
30 }

```

R code Exa 10.3.b Dependence of mileage on gas used

```

1 Xijold<-matrix(c(220, 251, 226, 246, 260, 244, 235,
                  232, 242, 225, 252, 272, 250, 238, 256),nrow = 3,
                  ncol = 5,byrow = TRUE)
2 Xij = Xijold - 220;
3 m=3;
4 n=5;
5 Xidot = matrix(0,3,1);
6 for (i in 1:m){
7     for (j in 1:n){
8         Xidot[i]=Xidot[i] + Xij[i,j]
9     }
10 }
11 Xidot = Xidot/n;
12 Xdotdot = sum(Xidot)/m;
13 SSb=0;
14 for (i in 1:m){
15     SSb = SSb + (Xidot[i]-Xdotdot)^2
16 }

```

```

17 SSb = SSb*n;
18 Xijsquared = Xij^2;
19 SSW = sum(Xijsquared) - (m*n*(Xdotdot^2)) - SSb;
20 sigma1 = SSW/((n*m)-m);
21 sigma2 = SSb/(m-1);
22 TS = sigma2/sigma1;
23 cat("Value of the test statistic is",TS)

```

R code Exa 10.3.c Difference in GPA

```

1 Xij<-matrix(c(3.2, 3.4, 3.3, 3.5, 3.4, 3.0, 3.7,
               3.3, 2.8, 2.6, 3.0, 2.7),nrow = 3,ncol = 4,byrow
               = TRUE)
2 Xi = matrix(0,3,1);
3 n= 4;
4 m=3;
5 for (i in 1:3){
6   for (j in 1:4){
7     Xi[i]= Xi[i] + Xij[i,j]
8   }
9 }
10 Xi = Xi/n;
11 SSW= 0;
12 for (i in 1:3){
13   for (j in 1:4){
14     SSW = SSW + ((Xij[i,j]-Xi[i])^2)
15   }
16 }
17 sigma1 = SSW/((n*m)-m);
18 Xdotdot = sum(Xi)/m;
19 new = (Xi - Xdotdot)^2;
20 SSb= n*sum(new);
21 sigma2 = SSb/(m-1);
22 TS = sigma2/sigma1;
23 cat("Value of the test statistic is",TS)

```

```

24 pvalue = 1 - pf(TS,m-1, ((n*m)-m) );
25 cat("The p-value is",pvalue)
26 C = 3.95; #from table A5
27 W = C*sqrt(SSW/(9*4));
28 W
29 cat("Mean1 - Mean2 lies between ", Xi[1]-Xi[2]-W ,"
      and ", Xi[1]-Xi[2]+W)
30 cat("Mean1 - Mean3 lies between ", Xi[1]-Xi[3]-W ,"
      and ", Xi[1]-Xi[3]+W)
31 cat("Mean2 - Mean3 lies between ", Xi[2]-Xi[3]-W ,"
      and ", Xi[2]-Xi[3]+W)

```

R code Exa 10.4.b Estimating Parameters

```

1 X<-matrix(c(75, 73, 60, 70, 86, 78, 71, 64, 72, 90,
             80, 69, 62, 70, 85, 73, 67, 63, 80, 92),nrow = 4,
           ncol = 5,byrow = TRUE)
2 Xidot = matrix(0,4,1);
3 for (i in 1:4){
4   for (j in 1:5){
5     Xidot[i]=Xidot[i] + X[i,j]
6   }
7 }
8 Xidot = Xidot/5;
9 Xjdot = matrix(0,5,1);
10 for (j in 1:5){
11   for (i in 1:4){
12     Xjdot[j]=Xjdot[j] + X[i,j]
13   }
14 }
15 Xjdot = Xjdot/4;
16 Xdotdot = sum(Xidot)/4;
17 meanhat = Xdotdot;
18 alphahat = Xidot - meanhat;
19 betahat = Xjdot - meanhat;

```

```

20 cat("The estimator of the mean is",meanhat)
21 cat("The alphas are-",alphahat)
22 cat("The betas are-",betahat)

```

R code Exa 10.5.a Species collected

```

1 X<-matrix(c(53, 35, 31, 37, 40, 43, 36, 34, 17, 21,
             30, 18, 47, 37, 17, 31, 45, 26, 55, 31, 17, 23,
             43, 37, 40, 32, 19, 26, 45, 37, 52, 42, 20, 27,
             26, 32, 39, 28, 21, 21, 36, 28, 40, 32, 21, 21,
             36, 35),nrow =8,ncol = 6, byrow = TRUE)
2 m= 8;
3 n = 6;
4 Xidot = matrix(0,8,1);
5
6 for (i in 1:8){
7   for (j in 1:6){
8     Xidot[i]=Xidot[i] + X[i,j]
9   }
10 }
11 Xidot = Xidot/6;
12 Xjdot = matrix(0,6,1);
13
14 for (j in 1:6){
15   for (i in 1:8){
16     Xjdot[j]=Xjdot[j] + X[i,j]
17   }
18 }
19
20 Xjdot = Xjdot/8;
21 Xdotdot = sum(Xidot)/8;
22 new = (Xidot - Xdotdot)^2;
23 SSr = n*sum(new);
24 new1 = (Xjdot - Xdotdot)^2;
25 SSc = m*sum(new1);

```

```

26 SSe = 0;
27 for (i in 1:m){
28     for (j in 1:n){
29         SSe = SSe + (X[i,j]-Xidot[i]-Xjdot[j]+ Xdotdot)
                ^2
30     }
31 }
32
33 N =(m-1)*(n-1)
34 TS1 = SSr*N/((m-1)*SSe)
35 TS2 = SSr*N/((n-1)*SSe)
36 pvaluec = 1- pf(TS1, m-1, N)
37 pvaluer = 1- pf(TS2, n-1, N)
38 cat("The value of the F-statistic for testing that
    there is no row effect is",TS1)
39 cat("The p-value for testing that there is no row
    effect is",pvaluec)
40 cat("The value of the F-statistic for testing that
    there is no column effect is",TS2)
41 cat("The p-value for testing that there is no column
    effect is",pvaluer)

```

Chapter 11

Goodness of Fit Tests and Categorical Data Analysis

R code Exa 11.2.a Relation between death date and birth date

```
1 X<-c(90, 100, 87, 96, 101, 86, 119, 118, 121, 114,
      113, 106)
2 pi= matrix(1,12,1);
3 pi= pi/12;
4 new = X^2;
5 npi= sum(X)*pi;
6 T = sum(new);
7 T = T/npi;
8 T = T - sum(X);
9 cat("When there are 12 regions")
10 cat("The test statistic is",T[1])
11 pvalue = 1- pchisq(T[1], 11);
12 cat("The pvalue is ",pvalue)
13
14 X<-c(277, 283, 358, 333)
15 pi= matrix(1,4,1);
16 pi= pi/4;
17 new = X^2;
18 npi= sum(X)*pi;
```

```

19 T = sum(new);
20 T = T/np;
21 T = T - sum(X);
22 cat("When there are 4 regions")
23 cat("The test statistic is",T[1])
24 pvalue = 1- pchisq(T[1], 3);
25 cat("The pvalue is ",pvalue)

```

R code Exa 11.2.b Quality of bulbs

```

1 X<-c(3, 6, 9, 7, 5)
2 p<-c(0.15, 0.25, 0.35, 0.20, 0.05)
3 T= 0;
4 n3=sum(X);
5 np = p*n3;
6 Xsqu = (X-np)^2;
7 cat(Xsqu);
8 XT = Xsqu/np;
9 T = sum(XT);
10
11 cat("The test statistic is",T)
12 pvalue = 1- pchisq(T[1], 4)
13 cat("The pvalue is ",pvalue)
14 cat("Thus, the hypothesis would not be rejected at
    5% level of significance")

```

R code Exa 11.2.d Six outcomes

```

1 X<-c(3, 3, 5, 18, 4, 7)
2 p<-c(0.1, 0.1, 0.05, 0.4, 0.2, 0.15)
3 psimu = 0.1843
4 num= 10000;
5 T= 0;

```



```

6 n=sum(X);
7 np = n*p;
8 Xsqu = X^2;
9 for (i in 1:6){
10   T = T + (Xsqu[i]/np[i])
11 }
12 T = T - sum(X)
13
14 cat("The test statistic is",T[1])
15 pvalue = 1- pchisq(T[1], 5);
16 int1 = psimu - (1.645*sqrt(psimu*(1-psimu)/num));
17 int2 = psimu + (1.645*sqrt(psimu*(1-psimu)/num));
18 cat("With 90% confidence p-value lies between ",int1
    , "and",int2)

```

R code Exa 11.3.a Weekly accidents

```

1 Y<-c(8, 0, 0, 1, 3, 4, 0, 2, 12, 5, 1, 8, 0, 2, 0,
      1, 9, 3, 4, 5, 3, 3, 4, 7, 4, 0, 1, 2, 1, 2)
2 weeks = 30;
3 lamda = sum(Y)/weeks;
4 p = matrix(0,5,1);
5 p[1] = ppois(0, lamda);
6 p[2] = ppois(1, lamda) - p[1];
7 p[3] = ppois(3, lamda) - ppois(1, lamda);
8 p[4] = ppois(5, lamda) - ppois(3, lamda);
9 p[5] = 1 - ppois(5, lamda);
10
11 X = matrix(0,5,1);
12 for (i in 1:30){
13   if(Y[i]==0){
14     X[1] = X[1] +1;
15   } else if(Y[i]==1){
16     X[2] = X[2] +1;
17   } else if(Y[i]==2){

```

```

18     X[3] = X[3] +1;
19 } else if(Y[i]==3){
20     X[3] = X[3] +1;
21 } else if(Y[i]==4){
22     X[4] = X[4] +1;
23 } else if(Y[i]==5){
24     X[4] = X[4] +1;
25 } else if(Y[i]>5){
26     X[5] = X[5] +1;
27 }
28 }
29
30 T= 0;
31 npi = weeks * p;
32 for (i in 1:5){
33     T = T + ((X[i]-npi[i])^2)/npi[i]
34 }
35 cat("T is",T)
36 pvalue = 1- pchisq(T, 3)
37 cat("The p-value is",pvalue)
38 cat("Hypothesis of an underlying poisson
      distribution is rejected")

```

R code Exa 11.4.a Political affiliation and Gender

```

1 Nij<-matrix(c(68, 56, 32, 52, 72, 20),nrow = 2,ncol
      = 3,byrow = TRUE)
2 n= sum(Nij);
3 Ni = matrix(0,2,1);
4 Mj = matrix(0,3,1);
5 for (i in 1:2){
6     for (j in 1:3){
7         Ni[i] = Ni[i] + Nij[i,j]
8     }
9 }

```

```

10 for (j in 1:3){
11   for (i in 1:2){
12     Mj[j] = Mj[j] + Nij[i,j]
13   }
14 }
15 NM = matrix(1,2,3);
16 for (i in 1:2){
17   for (j in 1:3){
18     NM[i,j]= Ni[i]*Mj[j]
19   }
20 }
21 NM= NM/n;
22 TS = 0
23 for (i in 1:2){
24   for (j in 1:3){
25     TS = TS + ((Nij[i,j]-NM[i,j])^2)/NM[i,j]
26   }
27 }
28 cat("The test statistic is",TS)
29 compare = pchisq(2, 0.95, 0.05)
30 if(TS>compare){
31   cat("The null hypothesis is rejected at the 5%
      level of significance")
32 } else {
33   disp("The null hypothesis is accepted at the 5%
      level of significance")
34 }

```

R code Exa 11.4.b Machine Breakdown and shift

```

1 Nij<-matrix(c(10, 12, 6, 7, 10, 24, 9, 10, 13, 20,
      7, 10),nrow = 3,ncol = 4,byrow = TRUE)
2 n= sum(Nij);
3 Ni = matrix(0,3,1);
4 Mj = matrix(0,4,1);

```

```

5 for (i in 1:3){
6   for (j in 1:4){
7     Ni[i] = Ni[i] + Nij[i,j]
8   }
9 }
10 for (j in 1:4){
11   for (i in 1:3){
12     Mj[j] = Mj[j] + Nij[i,j]
13   }
14 }
15 NM = matrix(1,3,4);
16 for (i in 1:3){
17   for (j in 1:4){
18     NM[i,j]= Ni[i]*Mj[j]
19   }
20 }
21 NM= NM/n;
22 TS = 0
23 for (i in 1:3){
24   for (j in 1:4){
25     TS = TS + ((Nij[i,j]-NM[i,j])^2)/NM[i,j]
26   }
27 }
28 cat("The test statistic is",TS)
29 pvalue = 1- pchisq(TS, 6)
30 cat("the hypothesis that the machine that causes a
    breakdown is independent of the shift on which
    the breakdown occurs is accepted.")

```

R code Exa 11.5.a Lung cancer and smoking

```

1 Nij<-matrix(c(62, 14, 9938, 19986),nrow = 2,ncol =
  2,byrow = TRUE)
2 n= sum(Nij);
3 Ni = matrix(0,2,1);

```

```

4  Mj = matrix(0,2,1);
5  for (i in 1:2){
6    for (j in 1:2){
7      Ni[i] = Ni[i] + Nij[i,j]
8    }
9  }
10 Ni
11 for (j in 1:2){
12   for (i in 1:2){
13     Mj[j] = Mj[j] + Nij[i,j]
14   }
15 }
16
17 NM = matrix(1,2,2);
18 for (i in 1:2){
19   for (j in 1:2){
20     NM[i,j]= Ni[i]*Mj[j]
21   }
22 }
23 NM= NM/n;
24 NM
25 TS = 0
26 for (i in 1:2){
27   for (j in 1:2){
28     TS = TS + ((Nij[i,j]-NM[i,j])^2)/NM[i,j]
29   }
30 }
31 cat("The test statistic is",TS)
32 compare = pchisq(1, 0.99, 0.01)
33 compare
34 if(TS>compare){
35   cat("The null hypothesis is rejected at the 1%
      level of significance");
36 } else{
37   cat("The null hypothesis is accepted at the 1%
      level of significance");
38 }

```

R code Exa 11.5.b Females reporting abuse

```
1 Nij<-matrix(c(28, 30, 58, 55, 472, 470, 442, 445),
  nrow = 2,ncol = 4,byrow = TRUE)
2 n= sum(Nij);
3 Ni = matrix(0,2,1);
4 Mj = matrix(0,4,1);
5 for (i in 1:2){
6   for (j in 1:4){
7     Ni[i] = Ni[i] + Nij[i,j]
8   }
9 }
10 for (j in 1:4){
11   for (i in 1:2){
12     Mj[j] = Mj[j] + Nij[i,j]
13   }
14 }
15 NM = matrix(1,2,4);
16 for (i in 1:2){
17   for (j in 1:4){
18     NM[i,j]= Ni[i]*Mj[j]
19   }
20 }
21 NM= NM/n;
22
23 TS = 0
24 for (i in 1:2){
25   for (j in 1:4){
26     TS = TS + ((Nij[i,j]-NM[i,j])^2)/NM[i,j]
27   }
28 }
29 cat("The test statistic is",TS)
30
31 compare = pchisq(3, 0.99, 0.01)
```

```

32 pvalue = 1- pchisq(TS, 3)
33 cat("The p-value is",pvalue)
34
35 compare
36 if(TS>compare){
37   cat("The null hypothesis is rejected at the 1%
      level of significance");
38 } else{
39   cat("The null hypothesis is accepted at the 1%
      level of significance");
40 }

```

R code Exa 11.6.a Testing distribution of a population

```

1 X<-c(66, 72, 81, 94, 112, 116, 124, 140, 145, 155);
2 D= 0.4831487;
3 n= 10;
4 Dgiven = 1.480;
5 Dstar = (sqrt(n) + 0.12 + (0.11/sqrt(n)))*D;
6 cat("Dstar is ",Dstar);
7 if(Dstar>Dgiven){
8   cat("Null hypothesis is rejected at 2.5% level of
      significance")
9 } else{
10   cat("Null hypothesis is accepted at 2.5% level
      of significance")
11 }

```

Chapter 12

Non parametric Hypothesis Tests

R code Exa 12.2.a testing the median

```
1 n= 200;
2 v = 120;
3 p =0.5;
4 if(v < (n/2)){
5   pvalue = 2*pbinom(v, n, p)
6 } else{
7   pvalue = 2*pbinom(n-v, n, p)
8 }
9 cat("Pvalue is ",pvalue)
```

R code Exa 12.2.b testing the median

```
1 n= 80;
2 v = 28;
3 p =0.5;
4
```



```

5 pvalue = pbinom(v, n, p);
6 cat("Pvalue is ",pvalue);
7 cat("Thus, the null hypothesis that the median
    income is less than or equal to $90,000 is
    rejected")

```

R code Exa 12.3.b Signed Rank Test

```

1 n =4;
2 mo = 2;
3 X<-c(4.2, 1.8, 5.3, 1.7)
4 t =3; #value of the test statistic (Ex12_3a)
5 tstar= min(t, (n*(n+1)/2) - t);
6 P = matrix(0,4,4);
7 P[1,1]= 0.5;
8 P[1,2] = 1;
9 P[1,3] = 1;
10 P[1,4] = 1;
11 for (i in 2:4){
12   for (j in 1:4){
13     if (j-i <1) {
14       P[i,j] = 0.5*P[i-1, j]
15     } else{
16       P[i,j] = 0.5*(P[i-1,j-i]+P[i-1,j])
17     }
18   }
19 }
20 P

```

R code Exa 12.3.c Determining Population Distribution

```

1 n =20;
2 t =142;

```

```

3  tstar= min(t, (n*(n+1)/2) - t);
4  P = matrix(1,20,tstar+1);
5  P[1,1]= 0.5;
6  P[1,2] = 1;
7  for (i in 2:20){
8    for (j in 1:(tstar+1)){
9      if (j-i <1){
10       P[i,j] = 0.5*P[i-1, j]
11     } else{
12       P[i,j] = 0.5*(P[i-1,j-i]+P[i-1,j])
13     }
14   }
15 }
16 pvalue= 2*P[20,tstar+1];
17 cat("Pvalue is",pvalue)
18 cat("Thus the hypothesis that the population
      distribution is symmetric about 0 is accepted at
      the alpha = 10% level of significance")

```

R code Exa 12.4.a Treatments against corrosion

```

1  X<-c(65.2, 67.1, 69.4, 78.2, 74, 80.3)
2  Y<-c(59.4, 72.1, 68, 66.2, 58.5)
3  Z = append(X,Y)
4  Z = sort(Z);
5  n= length(X);
6  q = length(Z)
7  T = 0;
8  for (i in 1:n){
9    test = X[i];
10   for (j in 1:q){
11     if(test== Z[j]){
12       T = T+ j;
13     }
14   }

```

```

15 }
16
17 cat("The test statistic is ",T)

```

R code Exa 12.4.b Determining P

```

1 prob <- function(N, M, K) {
2   if(N==1 & M==0){
3     if(K >0){
4       cat(result = 1)
5     } else {
6       cat(result =0)
7     }
8   } else if (N==0 & M==1){
9     if(K <0){
10      cat(result = 0)
11    } else {
12      cat(result =1)
13    }
14  } else if (N==0 & M==0 &K==0){
15    cat(result =1)
16  } else{
17    cat(result = (prob(N-1, M, K-N-M)*(N/(N+M)))
18              + (prob(N, M-1, K)*(M/(N+M))))
19  }
20
21 pval <- function(n,m,t) {
22   result = 2*min(prob(n,m,t), 1-prob(n,m,t-1));
23 }
24
25 '> pval(2,1,3)
26 001
27 Error: C stack usage 19923984 is too close to the
   limit

```

```

28
29 C:\Program Files\RStudio\bin>rstudio.exe --max-
    ppsize=5000000000
30
31 '

```

R code Exa 12.4.c Finding p value

```

1  prob <- function(N, M, K) {
2    if(N==1 & M==0){
3      if(K >0){
4        cat(result = 1)
5      } else {
6        cat(result =0)
7      }
8    } else if (N==0 & M==1){
9      if(K <0){
10       cat(result = 0)
11     } else {
12       cat(result =1)
13     }
14   } else if (N==0 & M==0 &K==0){
15     cat(result =1)
16   } else{
17     cat(result = (prob(N-1, M, K-N-M)*(N/(N+M))) + (
18       prob(N, M-1, K)*(M/(N+M))))
19   }
20 }
21 pval <- function(n,m,t) {
22   result = 2*min(prob(n,m,t), 1-prob(n,m,t-1));
23 }
24
25 '> pval(2,1,3)
26 001

```

```

27 Error: C stack usage 19923984 is too close to the
    limit
28
29 C:\Program Files\RStudio\bin>rstudio.exe --max-
    ppsize=5000000000
30
31 '
32 '> pval(5,6,21)
33 Error: C stack usage 19923984 is too close to the
    limit '

```

R code Exa 12.4.d Comparing production methods

```

1 prob <- function(N, M, K) {
2   if(N==1 & M==0){
3     if(K >0){
4       cat(result = 1)
5     } else {
6       cat(result =0)
7     }
8   } else if (N==0 & M==1){
9     if(K <0){
10      cat(result = 0)
11    } else {
12      cat(result =1)
13    }
14  } else if (N==0 & M==0 &K==0){
15    cat(result =1)
16  } else{
17    cat(result = (prob(N-1, M, K-N-M)*(N/(N+M))) + (
18      prob(N, M-1, K)*(M/(N+M))))
19  }
20
21 pval <- function(n,m,t) {

```

```

22   result = 2*min(prob(n,m,t), 1-prob(n,m,t-1));
23 }
24
25 '> pval(9,13,72)
26 001
27 Error: C stack usage 19923984 is too close to the
    limit
28
29 C:\Program Files\RStudio\bin>rstudio.exe --max-
    ppsize=5000000000
30
31 '

```

R code Exa 12.4.e Determining p value

```

1  n1 =5;
2  m1= 6;
3
4  t1 =21;
5  num1 = n1*(n1+m1+1)/2;
6  d1=abs(t1 - num1);
7  val = d1/sqrt(n1*m1*(n1+m1+1)/12);
8  pval = 2*(1- pnorm(val, 0,1))
9  cat("The p-value for eg 12.4a is",pval)
10 n2 =9;
11 m2= 13;
12 t2 =72;
13 d2=abs(t2 - n2*(n2+m2+1)/2);
14 val = d2/sqrt(n2*m2*(n2+m2+1)/12);
15 pval = 2*(1- pnorm(val, 0,1));
16 cat("The p-value for eg 12.4d is",pval)

```

R code Exa 12.5.a Testing randomness

```

1 fact <- function(num) {
2   if(num<=0){
3     result= 1
4   } else {
5     result = factorial(num)
6   }
7 }
8
9 proba <- function(n,m,k) {
10  if(k%%2==0){
11    k=k/2;
12    result = 2*fact(m-1)*fact(n-1)*fact(n)*fact(m)/(
13      fact(k-1)^2*fact(m-k)*fact(n-k)*fact(n+m));
14  } else {
15    k = (k-1)/2;
16    result = fact(m-1)*fact(n-1)*fact(n)*fact(m)/(
17      fact(k-1)*fact(k)*fact(m-k)*fact(n-k-1)*fact(
18        n+m)) + fact(m-1)*fact(n-1)*fact(n)*fact(m)/(
19        fact(k-1)*fact(k)*fact(m-k-1)*fact(n-k)*fact(
20        n+m));
21  }
22 }
23
24 r1 = 20;
25 n1 = 20;
26 m1=10;
27 ans1 =0;
28 for (i in 1:19){
29   ans1 =ans1 + proba(n1,m1,i);
30 }
31 if(ans1<0.5){
32   pvalue1 = 2*ans1;
33 } else {
34   pvalue1 = 2*(1-ans1);
35 }
36
37 cat("P-value is",pvalue1)

```

R code Exa 12.5.c Determining p value

```
1 u = 61;
2 sigma = 5.454;
3 r =75;
4 val = pnorm((r-u)/sigma, 0,1)
5 if(val>0.5){
6   pvalue = 2*(1-val)
7 } else{
8   pvalue = 2*val
9 }
10 cat("P-value is",pvalue)
```

Chapter 13

Quality Control

R code Exa 13.2.a Steel shaft diameter

```
1 X<-c(3.01, 2.97, 3.12, 2.99, 3.03, 3.02, 3.10, 3.14,
      3.09, 3.20)
2 Y<-seq(1,10,1)
3 u = 3;
4 sigma = 0.1;
5 n=4;
6 ucl = u + (3*sigma/sqrt(n));
7 lcl = u - (3*sigma/sqrt(n));
8 Z<-seq(0.1,10,0.1)
9 P= matrix(1,1,100)
10 Q= matrix(1,1,100)
11 P =P*ucl;
12 Q =Q*lcl;
13 plot(Y, X)
14 plot(Z, P)
15 plot(Z, Q)
16 cat('ucl is ',ucl)
17 cat('lcl is ',lcl)
```

R code Exa 13.2.b unknown mean and variance

```
1 Xbar<-c(3.01, 2.97, 3.12, 2.99, 3.03, 3.02, 3.10,
          3.14, 3.09, 3.20)
2 S<-c(0.12, 0.14, 0.08, 0.11, 0.09, 0.08, 0.15, 0.16,
        0.13, 0.16)
3 c<-c(0.7978849, 0.8862266, 0.9213181, 0.9399851,
        0.9515332, 0.9593684, 0.9650309, 0.9693103,
        0.9726596)
4 n=4;
5 Xbarbar= mean(Xbar);
6 Sbar =mean(S);
7 lcl = Xbarbar - (3*Sbar/(sqrt(n)*c(n-1)));
8 ucl = Xbarbar + (3*Sbar/(sqrt(n)*c(n-1)));
9 u = Xbarbar;
10 sigma= Sbar/c[n-1];
11 prob = pnorm(3.1, u, sigma) - pnorm(2.9, u, sigma);
12 cat("Percentage of the items that will meet the
      specifications is",prob*100)
```

R code Exa 13.3.a determining control limits

```
1 Xbar<-c(35.1, 33.2, 31.7, 35.4, 34.5, 36.4, 35.9,
          38.4, 35.7, 27.2, 38.1, 37.6, 38.8, 34.3, 43.2,
          41.3, 35.7, 36.3, 35.4, 34.6)
2 S<-c(4.2, 4.4, 2.5, 3.2, 2.6, 4.5, 3.4, 5.1, 3.8,
        6.2, 4.2, 3.9, 3.2, 4, 3.5, 8.2, 8.1, 4.2, 4.1,
        3.7)
3 c<-c(0.7978849, 0.8862266, 0.9213181, 0.9399851,
        0.9515332, 0.9593684, 0.9650309, 0.9693103,
        0.9726596)
4 Y<-seq(1,20,1)
5 n =5;
6 Z<-seq(0.1,20,0.1)
7 Xbarbar = mean(Xbar);
```

```

8 Sbar = mean(S);
9 lclX = Xbarbar - (3*Sbar/(sqrt(n)*c[n-1]));
10 uclX = Xbarbar + (3*Sbar/(sqrt(n)*c[n-1]));
11 val1 = 1/c[n-1]
12 val1 = val1^2;
13 val1 = val1 - 1;
14 val = sqrt(val1);
15 ucls = Sbar*(1+(3*val));
16 lcls = Sbar*(1-(3*val));
17 plot(Y, Xbar)
18 P<-matrix(1,1, 200)
19 Q<-matrix(1,1, 200)
20 P= P*lclX;
21 Q=Q*uclX;
22 cat('UCL(X)=',uclX)
23 cat('LCL(X)=',lclX)
24 plot(Z, P)
25 plot(Z, Q,main = 'Control Chart for X')
26 cat('UCL(S)=',uclX)
27 cat('LCL(S)=',lclX)
28 plot(Y, S)
29 P= P*lcls/lclX;
30 Q=Q*ucls/uclX;
31 plot(Z, P)
32 plot(Z, Q,main = 'Control Chart for S')

```

R code Exa 13.4.a Defectives Screws

```

1 defect<-c(6, 5, 3, 0, 1, 2, 1, 0, 2, 1, 1, 3, 2, 0,
            1, 1, 0, 2, 1, 2)
2 F<-c(0.12, 0.10, 0.06, 0.00, 0.02, 0.04, 0.02, 0.00,
       0.04, 0.02, 0.02, 0.06, 0.04, 0.00, .02, 0.02,
       0.00, 0.04, 0.02, 0.04)
3 total = 1000;
4 Fbar = sum(defect)/total;

```

```

5 n=50;
6 val = sqrt(Fbar*(1-Fbar)/n);
7 lcl = Fbar - (3*val);
8 ucl = Fbar + (3*val);
9 cat("LCL is" ,lcl)
10 cat("UCL is",ucl)
11 for (i in 1:20){
12     if( F[i]>ucl | F[i]<lcl) {
13         totald=sum(defect)-defect[i]
14     }
15 }
16 total = total - 50;
17 Fbar = totald/total;
18 val = sqrt(Fbar*(1-Fbar)/n);
19 cat("After recomputation");
20 lcl = Fbar - (3*val);
21 ucl = Fbar + (3*val);
22 cat("LCL is",lcl)
23 cat("UCL is",ucl)

```

R code Exa 13.5.a Control during production of cars

```

1 X<-c(141, 162, 150, 111, 92, 74, 85, 95, 76, 68, 63,
      74, 103, 81, 94, 68, 95, 81, 102, 73)
2 total = sum(X);
3 num = 20;
4 Xbar = mean(X);
5 lcl = Xbar - 3*sqrt(Xbar);
6 ucl = Xbar + 3*sqrt(Xbar);
7 cat("UCL is",ucl)
8 cat("LCL is",lcl)
9 for (i in 1:20){
10     if(X[i]> ucl){
11         total = total - X[i]
12         num= num -1

```

```

13   }
14 }
15 Xbar = total/num
16 lcl = Xbar - 3*sqrt(Xbar);
17 ucl = Xbar + 3*sqrt(Xbar);
18 cat("After recomputation")
19 cat("UCL is",ucl)
20 cat("LCL is",lcl)
21 total = total - X[4]
22 num = num-1;
23 cat("Xbar is",Xbar)
24 cat(" is",X[4])
25 Xbar = total/num
26 lcl = Xbar - 3*sqrt(Xbar);
27 ucl = Xbar + 3*sqrt(Xbar);
28 cat("After second recomputation")
29 cat("UCL is",ucl);
30 cat("LCL is",lcl);
31 cat("It appears that the process is in control with
    mean",Xbar);
32
33 'The mean after the second recomputation is
    incoreectly calculated in the textbook. It should
    be
34  $((17*84.41)-111)/16 = 82.748$  whereas the value
    given in the book is 82.56. The values of UCL and
    LCL
35 change accordingly.'

```

R code Exa 13.6.b Service Time

```

1 X<-c(48, 52, 70, 62, 57, 81, 56, 59, 77, 82, 78, 80,
      74, 82, 68, 84)
2 u = 62;
3 n = 4;

```

```

4 sigma = 24;
5 alpha = 0.25;
6 W = matrix(0,17);
7 W[1] = 60;
8 for (i in 2:17){
9   W[i] = (0.25*X[i-1]) + (0.75*W[i-1])
10 }
11 cat("The values of W are",W)
12 val = 3*sigma*sqrt(alpha/(n*(2-alpha)));
13 lcl = u- val;
14 ucl = u+ val;
15 cat("LCL is",lcl)
16 cat("UCL is",ucl)

```

R code Exa 13.6.c Exponentially weighted moving average control

```

1 X<-c(9.617728, 10.25437, 9.867195, 10.79338,
      10.60699, 10.48396, 13.33961, 9.462969, 10.14556,
      11.66342, 11.55484, 11.26203, 12.31473,
      9.220009, 11.25206, 10.48662, 9.025091, 9.693386,
      11.45989, 12.44213, 11.18981, 11.56674,
      9.869849, 12.11311, 11.48656)
2 t<-seq(1,26,1)
3 alpha = 2/9;
4 val = 9.915051 - (alpha*9.617728);
5 val = val/(1-alpha);
6 cat("val is",val)
7 u = 10;
8 n = 5;
9 sigma = 2;
10
11 W = matrix(0,26)
12 W[1] = 10.;
13 for (i in 2:26){
14   W[i] = (alpha*X[i-1]) + ((1-alpha)*W[i-1])

```

```

15 }
16 cat("The values of W are",W)
17 val = 3*sigma*sqrt(alpha/(n*(2-alpha)));
18 lcl = u- val;
19 ucl = u+ val;
20 cat("LCL is",lcl)
21 cat("UCL is",ucl)
22 plot(t,W,xlab ="t",ylab = "W")
23 nlcl = matrix(1,1, 26)
24 nlcl= nlcl* lcl;
25 lines(t,nlcl)
26 nucl = matrix(1, 1, 26);
27 nucl= nucl * ucl;
28 lines(t,nucl)
29
30 'The asymptotic lines for UCL and LCL have been
    plotted '

```

R code Exa 13.6.d Finding control limit

```

1 X<-c(29, 33, 35, 42, 36, 44, 43, 45)
2 u =30;
3 sig = 8;
4 d =0.5;
5 B =5;
6 Y = X - u - (d*sig);
7 S = matrix(0,9);
8 S[1] =0;
9 for (i in 2:9){
10   S[i]= max(S[i-1] + Y[i-1], 0)
11 }
12 cat("S is",S)
13 cl = B*sig;
14 cat(cl)
15 answer =100;

```

```
16 for ( i in 1:9){  
17   if(S[i]>c1){  
18     answer = i  
19   }  
20 }  
21 cat("The mean has increased after observing the ",  
      answer-1," subgroup average")
```

Chapter 14

Life Testing

R code Exa 14.3.a Lifetime of a transistor

```
1 total =50;
2 failure = 15;
3 alpha = 0.05;
4 t =525;
5 val1 = qchisq(alpha/2,2*failure)
6 val2 = qchisq(1-alpha/2,2*failure)
7
8 int1 = 2*t/val1;
9 int2 = 2*t/val2;
10 cat("The 95% confidence interval is",int2,"to",int1)
11
12 'The confidence interval is from 22.35 to 62.17
   whereas solution in R is 22.35 to 62.53
13 because of the difference in the value of chi-square
   (0.975, 30).
14 The textbook says the value is 16.89 whereas R
   calculates its value as 16.79 '
```

R code Exa 14.3.b Lifetime of Battery

```

1 t = 1800;
2 theta = 150;
3 r = 20;
4 pvalue = 1 - pchisq(2*r, 2*t/theta)
5 cat("P-value is ", pvalue)

```

R code Exa 14.3.c One at a time sequential test

```

1 T = 500;
2 alpha = 0.05;
3 r = 10;
4 val1 = qchisq(1-alpha/2, 2*r)
5 val2 = qchisq(alpha/2, 2*r)
6 int1 = 2*T/val1;
7 int2 = 2*T/val2;
8 cat("The 95% confidence interval is", int1, "to", int2)
9
10 'The confidence interval is from 29.27 to 103.52
    whereas solution in R is 29.265774 to
11 104.26683 because of the difference in the value of
    chi-square(0.975, 30).
12 The textbook says the value is 9.66 whereas scilab
    calculates its value as 9.5907774 '

```

R code Exa 14.3.d Lifetime of semiconductors

```

1 r = 30;
2 T = 600;
3 theta = 25;
4 val1 = pchisq(2*T/theta, 2*r)
5 val2 = 1 - pchisq(2*T/theta, 2*(r+1))
6 val2
7 pvalue = min(val1, val2);

```

```
8 cat("The pvalue is",pvalue)
9 cat("H0 would be accepted when the significance
    level is 0.10")
```

R code Exa 14.3.e Bayes estimator

```
1 X<-c(5, 7, 6.2, 8.1, 7.9, 15, 18, 3.9, 4.6, 5.8)
2 Y<-c(3, 3.2, 4.1, 1.8, 1.6, 2.7, 1.2, 5.4, 10.3,
    1.5)
3 t = sum(X)+sum(Y);
4 R =10;
5 a = 20;
6 b = 2;
7 estimate = (R+b)/(a+t);
8 cat("Bayes estimate of lambda is",estimate)
```

R code Exa 14.4.a Lifetime of items produced by two plants

```
1 Xlife = 420;
2 Ylife = 510;
3 Xnum= 10;
4 Ynum =15;
5 ts = Xlife*Ynum/(Ylife*Xnum);
6 cat("The value of the test statistic is",ts)
7 val = pf(ts, Xnum, Ynum)
8 pvalue = 2*(1-val);
9 cat("The p-value is",pvalue)
10 cat("We cannot reject H0");
```
