Standard Reduction Potentials of Half-Cells

(Ionic concentrations are at 1M in water @ 25° C)

•	sure	ii 1141 in water & 25°C)		
Oxidizing Agents		Reducing Agents	E ⁰ (Volts)	
$F_2(g) + 2e^{-}$	\rightarrow	2F ⁻ (aq)	+2.87	
$PbO_2(s) + SO_4^{2-}(aq) + 4H^+(aq) + 2e^-$		$PbSO_4(s) + 2H_2O(\ell)$	+1.69	
$MnO_4^-(aq) + 8H^+(aq) + 5e^-$	\rightarrow	$Mn^{2+}(aq) + 4H_2O(\ell)$	+1.51	
$Au^{3+}(aq) + 3e^{-}$		Au(s)	+1.50	
$ClO_4^-(aq) + 8H^+(aq) + 8e^-$	\rightarrow	$Cl^{-}(aq) + 4H_2O(\ell)$	+1.39	
$\text{Cl}_2(g) + 2e^-$	\rightarrow	2Cl ⁻ (aq)	+1.36	
$Cr_2O_7^{2-}(aq) + 14H^+(aq) + 6e^-$	\rightarrow	$2Cr^{3+}(aq) + 7H_2O(\ell)$	+1.33	
$2HNO_2(aq) + 4H^+(aq) + 4e^-$	\rightarrow	$N_2O(g) + 3H_2O(\ell)$	+1.30	
$O_2(g) + 4H^+(aq) + 4e^-$	\rightarrow	$2H_2O(\ell)$	+1.23	
$MnO_2(s) + 4H^+(aq) + 2e^-$	\rightarrow	$Mn^{2+}(aq) + 2H_2O(\ell)$	+1.22	
$Br_2(aq) + 2e^{-}$	\rightarrow	2Br ⁻ (aq)	+1.07	
$Hg^{2+}(aq) + 2e^{-}$	\rightarrow	$Hg(\ell)$	+0.85	
$ClO^{-}(aq) + H_2O(\ell) + 2e^{-}$	\rightarrow	$Cl^{-}(aq) + 2OH^{-}(aq)$	+0.84	
$Ag^{+}(aq) + e^{-}$		Ag(s)	+0.80	
$NO_3^-(aq) + 2H^+(aq) + e^-$	\rightarrow	$NO_2(g) + H_2O(\ell)$	+0.80	
$Fe^{3+}(aq) + e^{-}$		$Fe^{2+}(aq)$	+0.77	?
$O_2(g) + 2H^+(aq) + 2e^-$		$H_2O_2(\ell)$	+0.70	7
$I_2(s) + 2e^{-s}$	\rightarrow	2I ⁻ (aq)	+0.54	(
$O_2(g) + 2H_2O(\ell) + 4e^{-1}$	\rightarrow	4OH-(aq)	+0.40	~
$Cu^{2+}(aq) + 2e^{-}$	\rightarrow	Cu(s)	+0.34	Į
$SO_4^{2-}(aq) + 4H^+(aq) + 2e^-$	\rightarrow	H_2SO_3 (aq) + $H_2O(\ell)$	+0.17	•
$Sn^{4+}(aq) + 2e^{-}$		$\operatorname{Sn}^{2+}(\operatorname{aq})$	+0.15	(
$S(s) + 2H^{+}(aq) + 2e^{-}$	\rightarrow	H_2S (aq)	+0.14	-
$AgBr(s) + e^{-}$	\rightarrow	Ag(s) + Br(aq)	+0.07	~
$2H^{+}(aq) + 2e^{-}$	\rightarrow	$H_{2(g)}$	0.00	J
$Pb^{2+}(aq) + 2e^{-}$	\rightarrow	Pb(s)	-0.13	(
$Sn^{2+}(aq) + 2e^{-}$	\rightarrow	Sn(s)	-0.14	-
$AgI(s) + e^{-}$	\rightarrow	Ag(s) + I(aq)	-0.15	- {
$Ni^{2+}(aq) + 2e^{-}$	\rightarrow	Ni(s)	-0.26	,
$Co^{2+}(aq) + 2e^{-}$	\rightarrow	Co(s)	-0.28	4.5
$PbSO_4(s) + 2e^{-s}$	\rightarrow	$Pb(s) + SO_4^{2-}(aq)$	-0.36	7
$Se(s) + 2H^{+}(aq) + 2e^{-}$	\rightarrow	H_2 Se (aq)	-0.40	- {
$Cd^{2+}(aq) + 2e^{-}$	\rightarrow	Cd(s)	-0.40	• ;
$\operatorname{Cr}^{3+}(\operatorname{aq}) + \operatorname{e}^{-}$	\rightarrow	$Cr^{2+}(aq)$	-0.41	(
$Fe^{2+}(aq) + 2e^{-}$	\rightarrow	Fe(s)	-0.45	
$NO_2^-(aq) + H_2O(\ell) + e^-$	\rightarrow	$NO(g) + 2OH^{-}(aq)$	-0.46	-
$Ag_2S(s) + 2e^{-s}$	\rightarrow	$2Ag(s) + S^{2}(aq)$	-0.69	–
$Zn^{2+}(aq) + 2e^{-}$	\rightarrow	Zn(s)	-0.76	
$2H_2O(\ell) + 2e^{-1}$	\rightarrow	$H_2(g) + 2OH^-(aq)$	-0.83	
<u>-</u>				
$Cr^{2+}(aq) + 2e^{-}$	\rightarrow	Cr(s)	-0.91	
$Se(s) + 2e^{-s}$	\rightarrow	Se ² -(aq)	-0.92	
$SO_4^{2-}(aq) + H_2O(\ell) + 2e^{-\ell}$	\rightarrow	$SO_3^{2-}(aq) + 2OH^{-}(aq)$	-0.93	
$A1^{3+}(aq) + 3e^{-}$	\rightarrow	Al(s)	-1.66	
$Mg^{2+}(aq) + 2e^{-}$		Mg(s)	-2.37	
$Na^+(aq) + e^-$		Na(s)	-2.71	
$Ca^{2+}(aq) + 2e^{-}$		Ca(s)	-2.87	1
$Ba^{2+}(aq) + 2e^{-}$	\rightarrow	Ba(s)	-2.91	
$Li^+(aq) + e^-$	\rightarrow	Li(s)	-3.04	