- Assessment of drought impact over land cover in continental Chile by the analysis of water supply and demand from the ERA5-Land and MODIS
  - Francisco Zambrano<sup>a,1,\*</sup>

<sup>a</sup> Universidad Mayor, Hémera Centro de Observación de la Tierra, Facultad de Ciencias, Escuela de Ingeniería en Medio Ambiente y Sustentabilidad, Santiago, Chile, 7500994

#### 4 Abstract

Human-induced greenhouse gas emissions have increased the frequency and/or intensity of weather and climate extremes. Central Chile has been affected by a persistent drought which is impacting the hydrological system and vegetation development. The region has been the focus of research studies due to the diminishing water supply, this persistent period of water scarcity has been defined as a "mega drought". Nevertheless, our results evidence that the water deficit has expanded beyond. Our goal is to analyze the impact of drought, measured by drought indices of water supply/demand and vegetation status, in the LULCC (land use land cover change) over continental Chile. For the analysis, continental Chile was divided into five zones according to a latitudinal gradient: "Norte Grande", "Norte Chico", "Zona Central", "Zona Sur", and "Zona Austral". We used monthly climatic re-analysis variables for precipitation, temperature and soil moisture for 1981-2023 from ERA5-Land; and MODIS (Moderate-Resolution Imaging Spectroradiometer) product MCD12Q1 for land cover for 2001-2021, and the NDVI vegetation index from product MOD13A2 collection 6.1 for 2000-2023, both from collection 6.1. We estimated atmospheric evaporative demand (AED) by combining the Hargreaves-Samani equation with the ERA5-Land temperature. We derived the drought indices SPI (Standardized Precipitation Index), SPEI (Standardized Precipitation Evapotranspiration Index), EDDI (Evaporative Demand Drought Index), zcSM (standardized anomaly of cumulative soil moisture), and the zcNDVI (standardized anomaly of cumulative NDVI). These indices were calculated for time scales of 1, 3, 6, 12, 24, and 36 months, except for zcNDVI (1, 3, and 6 months). We analyzed the temporal correlation of SPI, SPEI, EDDI, and zcSM with zcNDVI to have insights into the impact of water supply and demand on vegetation. Our results showed that LULCC had an increasing trend of 412 [km2yr-1] of forest expansion in the "Zona Sur", together with a decreasing trend of 24 [km2yr-1] of cropland contraction in the "Zona Central" meanwhile the "Zona Sur" showed an increase of 31 [km2yr-1], and a contraction of 80 [km2yr-1] of bare soil in the "Zona Austral". The EDDI was the less correlated index for the five macro zones and the five types of land cover, showing that the temperature in Chile has little impact on vegetation. Higher r-squared values, between 0.5 and 0.8, were obtained at "Norte Chico" and "Zona Central" for the land cover types of savanna, shrubland, grassland, and croplands for the indices SPEI and zcSM at time scales of 12 and 24 months. The forest type reaches a r-squared of ~0.5 for zcSM of 12 months. The results indicate that the "Norte Chico" and "Zona Central" are the most sensitive regions to water supply deficits longer than a year, potentially explained by a low capacity of water storage in those zones that should be further investigated.

5 Keywords: drought, land cover change, satellite

Email address: francisco.zambrano@umayor.cl (Francisco Zambrano)

<sup>\*</sup>Corresponding author

<sup>&</sup>lt;sup>1</sup>This is the first author footnote.

### 1. Introduction

10

12

13

15

16

17

19

20

21

23

24

25

27

28

31

32

34

37

39

41

42

47

50

51

52

53

The sixth assessment report (AR6) of the IPCC [1] indicates that human-induced greenhouse gas emissions have increased the frequency and/or intensity of some weather and climate extremes, and the evidence has been strengthened since AR5 [2]. There is high confidence that increasing global warming can expand the land area affected by increasing drought frequency and severity [3]. Furthermore, drought increases tree mortality and triggers changes in land cover and, consequently, land use, thus impacting ecosystems [4]. Nevertheless, there is a lack of understanding of how the alteration in water supply and demand is affecting land cover transformations.

Precipitation is the primary driver of drought and is intensified by temperature [5]. Drought impacts soil moisture, hydrological regimes, and vegetation productivity. Initially, drought was commonly classified as meteorological, hydrological, and agricultural [6]. Lately, [7] and [8] have given an updated definition of drought for the Anthropocene, suggesting that it should be considered the feedback of humans' decisions and activities that drives the anthropogenic drought. Even though it has been argued that those definitions do not fully address the ecological dimensions of drought. [4] proposed the ecological drought definition as "an episodic deficit in water availability that drives ecosystems beyond thresholds of vulnerability, impacts ecosystem services, and triggers feedback in natural and/or human systems". Moreover, many ecological studies have misinterpreted how to characterize drought, for example, sometimes considering "dry" conditions as "drought" [9]. On the other hand, the AR6 [1] states that even if global warming is stabilized at 1.5°-2°C, many parts of the world will be impacted by more severe agricultural and ecological droughts. Then, there is a challenge in conducting drought research, especially to evaluate its impact on ecosystems.

Chile has been facing a persistent rainfall deficit for more than a decade [10], which has impacted vegetation development [11] and the hydrological system [12]. Current drought conditions have affected crop productivity [13, 14], forest development [15, 16], forest fire occurrence [17], land cover change [18], water supply in watersheds [19], and have had economic impacts [20]. In 2019–2020, the drought severity reached an extreme condition in Central Chile (30–34°S) not seen for at least 40 years, and the evidence indicates that the impact is transversal to the land cover classes of forest, grassland, and cropland [11]. The prolonged lack of precipitation in Central Chile is producing changes in ecosystem dynamics that must be studied.

For the spatiotemporal assessment of drought impact (i.e., by water supply and demand) on land cover changes, we need climatic realiable variables such as precipitation, temperature, soil moisture, land cover, and vegetation status. For developing countries like Chile, the weather networks present several disadvantages, such as gaps, a short history, and low-quality data. Reanalysis data, as the ERA5-Land (ERA5L) [21] provides hourly climatic information (precipitation, temperature, and soil moisture) without gaps since 1950 with global extension. ERA5L has already been used for drought assessment using the Standardized Precipitation-Evapotranspiration Index (SPEI) [22] and for flash drought [23] by analyzing soil mositure and evapotranspiration. On the other hand, satellite remote sensing [24, 25] is the primary method to evaluate how drought impacts vegetation dynamics. Vegetation drought indices (VDI) are commonly used as proxies of productivity [26, 27], which can be derived from the MODIS (Moderate-Resolution Imaging Spectroradiometer). Besides, land use and land cover (LULC) change can be driven by drought [28, 29]. To analyze these changes, multiple LULC products exist [30], one of those that provides time series since 2001 is the MCD12Q1 [31] from MODIS. The variation in water supply and demand is finally reflected in the total water storage (TWS). The TWS can be retrieved by the Gravity Recovery and Climate Experiment (GRACE), which allows analyzing water availability changes at coarse resolution [32, 33]. We can use climatic reanalysis (ERA5L) and vegetation data (MODIS) to derive drought indices of supply (i.e., precipitation) and demand (i.e., temperature) and thus evaluate the impact of drought on LULC changes. Further, the TWS can be assessed with regard to the changes in water supply and demand to gain insight into the impact on water storage.

To evaluate meteorological drought (i.e., water supply), the World Meteorological Organization (WMO; [34]) recommends the Standardized Precipitation Index (SPI; [35]), a multiscalar drought index that allows to monitor precipitation deficits from short- to long-term. Following the same approach, [36] incorporates

into the SPI the effect of temperature through the use of potential evapotranspiration, thus proposing the SPEI (Standardized Precipitation Evapotranspiration Index). Similarly, to evaluate solely the evaporative demand driven by temperature, [37] and [38] came up with the Evaporative Demand Drought Index (EDDI). For vegetation, in a similar manner as the SPI, SPEI and EDDI; [14] proposed the zcNDVI, a standardized anomaly of the cumulative Normalized Difference Vegetation Index (NDVI), which could be acumulated over the growing season or any period (e.g., months), resulting in a multiscalar drought index. For soil moisture, several drought indices exist, such as the Soil Moisture Deficit Index (SDMI) a normalized index [39] and the Soil Moisture Agricultural Drought Index (SMADI) [40] which is a normalized index using vegetation, land surface temperature, and a vegetation condition index (VCI, [41]). From TWS, we can estimate the standardized terrestrial water storage index (STI) [42], a standardized anomaly that follows the methodology of the SPI, SPEI, EDDI, and zcNDVI. Thereby, we have drought indices for water supply, demand, and storage, which can help to make a comprehensive assessment of drought.

In this research, we aim to analyze the impact of drought on different types of land cover classes in continental Chile by combining environmental variables such as biomass productivity, and water demand and supply gathered from earth observation products. The specific objectives of the study are: i) to calculate multi-scalar drought indices for water demand and supply for 1981–2023; ii) to evaluate LULC change for 2001–2021 and its relation to drought indices; iii) to analyze the relationship of a proxy of biomass (zcNDVI) with drought indices; and iv) to assess if the observed changes in the drought indices are linked to the TWS.

# 2. Study area

### 3. Materials and Methods

75 3.1. Data

69

70

71

- 76 3.1.1. Earth observation data
- 77 3.1.2. in-situ data
- 78 3.2. Drought indices for water demand and supply
- 3.3. Analysis of a biomass proxy with drought indices of supply and demand
- 3.4. LULC change for 2001-2021 and its relation with water supply and demand
- <sup>81</sup> ![Proportion of land cover class from the persistent land cover for 2001-2021 (>80%) per macrozone.

# 3.5. Total water storage and drought indices

Terrestrial Water Storage (TWS) is defined as the total amount of water stored on land that includes any natural or artificial water bodies, such as groundwater, soil moisture, rivers, lakes, snowpack, ice, and biomass water [43, 44]. TWS changes evidence the effects of multiple water fluxes on the hydrological cycle [44]. These are reflected in the temporal variation of observations of the Earth's gravitational field [45, 46] and, in recent decades, the twin Gravity Recovery and Climate Experiment (GRACE) satellites and their Follow-On mission (GRACE-FO) have provided valuable results on globally distributed TWS anomalies [47, 48]. The GRACE mission was launched in March 2002 and was operational until October 2017 [49]. Then its GRACE Follow-On successor was launched in May 2018 [50, 51]. The information provided by these satellites is used to construct monthly maps of the Earth's average gravity field, providing details of the movement of water masses or water mass anomaly estimates relative to the long-term average gravity field [43, 52].

In this study, RL06.1\_V3 GRACE mascon (mass concentration) monthly solutions were used, which are provided by the Jet Propulsion Laboratory (JPL-M, https://grace.jpl.nasa.gov). Each GRACE Tellus monthly grid at 0.5 degrees represents the deviation of surface mass for that month relative to a reference time average (2004-2009 baseline), which is subtracted from all other monthly grids to provide terrestrial water storage anomalies (TWSA) [49, 51]. This JPL-M version of the data employs a Coastal Resolution Improvement (CRI) filter that reduces signal leakage errors across coastlines [? ? ]. Although the mascon





Figure 1: (a) Chile and the five zones "norte grande", "norte chico", "zona central", "zona sur", and "zona austral". (b) Topography reference map. (c) Land cover classes for 2021. (d) Persistent land cover classes (>80%) for 2001-2021.

solutions greatly reduced leakage errors, a gain factor, which is used to enhance the spatial resolution, was applied to the dataset as recommended for hydrological studies [49, 51]. The water storage and height anomalies are given in Equivalent Water Height or Thickness units (EWH, cm) [46], and their temporal resolution monthly is from April 2002 to May 2023.



Trend of change [km² year-1]												
zone norte grande	Forest		Cropland		Grassland		Savanna		Shrubland		Barren land	
					-31.2	<u> </u>			-2.3	~~.	32.8	
norte chico					-100.2	^	-66.1	·	79.3		104.0	· ~ ~
zona central	-66.2	<u> </u>	-24.4	·~~.	89.9		-128.6	<u>~~</u>	130.2		23.3	~~~
zona sur	412.4		30.8	~_·	-55.9	<u></u>	-316.2	<u>~</u>	-14.6	·	2.1	~~~
zona austral	-9.1	~~~			226.1	_	163.9				-80.2	~~~

Figure 2: Proportion of land cover class from the persistent land cover for 2001-2021 (>80%) per macrozone. The table on the left shows the linear change trend next to time-series plot of surface, per land cover class (IGBP MCD12Q1.006) for 2001-2021 through the fives zones on continental Chile. Blue dots on the plots indicate the maximum and red dots minimum surface reach.

## 3.6. Validation of ERA5-Land variables

### 4. Results

105

113

114

115

116

117

119

120

121

122

123

124

125

126

127

128

129 130

131

132

133

134

135

136

137

138

139

- 106 4.1. Drought indices for water demand and supply
- 107 4.2. Biomass proxy with drought indices of supply and demand
- 108 4.3. LULC change for 2001-2021 and its relation with water supply and demand
- 109 4.4. Total water storage (TWS) and drought indices
- 110 4.5. Validation of ERA5-Land variables

### 5. Discussion

## 112 6. Conclusion

# References

- [1] K. Calvin, D. Dasgupta, G. Krinner, A. Mukherji, P. W. Thorne, C. Trisos, J. Romero, P. Aldunce, K. Barrett, G. Blanco, W. W. Cheung, S. Connors, F. Denton, A. Diongue-Niang, D. Dodman, M. Garschagen, O. Geden, B. Hayward, C. Jones, F. Jotzo, T. Krug, R. Lasco, Y.-Y. Lee, V. Masson-Delmotte, M. Meinshausen, K. Mintenbeck, A. Mokssit, F. E. Otto, M. Pathak, A. Pirani, E. Poloczanska, H.-O. Pörtner, A. Revi, D. C. Roberts, J. Roy, A. C. Ruane, J. Skea, P. R. Shukla, R. Slade, A. Slangen, Y. Sokona, A. A. Sörensson, M. Tignor, D. Van Vuuren, Y.-M. Wei, H. Winkler, P. Zhai, Z. Zommers, J.-C. Hourcade, F. X. Johnson, S. Pachauri, N. P. Simpson, C. Singh, A. Thomas, E. Totin, P. Arias, M. Bustamante, I. Elgizouli, G. Flato, M. Howden, C. Méndez-Vallejo, J. J. Pereira, R. Pichs-Madruga, S. K. Rose, Y. Saheb, R. Sánchez Rodríguez, D. Ürge Vorsatz, C. Xiao, N. Yassaa, A. Alegría, K. Armour, B. Bednar-Friedl, K. Blok, G. Cissé, F. Dentener, S. Eriksen, E. Fischer, G. Garner, C. Guivarch, M. Haasnoot, G. Hansen, M. Hauser, E. Hawkins, T. Hermans, R. Kopp, N. Leprince-Ringuet, J. Lewis, D. Ley, C. Ludden, L. Niamir, Z. Nicholls, S. Some, S. Szopa, B. Trewin, K.-I. Van Der Wijst, G. Winter, M. Witting, A. Birt, M. Ha, J. Romero, J. Kim, E. F. Haites, Y. Jung, R. Stavins, A. Birt, M. Ha, D. J. A. Orendain, L. Ignon, S. Park, Y. Park, A. Reisinger, D. Cammaramo, A. Fischlin, J. S. Fuglestvedt, G. Hansen, C. Ludden, V. Masson-Delmotte, J. R. Matthews, K. Mintenbeck, A. Pirani, E. Poloczanska, N. Leprince-Ringuet, C. Péan, IPCC, 2023: Climate Change 2023: Synthesis Report. Contribution of Working Groups I, II and III to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change [Core Writing Team, H. Lee and J. Romero (eds.)]. IPCC, Geneva, Switzerland., Tech. rep., Intergovernmental Panel on Climate Change (IPCC) (Jul.
  - URL https://www.ipcc.ch/report/ar6/syr/
- [2] IPCC, Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change, Cambridge University Press, Cambridge, UK; New York, USA, 2013. doi:10.1017/CB09781107415324.
  - URL www.climatechange2013.org
- [3] X. a. A. M. a. B. W. a. D. C. a. L. A. a. G. S. a. I. I. a. K. J. a. L. S. a. O. F. a. P. I. a. S. M. a. V.-S. S. a. W. M. a. Z. M.-D. B. a. V. a. O. Seneviratne, S and Zhang, Weather and Climate Extreme Events in a Changing Climate, Cambridge University Press. In Press., 2021, publication Title: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change.

[4] S. D. Crausbay, A. R. Ramirez, S. L. Carter, M. S. Cross, K. R. Hall, D. J. Bathke, J. L. Betancourt, S. Colt, A. E. Cravens, M. S. Dalton, J. B. Dunham, L. E. Hay, M. J. Hayes, J. McEvoy, C. A. McNutt, M. A. Moritz, K. H. Nislow, N. Raheem, T. Sanford, Defining Ecological Drought for the Twenty-First Century, Bulletin of the American Meteorological Society 98 (12) (2017) 2543-2550, publisher: American Meteorological Society. doi:10.1175/BAMS-D-16-0292.1. URL https://journals.ametsoc.org/view/journals/bams/98/12/bams-d-16-0292.1.xml

140

141

142

143 144

145

146

147

151

152

154

155

156 157

158

- L. Luo, D. Apps, S. Arcand, H. Xu, M. Pan, M. Hoerling, Contribution of temperature and precipitation anomalies to the California drought during 2012–2015, Geophysical Research Letters 44 (7) (2017) 3184–3192. doi:10.1002/2016GL072027. URL https://agupubs.onlinelibrary.wiley.com/doi/10.1002/2016GL072027
- [6] D. A. Wilhite, M. H. Glantz, Understanding: The drought phenomenon: The role of definitions, Water International 10 (3) 148 (1985) 111-120. doi:10.1080/02508068508686328. 149 URL http://dx.doi.org/10.1080/02508068508686328 150
- A. F. Van Loon, T. Gleeson, J. Clark, A. I. Van Dijk, K. Stahl, J. Hannaford, G. Di Baldassarre, A. J. Teuling, L. M. Tallaksen, R. Uijlenhoet, D. M. Hannah, J. Sheffield, M. Svoboda, B. Verbeiren, T. Wagener, S. Rangecroft, N. Wanders, H. A. Van Lanen, Drought in the Anthropocene, Nature Geoscience 9 (2) (2016) 89-91. doi:10.1038/ngeo2646. 153
  - A. AghaKouchak, A. Mirchi, K. Madani, G. Di Baldassarre, A. Nazemi, A. Alborzi, H. Anjileli, M. Azarderakhsh, F. Chiang, E. Hassanzadeh, L. S. Huning, I. Mallakpour, A. Martinez, O. Mazdiyasni, H. Moftakhari, H. Norouzi, M. Sadegh, D. Sadeqi, A. F. Van Loon, N. Wanders, Anthropogenic Drought: Definition, Challenges, and Opportunities, Reviews of Geophysics 59 (2) (2021) e2019RG000683. doi:10.1029/2019RG000683.  $URL\ https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2019RG000683$
- I. J. Slette, A. K. Post, M. Awad, T. Even, A. Punzalan, S. Williams, M. D. Smith, A. K. Knapp, How ecologists define 159 drought, and why we should do better, Global Change Biology 25 (10) (2019) 3193-3200. doi:10.1111/gcb.14747. 160 URL https://onlinelibrary.wiley.com/doi/10.1111/gcb.14747 161
- R. Garreaud, C. Alvarez-Garreton, J. Barichivich, J. P. Boisier, D. Christie, M. Galleguillos, C. LeQuesne, J. McPhee, 162 163 M. Zambrano-Bigiarini, The 2010-2015 mega drought in Central Chile: Impacts on regional hydroclimate and vegetation, Hydrology and Earth System Sciences Discussions 2017 (2017) 1-37. doi:10.5194/hess-2017-191. 164 URL http://www.hydrol-earth-syst-sci-discuss.net/hess-2017-191/ 165
- F. Zambrano, Four decades of satellite data for agricultural drought monitoring throughout the growing season in Central 166 167 Chile, in: R. M. Vijay P. Singh Deepak Jhajharia, R. Kumar (Eds.), Integrated Drought Management, Two Volume Set, CRC Press, 2023, p. 28.
- J. P. Boisier, C. Alvarez-Garreton, R. R. Cordero, A. Damiani, L. Gallardo, R. D. Garreaud, F. Lambert, C. Ramallo, 169 M. Rojas, R. Rondanelli, Anthropogenic drying in central-southern Chile evidenced by long-term observations and climate 170 model simulations, Elementa 6 (1) (2018) 74. doi:10.1525/elementa.328. URL https://www.elementascience.org/article/10.1525/elementa.328/ 172
- [13] F. Zambrano, M. Lillo-Saavedra, K. Verbist, O. Lagos, Sixteen years of agricultural drought assessment of the biobío 173 region in chile using a 250 m resolution vegetation condition index (VCI), Remote Sensing 8 (6) (2016) 1-20, publisher: 174 Multidisciplinary Digital Publishing Institute. doi:10.3390/rs8060530. 175 176 URL http://www.mdpi.com/2072-4292/8/6/530
- [14] F. Zambrano, A. Vrieling, A. Nelson, M. Meroni, T. Tadesse, Prediction of drought-induced reduction of agricultural 177 178 productivity in Chile from MODIS, rainfall estimates, and climate oscillation indices, Remote Sensing of Environment 219 (2018) 15-30, publisher: Elsevier. doi:10.1016/j.rse.2018.10.006. 179 URL https://www.sciencedirect.com/science/article/pii/S0034425718304541 180
- A. Miranda, A. Lara, A. Altamirano, C. Di Bella, M. E. González, J. Julio Camarero, Forest browning trends in response 181 to drought in a highly threatened mediterranean landscape of South America, Ecological Indicators 115 (2020) 106401. 182 doi:10.1016/i.ecolind.2020.106401. 183 URL https://linkinghub.elsevier.com/retrieve/pii/S1470160X20303381
- A. Venegas-González, F. R. Juñent, A. G. Gutiérrez, M. T. Filho, Recent radial growth decline in response to increased 185 drought conditions in the northernmost Nothofagus populations from South America, Forest Ecology and Management 186  $409\ (2018)\ 94-104.\ {\tt doi:10.1016/j.foreco.2017.11.006}.$ 187 URL https://linkinghub.elsevier.com/retrieve/pii/S0378112717313993 188
- R. Urrutia-Jalabert, M. E. González, . González-Reyes, A. Lara, R. Garreaud, Climate variability and forest fires in 189 central and south-central Chile, Ecosphere 9 (4) (2018) e02171. doi:10.1002/ecs2.2171. 190 URL https://esajournals.onlinelibrary.wiley.com/doi/10.1002/ecs2.2171 191
- [18] I. Fuentes, R. Fuster, D. Avilés, W. Vervoort, Water scarcity in central Chile: the effect of climate and land cover changes 192 on hydrologic resources, Hydrological Sciences Journal 66 (6) (2021) 1028-1044. doi:10.1080/02626667.2021.1903475. 193 URL https://www.tandfonline.com/doi/full/10.1080/02626667.2021.1903475 194
- C. Alvarez-Garreton, J. P. Boisier, R. Garreaud, J. Seibert, M. Vis, Progressive water deficits during multiyear droughts 195 in basins with long hydrological memory in Chile, Hydrology and Earth System Sciences 25 (1) (2021) 429-446. doi: 196 10.5194/hess-25-429-2021. 197 URL https://hess.copernicus.org/articles/25/429/2021/ 198
- [20] F. J. Fernández, F. Vásquez-Lavín, R. D. Ponce, R. Garreaud, F. Hernández, O. Link, F. Zambrano, M. Hanemann, The 199 economics impacts of long-run droughts: Challenges, gaps, and way forward, Journal of Environmental Management 344 200 (2023) 118726. doi:10.1016/j.jenvman.2023.118726. 201  $URL\ https://linkinghub.elsevier.com/retrieve/pii/S0301479723015141$ 202
- 203 J. Muñoz-Sabater, E. Dutra, A. Agustí-Panareda, C. Albergel, G. Arduini, G. Balsamo, S. Boussetta, M. Choulga, S. Harrigan, H. Hersbach, B. Martens, D. G. Miralles, M. Piles, N. J. Rodríguez-Fernández, E. Zsoter, C. Buontempo, 204

- J.-N. Thépaut, ERA5-Land: a state-of-the-art global reanalysis dataset for land applications, Earth System Science Data 13 (9) (2021) 4349–4383. doi:10.5194/essd-13-4349-2021.

  URL https://essd.copernicus.org/articles/13/4349/2021/
- [22] M. Nouri, Drought Assessment Using Gridded Data Sources in Data-Poor Areas with Different Aridity Conditions, Water
   Resources Management 37 (11) (2023) 4327-4343. doi:10.1007/s11269-023-03555-4.
   URL https://link.springer.com/10.1007/s11269-023-03555-4
- 211 [23] M. Wang, L. Menzel, S. Jiang, L. Ren, C.-Y. Xu, H. Cui, Evaluation of flash drought under the impact of heat wave events 212 in southwestern Germany, Science of The Total Environment 904 (2023) 166815. doi:10.1016/j.scitotenv.2023.166815. 213 URL https://linkinghub.elsevier.com/retrieve/pii/S0048969723054402
- [24] H. West, N. Quinn, M. Horswell, Remote sensing for drought monitoring \& impact assessment: Progress, past challenges and future opportunities, Remote Sensing of Environment 232, publisher: Elsevier Inc. (Oct. 2019). doi:10.1016/j.rse. 2019.111291.
- [25] A. AghaKouchak, A. Farahmand, F. S. Melton, J. Teixeira, M. C. Anderson, B. D. Wardlow, C. R. Hain, Remote sensing of drought: Progress, challenges and opportunities, Reviews of Geophysics 53 (2) (2015) 452–480. doi:10.1002/2014RG00 0456.
   URL http://dx.doi.org/10.1002/2014RG000456.
- 221 [26] J. M. Paruelo, M. Texeira, L. Staiano, M. Mastrángelo, L. Amdan, F. Gallego, An integrative index of Ecosystem Services provision based on remotely sensed data, Ecological Indicators 71 (2016) 145–154, publisher: Elsevier. doi:10.1016/J.EC 0LIND.2016.06.054.
  - URL https://www.sciencedirect.com/science/article/pii/S1470160X16303843

224

- [27] A. Schucknecht, M. Meroni, F. Kayitakire, A. Boureima, A. Schucknecht, M. Meroni, F. Kayitakire, A. Boureima,
   Phenology-Based Biomass Estimation to Support Rangeland Management in Semi-Arid Environments, Remote Sensing 9 (5) (2017) 463, publisher: Multidisciplinary Digital Publishing Institute. doi:10.3390/rs9050463.
   URL http://www.mdpi.com/2072-4292/9/5/463
- 229 [28] H. T. Tran, J. B. Campbell, R. H. Wynne, Y. Shao, S. V. Phan, Drought and Human Impacts on Land Use and Land
  230 Cover Change in a Vietnamese Coastal Area, Remote Sensing 2019, Vol. 11, Page 333 11 (3) (2019) 333, publisher:
  231 Multidisciplinary Digital Publishing Institute. doi:10.3390/RS11030333.
  232 URL https://www.mdpi.com/2072-4292/11/3/333/htm
- [29] F. O. Akinyemi, Vegetation Trends, Drought Severity and Land Use-Land Cover Change during the Growing Season in Semi-Arid Contexts, Remote Sensing 2021, Vol. 13, Page 836 13 (5) (2021) 836, publisher: Multidisciplinary Digital Publishing Institute. doi:10.3390/RS13050836.
   URL https://www.mdpi.com/2072-4292/13/5/836/htm
- [30] G. Grekousis, G. Mountrakis, M. Kavouras, An overview of 21 global and 43 regional land-cover mapping products,
   International Journal of Remote Sensing 36 (21) (2015) 5309-5335. doi:10.1080/01431161.2015.1093195.
   URL https://www.tandfonline.com/doi/full/10.1080/01431161.2015.1093195
- [31] M. Friedl, D. Sulla-Menashe, MCD12Q1 MODIS/Terra+Aqua Land Cover Type Yearly L3 Global 500m SIN Grid V006
   [Data set]. NASA EOSDIS Land Processes DAAC (2019). doi:10.5067/M0DIS/MCD12Q1.006.
- [32] M. Ahmed, M. Sultan, J. Wahr, E. Yan, The use of GRACE data to monitor natural and anthropogenic induced variations in water availability across Africa, Earth-Science Reviews 136 (2014) 289–300, publisher: Elsevier. doi:10.1016/J.EARS
   CIREV.2014.05.009.
- [33] S. Ma, Q. Wu, J. Wang, S. Zhang, Temporal Evolution of Regional Drought Detected from GRACE TWSA and CCI
   SM in Yunnan Province, China, Remote Sensing 2017, Vol. 9, Page 1124 9 (11) (2017) 1124, publisher: Multidisciplinary
   Digital Publishing Institute. doi:10.3390/RS9111124.
   URL https://www.mdpi.com/2072-4292/9/11/1124/htm
- 249 [34] WMO, M. Svoboda, M. Hayes, D. A. Wood, Standardized Precipitation Index User Guide, WMO, Geneva, 2012, series
  250 Title: WMO Publication Title: WMO-No. 1090 © Issue: 1090.
  251 URL http://library.wmo.int/opac/index.php?lvl=notice\_display&id=13682
- <sup>252</sup> [35] T. B. Mckee, N. J. Doesken, J. Kleist, The relationship of drought frequency and duration to time scales. In: Proceedings of the Ninth Conference on Applied Climatology., American Metereological Society (Boston) (1993) 179–184.
- [36] S. M. Vicente-Serrano, S. Beguería, J. I. López-Moreno, A multiscalar drought index sensitive to global warming: The standardized precipitation evapotranspiration index, Journal of Climate 23 (7) (2010) 1696–1718. doi:10.1175/2009JCLI
   2909.1.
  - URL http://dx.doi.org/10.1175/2009JCLI2909.1
- [37] M. T. Hobbins, A. Wood, D. J. McEvoy, J. L. Huntington, C. Morton, M. Anderson, C. Hain, The Evaporative Demand
   Drought Index. Part I: Linking Drought Evolution to Variations in Evaporative Demand, Journal of Hydrometeorology
   [17] (6) (2016) 1745–1761. doi:10.1175/JHM-D-15-0121.1.
   URL http://journals.ametsoc.org/doi/10.1175/JHM-D-15-0121.1
- [38] D. J. McEvoy, J. L. Huntington, M. T. Hobbins, A. Wood, C. Morton, M. Anderson, C. Hain, The Evaporative Demand
   Drought Index. Part II: CONUS-Wide Assessment against Common Drought Indicators, Journal of Hydrometeorology
   [17] (6) (2016) 1763–1779. doi:10.1175/JHM-D-15-0122.1.
   URL http://journals.ametsoc.org/doi/10.1175/JHM-D-15-0122.1
- [39] B. Narasimhan, R. Srinivasan, Development and evaluation of Soil Moisture Deficit Index (SMDI) and Evapotranspiration
   Deficit Index (ETDI) for agricultural drought monitoring, Agricultural and Forest Meteorology 133 (1-4) (2005) 69–88.
   doi:10.1016/j.agrformet.2005.07.012.
- URL https://linkinghub.elsevier.com/retrieve/pii/S0168192305001565

- [40] A. G. S. S. Souza, A. Ribeiro Neto, L. L. D. Souza, Soil moisture-based index for agricultural drought assessment: SMADI application in Pernambuco State-Brazil, Remote Sensing of Environment 252 (2021) 112124. doi:10.1016/j.rse.2020.1
   12124.
  - URL https://linkinghub.elsevier.com/retrieve/pii/S0034425720304971

273

279

286

- <sup>274</sup> [41] F. N. Kogan, Application of vegetation index and brightness temperature for drought detection, Advances in Space Research 15 (11) (1995) 91–100. doi:10.1016/0273-1177(95)00079-T.
- [42] A. Cui, J. Li, Q. Zhou, R. Zhu, H. Liu, G. Wu, Q. Li, Use of a multiscalar GRACE-based standardized terrestrial water
   storage index for assessing global hydrological droughts, Journal of Hydrology 603 (2021) 126871. doi:10.1016/j.jhydro
   1.2021.126871.
  - $URL\ https://linkinghub.elsevier.com/retrieve/pii/S0022169421009215$
- [43] V. Humphrey, M. Rodell, A. Eicker, Using Satellite-Based Terrestrial Water Storage Data: A Review, Surveys in Geo-physics 44 (5) (2023) 1489–1517. doi:10.1007/s10712-022-09754-9.
   URL https://link.springer.com/10.1007/s10712-022-09754-9
- [44] S. Deng, Y. Liu, W. Zhang, A Comprehensive Evaluation of GRACE-Like Terrestrial Water Storage (TWS) Reconstruction
   Products at an Interannual Scale During 1981–2019, Water Resources Research 59 (3) (2023) e2022WR034381. doi:
   10.1029/2022WR034381.
  - URL https://agupubs.onlinelibrary.wilev.com/doi/10.1029/2022WR034381
- [45] R. Abolafia-Rosenzweig, M. Pan, J. Zeng, B. Livneh, Remotely sensed ensembles of the terrestrial water budget over
   major global river basins: An assessment of three closure techniques, Remote Sensing of Environment 252 (2021) 112191.
   doi:10.1016/j.rse.2020.112191.
- URL https://linkinghub.elsevier.com/retrieve/pii/S0034425720305642
- [46] F. Sabzehee, A. Amiri-Simkooei, S. Iran-Pour, B. Vishwakarma, R. Kerachian, Enhancing spatial resolution of GRACE-derived groundwater storage anomalies in Urmia catchment using machine learning downscaling methods, Journal of Environmental Management 330 (2023) 117180. doi:10.1016/j.jenvman.2022.117180.
   URL https://linkinghub.elsevier.com/retrieve/pii/S0301479722027530
- [47] B. D. Tapley, M. M. Watkins, F. Flechtner, C. Reigber, S. Bettadpur, M. Rodell, I. Sasgen, J. S. Famiglietti, F. W. Landerer, D. P. Chambers, J. T. Reager, A. S. Gardner, H. Save, E. R. Ivins, S. C. Swenson, C. Boening, C. Dahle, D. N. Wiese, H. Dobslaw, M. E. Tamisiea, I. Velicogna, Contributions of GRACE to understanding climate change, Nature Climate Change 9 (5) (2019) 358–369. doi:10.1038/s41558-019-0456-2.
   URL https://www.nature.com/articles/s41558-019-0456-2
- [48] V. Ferreira, B. Yong, H. Montecino, C. E. Ndehedehe, K. Seitz, H. Kutterer, K. Yang, Estimating GRACE terrestrial
   water storage anomaly using an improved point mass solution, Scientific Data 10 (1) (2023) 234. doi:10.1038/s41597-023-02122-1.
- 303 URL https://www.nature.com/articles/s41597-023-02122-1
- [49] M. Ramjeawon, M. Demlie, M. Toucher, Analyses of groundwater storage change using GRACE satellite data in the
   Usutu-Mhlatuze drainage region, north-eastern South Africa, Journal of Hydrology: Regional Studies 42 (2022) 101118.
   doi:10.1016/j.ejrh.2022.101118.
   URL https://linkinghub.elsevier.com/retrieve/pii/S2214581822001318
- [50] F. W. Landerer, F. M. Flechtner, H. Save, F. H. Webb, T. Bandikova, W. I. Bertiger, S. V. Bettadpur, S. H. Byun,
   C. Dahle, H. Dobslaw, E. Fahnestock, N. Harvey, Z. Kang, G. L. H. Kruizinga, B. D. Loomis, C. McCullough, M. Murböck,
   P. Nagel, M. Paik, N. Pie, S. Poole, D. Strekalov, M. E. Tamisiea, F. Wang, M. M. Watkins, H. Wen, D. N. Wiese,
   D. Yuan, Extending the Global Mass Change Data Record: GRACE Follow-On Instrument and Science Data Performance,
   Geophysical Research Letters 47 (12) (2020) e2020GL088306. doi:10.1029/2020GL088306.
   URL https://agupubs.onlinelibrary.wiley.com/doi/10.1029/2020GL088306
- 314 [51] W. Yin, S. Yang, L. Hu, S. Tian, X. Wang, R. Zhao, P. Li, Improving understanding of spatiotemporal water storage changes over China based on multiple datasets, Journal of Hydrology 612 (2022) 128098. doi:10.1016/j.jhydrol.2022 .128098.
  - URL https://linkinghub.elsevier.com/retrieve/pii/S0022169422006734
- 52] J. Wahr, S. Swenson, V. Zlotnicki, I. Velicogna, Time-variable gravity from GRACE: First results: TIME-VARIABLE GRAVITY FROM GRACE, Geophysical Research Letters 31 (11) (2004) n/a-n/a. doi:10.1029/2004GL019779.

  URL http://doi.wiley.com/10.1029/2004GL019779