Proof Rules for Termination of Loops: A Comprehensive Analysis

Krzysztof R. Apt[1], Frank S. de Boer[2], and Ernst-Rüdiger Olderog[3]

[1] CWI, Amsterdam, The Netherlands
MIMUW, University of Warsaw, Poland
apt@cwi.nl
[2] CWI, Amsterdam, The Netherlands
F.S.de.Boer@cwi.nl
[3] Carl von Ossietzky University of Oldenburg, Oldenburg, Germany
olderog@informatik.uni-oldenburg.de

Abstract. We investigate four proof rules for proving termination of **while** programs and show their proof-theoretic equivalence. This involves a proof-theoretic analysis of various auxiliary proof rules in Hoare's logic. By discussing representations of proofs in the form of proof outlines, we reveal differences between these equivalent proof rules when used in practice. We also address applications in the context of the paradigm of design by contract.

1 Introduction

In his seminal paper [**?**], Hoare introduced an axiomatic method of reasoning about correctness of **while** programs, now called Hoare's logic. It is based on correctness formulas $\{p\}\ S\ \{q\}$, where S is a program and p and q are assertions (logical formulas), with the interpretation

> "If the assertion p is true before initiation of a program S, then the assertion q will be true on its completion."

In this context p is referred to as a *precondition* and q as a *postcondition* of S.

However, in contrast to Floyd's earlier paper [**?**] that dealt with correctness of flowchart programs, program termination was not addressed. To stress this difference one distinguishes now between *partial correctness* that only focuses on the delivery of correct results, and *total correctness*, that in addition stipulates that the program terminates. So the original proposal of Hoare dealt with partial correctness.

All approaches to proving program termination within Hoare's logic formalize Floyd's [**?**] observation that

> "Proofs of termination are dealt with by showing that each step of a program decreases some entity which cannot decrease indefinitely."

The first extension of Hoare's logic to deal with total correctness is due to [**?**]. Since then substantially simpler proof rules were proposed. In these proof rules variables that range over natural numbers are used. An appropriate relative completeness result, see for example [**?**], shows that variables ranging over more general well-founded orderings are not needed.

Termination continues to be a relevant and vibrant topic in program analysis, see for example [**?**] and the Annual International Termination Competition[4]. The latter comprises various competition categories, for instance proving termination of C programs, Java bytecode programs, logic programs, functional programs, and term rewriting systems. Here we focus on the shape of termination proofs in the context of Hoare's logic. To this end, we investigate four natural proof rules from the proof-theoretic point of view. More specifically, we study **while** programs with Hoare's original proof system for program correctness, in which the well-known proof rule for partial correctness of the **while** loops, which we call the LOOP I rule, is replaced by a suitable proof rule for establishing total correctness. We analyze four versions of such a proof rule.

- The LOOP II rule involves a single, monolithic proof which establishes an invariant, termination of the body of the while loop, and which ensures that the body itself cannot be executed infinitely often by a so-called bound function.
- The LOOP III rule achieves a separation of the reasoning about the invariant and the bound function.
- The LOOP IV rule not only achieves a separation of the reasoning about the invariant and the bound function, it additionally involves a separate proof of the termination the body itself. As such it is particularly well-suited when dealing with nested loops because it modularizes the correctness proof of the outer loop from the ones of the inner loops. It is a *hybrid* proof rule with premises referring to proof systems for both partial and total correctness.
- The LOOP V rule, like the LOOP II rule, is a monolithic rule which formalizes reasoning about termination of a while loop in a single proof. However, instead of an explicit representation of the bound function it uses a parameter of the invariant itself.

Depending on the choice of the loop rule, we obtain proof systems that we refer to by II, III, IV, and V, respectively. We show that these four proof systems are equivalent in the sense that every proof of a correctness formula $\{p\}\ S\ \{q\}$ carried out in one of these systems can be effectively transformed into a correctness proof in any other of these proof systems. This result is obtained by a detailed proof-theoretic analysis of the three loop rules in the context of these proof systems. To structure the proof well, we make use of auxiliary proof rules, which we show to be admissible in the proof systems.

Even though Hoare's logic has been extensively studied (see for example our survey [**?**]), little work has been done on the analysis of proofs in Hoare's logic.

[4] `https://termination-portal.org/wiki/Termination_Competition`

We are familiar with only three references, [**?**], [**?**], and [**?**], in which transformations of proofs in a Hoare logic are discussed.

While these proof rules are equivalent, their use and representation in the form of proof outlines, which are programs annotated by assertions, differs. We illustrate this by analyzing a termination proof of a program with nested loops. We also address applications in the context of assertions used as annotations in the design by contract paradigm.

2 Preliminaries

NAMES OF TWO SUBSECTIONS ADDED

2.1 Admissibility and equivalence

Assume a given language that is determined by its set of formulas. In what follows we assume that all considered axioms, proof rules, and proof systems are concerned with the same language.

Given a proof system PR and two sequences of formulas $\phi_1, \ldots, \phi_m$ and $\varphi_1, \ldots, \varphi_n$ we write

$$\phi_1, \ldots, \phi_m \vdash_{PR} \varphi_1, \ldots, \varphi_n$$

to denote the fact that each formula φ_i can be proved in PR using as additional axioms the formulas $\phi_1, \ldots, \phi_m$. We also use this notation when the sequence $\phi_1, \ldots, \phi_m$ is empty and when PR is a set of proof rules.

A proof rule

$$(R) \qquad \frac{\varphi_1, \ldots, \varphi_k}{\varphi}$$

is called ***admissible in*** PR (see, e.g. [**?**]) if

$$\vdash_{PR} \varphi_1, \ldots, \vdash_{PR} \varphi_k \text{ implies } \vdash_{PR} \varphi.$$

Intuitively, if a rule is admissible in PR it does not increase the power of the proof system PR [**?**], but it serves as a lemma that simplifies proofs in PR by condensing a detailed proof argument into one application of (R).

We say that two proof systems PR_1 and PR_2 are ***equivalent*** if for all formulas φ

$$\vdash_{PR_1} \varphi \text{ iff } \vdash_{PR_2} \varphi.$$

The notions of admissibility and equivalence are connected by the following simple observation.

Lemma 1. *Consider a proof system PR and proof rules R_1 and R_2.*

The proof systems $PR \cup \{R_1\}$ and $PR \cup \{R_2\}$ are equivalent iff R_1 is admissible in $PR \cup \{R_2\}$ and R_2 is admissible in $PR \cup \{R_1\}$.

2.2 LOOP rules

From now on we shall be concerned with the language, the formulas of which are either first-order formulas, called ***assertions***, or ***correctness formulas***, which are constructs of the form $\{p\}\ S\ \{q\}$, where p and q are assertions and S is a **while** program. Below we denote by $free(p)$ the set of free variables of the assertion p and by $var(t)$, $var(B)$, and $var(S)$ the set of variables that appear in the expression t, the Boolean expression B, and the program S, respectively.

We shall consider in total five proof systems concerned with the correctness formulas. They only differ in the used LOOP rule.

IN THE PARAGRAPH BELOW I CHANGED THE REFERENCE FROM OUR BOOK TO HOARE 69 (THOUGH THE RULE FOR THE IF STATEMENT WAS ADDED LATER BY LAUER. OTHERWISE IT SUGGESTS THAT WE ARE THE MAIN SOURCE FOR THIS PROOF SYSTEM

Proof system I denotes the customary proof system allowing us to prove partial correctness of **while** programs. Its axioms and proof rules, due to [**?**], are listed in the Appendix. Its LOOP rule has the following form:

RULE LOOP I

$$\frac{\{p \wedge B\}\ S\ \{p\}}{\{p\}\ \textbf{while}\ B\ \textbf{do}\ S\ \textbf{od}\ \{p \wedge \neg B\}}$$

In the proof system II this rule is replaced by

RULE LOOP II

$$\frac{\begin{array}{l}\{p \wedge B \wedge t = z\}\ S\ \{p \wedge t < z\},\\ p\ \rightarrow t \geq 0\end{array}}{\{p\}\ \textbf{while}\ B\ \textbf{do}\ S\ \textbf{od}\ \{p \wedge \neg\ B\}}$$

where t is an integer expression such that $var(t) \subseteq var(B) \cup var(S)$ and z is an integer variable that does not appear in p, B, t or S.

In the proof system III LOOP I rule is replaced by

RULE LOOP III

$$\frac{\begin{array}{l}\{p \wedge B\}\ S\ \{p\},\\ \{p \wedge B \wedge t = z\}\ S\ \{t < z\},\\ p\ \rightarrow t \geq 0\end{array}}{\{p\}\ \textbf{while}\ B\ \textbf{do}\ S\ \textbf{od}\ \{p \wedge \neg\ B\}}$$

where t and z are as above.

In the context of the LOOP rules discussed here, the assertion p is called the ***loop invariant*** and the expression t is called the ***bound function***. It provides an estimate how many iterations the loop will still perform before termination. The restriction $var(t) \subseteq var(B) \cup var(S)$ is added to simplify the subsequent proofs. However, it is inessential as shown in Appendix **??**.

We shall next consider the following hybrid rule that combines provability in two proof systems.

RULE LOOP IV

$$\frac{\begin{array}{l}\vdash_{\mathrm{I}} \{p \wedge B\}\ S\ \{p\},\\ \vdash_{\mathrm{I}} \{p \wedge B \wedge t = z\}\ S\ \{t < z\},\\ \{p \wedge B\}\ S\ \{\mathbf{true}\},\\ p\ \rightarrow t \geq 0\end{array}}{\{p\}\ \mathbf{while}\ B\ \mathbf{do}\ S\ \mathbf{od}\ \{p \wedge \neg\ B\}}$$

where t and z are as above.

Proof system IV is obtained from proof system I by replacing the LOOP I rule by the LOOP IV rule. The use of two forms of provability in the premises of this rule can be circumvented by the following modification of the notation. Denote the correctness formulas in the sense of partial correctness by $\{p\}\ S\ \{q\}$ and in the sense of total correctness by $[p]\ S\ [q]$. Then combine the proof system I with the proof system in which the axioms and proof rules of I except the LOOP I rule are rewritten using the $[p]\ S\ [q]$ syntax. Finally, add to this proof system the LOOP IV rewritten as follows:

$$\frac{\begin{array}{l}\{p \wedge B\}\ S\ \{p\},\\ \{p \wedge B \wedge t = z\}\ S\ \{t < z\},\\ [p \wedge B]\ S\ [\mathbf{true}],\\ p\ \rightarrow t \geq 0\end{array}}{[p]\ \mathbf{while}\ B\ \mathbf{do}\ S\ \mathbf{od}\ [p \wedge \neg\ B]}$$

where t and z are as above.

In what follows we use the original formulation of this rule, as it will not lead to any ambiguities.

The final LOOP V rule introduces an additional parameter in the loop invariant instead of a separate bound function.

RULE LOOP V

$$\frac{\begin{array}{l}\{p(x) \wedge B\}\ S\ \{\exists y : p(y) \wedge y < x\},\\ p(x) \rightarrow x \geq 0\end{array}}{\{\exists x : p(x)\}\ \mathbf{while}\ B\ \mathbf{do}\ S\ \mathbf{od}\ \{(\exists x : p(x)) \wedge \neg\ B\}}$$

where x is an integer variable which does not appear in **while** B **do** S **od**, and $p(y)$ denotes the result of replacing x in $p(x)$ by the fresh integer variable y.

ADDED:

Note that the brackets around $(\exists x : p(x))$ can be dropped thanks to the assumption about the variable x.

END ADDED

It is difficult to determine where the LOOP II rule was introduced first. We mentioned it in [**?**]. It also appears in [**?**] on page 64 and in [**?**] on page 151. The LOOP III rule was introduced in [**?**]. It corresponds to Dijkstra's modification of his weakest precondition semantics proposed in [**?**] and reproduced as [**?**].

The LOOP IV rule was introduced in [**?**]. It formalizes the following intuition. In order to prove the termination of a **while** loop it suffices to find a loop invariant and a bound function such that

(i) the loop invariant is maintained by each loop body execution, in the sense of partial correctness,
(ii) each loop body execution decreases the bound function, also in the sense of partial correctness,
(iii) the loop body terminates, and
(iv) the loop invariant implies that the bound function remains non-negative.

This rule supports modular reasoning about program correctness by separating the premises into partial correctness and termination properties. This is of particular relevance in the presence of nested loops, i.e., when the loop body contains inner loops. Then (i) establishes only the partial correctness of the loop body, whereas its termination is relegated to (iii). That the loop body can be iterated only finitely often is established in (ii) in combination with (iv). Note that for loop bodies without inner loops, partial and total correctness coincide. In this case the third premise, $\{p \wedge B\}\ S\ \{\mathbf{true}\}$, can then be dropped, and we arrive at the LOOP III rule. So the hybrid form of this rule arises only for nested loops.

Note that the LOOP II rule results from combining the first two premises of the LOOP III into one. An analogous modification can be carried out in the case of the LOOP IV rule. In what follows we disregard this possibility, given that the resulting analysis is analogous to the one concerning the LOOP II and LOOP III rules.

A version of the LOOP V rule has been introduced in [**?**]. This rule allows to generalize proofs of termination to arbitrary well-founded domains. Such a generalization is for example required for reasoning about termination in presence of *countable non-determinism* [**?**]. The following version of LOOP V separates reasoning about termination along the lines of rule LOOP IV.

$$\frac{\begin{array}{l}\vdash_{\mathrm{I}} \{p(x) \wedge B\}\ S\ \{\exists y : p(y) \wedge y < x\},\\ \{(\exists x : p) \wedge B\}\ S\ \{\mathbf{true}\},\\ p(x) \rightarrow x \geq 0\end{array}}{\{\exists x : p(x)\}\ \mathbf{while}\ B\ \mathbf{do}\ S\ \mathbf{od}\ \{(\exists x : p(x)) \wedge \neg\ B\}}$$

We will restrict our proof-theoretic analysis to the rules LOOP II, III, IV and V. However, it is straightforward to apply the underlying techniques of our analysis to prove the equivalence between LOOP V and this version.

2.3 Auxiliary Rules

In the proof-theoretic analysis of the LOOP II and III rules, we make use of the following two auxiliary rules, see, e.g., [**?**].

RULE CONJUNCTION

$$\frac{\{p_1\}\ S\ \{q_1\}, \{p_2\}\ S\ \{q_2\}}{\{p_1 \wedge p_2\}\ S\ \{q_1 \wedge q_2\}}$$

RULE ∃-INTRODUCTION

$$\frac{\{p\}\ S\ \{q\}}{\{\exists x : p\}\ S\ \{q\}}$$

where x does not occur in S or in $\mathit{free}(q)$.

To reason about the LOOP IV rule we shall consider the following auxiliary rule that combines provability in two proof systems.

RULE HYBRID CONJUNCTION

$$\frac{\begin{array}{l}\vdash_{\mathrm{I}} \{p_1\}\ S\ \{q_1\},\\ \{p_2\}\ S\ \{q_2\}\end{array}}{\{p_1 \wedge p_2\}\ S\ \{q_1 \wedge q_2\}}$$

A special case of this rule is the following rule from [**?**].

RULE DECOMPOSITION

$$\frac{\begin{array}{l}\vdash_{\mathrm{I}} \{p\}\ S\ \{q\},\\ \{p\}\ S\ \{\mathbf{true}\}\end{array}}{\{p\}\ S\ \{q\}}$$

The following theorem states that these rules are indeed auxiliary in the formal sense of admissibility.

THEOREM REFORMULATED

Theorem 1. *The above auxiliary rules are admissible in each proof system I-V.*

Though tedious this is rather straightforward to prove. In the appendix **??** a proof of Theorem **??** is given which restricts to those cases which are actually used in the proof of the main Theorem **??** (in the next section).

3 Admissibility of the Termination Rules

The LOOP II and LOOP III rules look very much the same and it is obvious that they are in some sense equivalent. The following main theorem, that also covers the LOOP IV and LOOP V rules, states this claim in the strongest way.

Theorem 2. *The loop rules are admissible in the other proof systems as follows:*

(i) The LOOP II rule is admissible in the proof system III.
(ii) The LOOP III rule is admissible in the proof system II.
(iii) The LOOP IV rule is admissible in the proof system III.
(iv) The LOOP III rule is admissible in the proof system IV.
(v) The LOOP V rule is admissible in the proof system II.
(vi) The LOOP II rule is admissible in the proof system V.

Together with Lemma **??**, we arrive at the following corollary.

Corollary 1.

(i) The proof systems II, III, IV and V are equivalent.
(ii) Each LOOP α rule is admissible in the proof system β, where $\alpha, \beta \in \{\mathrm{II}, \mathrm{III}, \mathrm{IV}, \mathrm{V}\}$.

Proof. (i) The equivalence relation between proof systems is transitive, so the claim follows by Lemma **??** and Theorem **??**.

(ii) Take a LOOP α rule and a proof system β, where $\alpha, \beta \in \{\mathrm{II}, \mathrm{III}, \mathrm{IV}, \mathrm{V}\}$. By part (i), α and β are equivalent. So by Lemma **??**, the LOOP α rule is admissible in the proof system β. □

The next lemma refers to the premises of the LOOP II and LOOP III rules. The names of the rules are abbreviated in the obvious way.

Lemma 2. *Suppose that z is an integer variable that does not appear in p, B, t or S. Then*

(i) $\{p \wedge B\}\ S\ \{p\},\ \{p \wedge B \wedge t = z\}\ S\ \{t < z\}$
$\vdash_{\{\mathrm{CONJ},\,\mathrm{CONS}\}}$
$\{p \wedge B \wedge t = z\}\ S\ \{p \wedge t < z\}.$

(ii) $\{p \wedge B \wedge t = z\}\ S\ \{p \wedge t < z\}$
$\vdash_{\{\exists-\mathrm{INTRO},\,\mathrm{CONS}\}}$
$\{p \wedge B\}\ S\ \{p\},\ \{p \wedge B \wedge t = z\}\ S\ \{t < z\}.$

Proof. (i) Immediate.

(ii) First note that by the CONSEQUENCE rule we can derive from

$$\{p \wedge B \wedge t = z\}\ S\ \{p \wedge t < z\}$$

both

$$\{p \wedge B \wedge t = z\}\ S\ \{p\}$$

and

$$\{p \wedge B \wedge t = z\}\ S\ \{t < z\}.$$

Next, by the $\exists$-INTRODUCTION rule, we derive from $\{p \wedge B \wedge t = z\}\ S\ \{p\}$

$$\{\exists z : p \wedge B \wedge t = z\}\ S\ \{p\}.$$

By the assumption about the variable z,

$$p \wedge B \rightarrow (p \wedge B \wedge \exists z : t = z) \rightarrow (\exists z : p \wedge B \wedge t = z),$$

so by the CONSEQUENCE rule, we derive $\{p \wedge B\}\ S\ \{p\}$, as desired. □

We are now prepared for the proof of our main result.

Proof of Theorem ??.
(i) Suppose

$$\vdash_{\mathrm{III}} \{p \wedge B \wedge t = z\}\ S\ \{p \wedge t < z\}$$

and that $p\ \rightarrow t \geq 0$ holds. By Lemma **??**(ii),

$$\vdash_{\mathrm{III} \cup \{\exists\text{–INTRO, CONS}\}} \{p \wedge B\}\ S\ \{p\}$$

and

$$\vdash_{\mathrm{III} \cup \{\exists\text{–INTRO, CONS}\}} \{p \wedge B \wedge t = z\}\ S\ \{t < z\}.$$

By Theorem **??** $\vdash_{\mathrm{III}} \{p \wedge B\}\ S\ \{p\}$ and $\vdash_{\mathrm{III}} \{p \wedge B \wedge t = z\}\ S\ \{t < z\}$, so by the LOOP III rule $\vdash_{\mathrm{II}} \{p\}$ **while** B **do** S **od** $\{p \wedge \neg B\}$.

(ii) Suppose

$$\vdash_{\mathrm{II}} \{p \wedge B\}\ S\ \{p\},$$

$$\vdash_{\mathrm{II}} \{p \wedge B \wedge t = z\}\ S\ \{t < z\},$$

and that $p\ \rightarrow t \geq 0$ holds. By Lemma **??**(i),

$$\vdash_{\mathrm{III} \cup \{\mathrm{CONJ, CONS}\}} \{p \wedge B \wedge t = z\}\ S\ \{p \wedge t < z\}.$$

By Theorem **??** $\vdash_{\mathrm{II}} \{p \wedge B \wedge t = z\}\ S\ \{p \wedge t < z\}$, so by the LOOP II rule $\vdash_{\mathrm{II}} \{p\}$ **while** B **do** S **od** $\{p \wedge \neg B\}$.

(iii) Suppose

$$\vdash_{\mathrm{I}} \{p \wedge B\}\ S\ \{p\},$$

$$\vdash_{\mathrm{I}} \{p \wedge B \wedge t = z\}\ S\ \{t < z\},$$

$$\vdash_{\mathrm{III}} \{p \wedge B\}\ S\ \{\mathbf{true}\},$$

and that $p\ \rightarrow t \geq 0$ holds. By Theorem **??** applied twice $\vdash_{\mathrm{III}} \{p \wedge B\}\ S\ \{p\}$ and $\vdash_{\mathrm{III}} \{p \wedge B \wedge t = z\}\ S\ \{t < z\}$.

So by the LOOP III rule $\vdash_{\mathrm{II}} \{p\}$ **while** B **do** S **od** $\{p \wedge \neg B\}$.

(iv) Suppose

$$\vdash_{\mathrm{IV}} \{p \wedge B\}\ S\ \{p\},$$

$$\vdash_{\mathrm{IV}} \{p \wedge B \wedge t = z\}\ S\ \{t < z\},$$

and that $p\ \rightarrow t \geq 0$ holds. By omitting everywhere in these two proofs the second and third premise of the LOOP IV rule whenever this rule is applied,

we get $\vdash_{\mathrm{I}} \{p \wedge B\}\ S\ \{p\}$ and $\vdash_{\mathrm{I}} \{p \wedge B \wedge t = z\}\ S\ \{t < z\}$. Further, by the CONSEQUENCE rule $\vdash_{\mathrm{IV}} \{p \wedge B\}\ S\ \{\mathbf{true}\}$.

So by the LOOP IV rule $\vdash_{\mathrm{IV}} \{p\}$ **while** B **do** S **od** $\{p \wedge \neg B\}$.

(v) Suppose

$$\vdash_{\mathrm{II}} \{p(x) \wedge B\}\ S\ \{\exists y : p(y) \wedge y < x\}$$

and $p(x) \rightarrow x \geq 0$. Let $\bar{u}$ denote the other free variables of $p(x)$ (different from x) and $f(\bar{u})$ be a *Skolem* function such that $(\exists x : p(x)) \rightarrow p(f(\bar{u}))$ and $p(x) \rightarrow f(\bar{u}) \leq x$, so that $f(\bar{u})$ denotes the minimal witness, which exists because $p(x) \rightarrow x \geq 0$. It follows that $\exists y : p(y) \wedge y < x$ implies $(\exists x : p(x)) \wedge f(\bar{u}) < x$ and $(\exists x : p(x)) \wedge f(\bar{u}) = x$ implies $p(x)$. Thus we derive

$$\vdash_{\mathrm{II}} \{(\exists x : p(x)) \wedge B \wedge f(\bar{u}) = x\}\ S\ \{(\exists x : p(x)) \wedge f(\bar{u}) < x\}$$

by the consequence rule (from the above premise of the rule LOOP V). Note that in general $\bar{u} \not\subseteq \mathit{var}(B) \cup \mathit{var}(S)$, so we cannot directly apply the LOOP II rule. Since $(\exists x : p(x)) \rightarrow f(\bar{u}) \geq 0$ and the unrestricted version of this rule is addmissible in the proof system II, we obtain

$$\vdash_{\mathrm{II}} \{\exists x : p(x)\}\ \mathbf{while}\ B\ \mathbf{do}\ S\ \mathbf{od}\ \{\exists x : p(x) \wedge \neg B\}$$

(vi) Suppose

$$\vdash_{\mathrm{V}} \{p \wedge B \wedge t = z\}\ S\ \{p \wedge t < z\}$$

and $p \rightarrow t \geq 0$. Let $p'(z)$ denote the assertion $p \wedge t = z$. We trivially have that $(p \wedge t < z) \rightarrow \exists y : p \wedge t = y \wedge y < z)$ (y fresh). By the consequence rule we thus obtain

$$\vdash_{\mathrm{V}} \{p'(z) \wedge B\}\ S\ \{\exists y : p'(y) \wedge y < z\}$$

from the above premise of the rule LOOP II. Further, $p \rightarrow t \geq 0$ trivially implies $p'(z) \rightarrow z \geq 0$. So we obtain

$$\vdash_{\mathrm{V}} \{\exists z : p'\}\ \mathbf{while}\ B\ \mathbf{do}\ S\ \mathbf{od}\ \{\exists z : p' \wedge \neg B\}$$

by rule LOOP V. Clearly, $(\exists z : p') \leftrightarrow p$. So by a trivial application of the consequence rule we conclude that

$$\vdash_{\mathrm{V}} \{p\}\ \mathbf{while}\ B\ \mathbf{do}\ S\ \mathbf{od}\ \{p \wedge \neg B\}$$

□

4 Soundness and Completeness Matters

Assume now that some notion of semantics of programs and assertions is given that includes the concept of a state, execution of a program, the notion of a state satisfying an assertion, and the truth of an assertion. We write then $\models \{p\}\ S\ \{q\}$ to denote the fact that every execution of S that starts in a state satisfying p

terminates in a state satisfying q and say then that $\{p\}\ S\ \{q\}$ is ***true***. (Thus we are referring to ***total correctness***.) Next, we say then that a proof rule

$$\frac{\varphi_1, \ldots, \varphi_k}{\varphi}$$

is ***sound*** if $\models \varphi_1, \ldots, \models \varphi_k$ implies $\models \varphi$.

In [**?**] we proved that the LOOP III rule is sound, while in [**?**] it was proved that the LOOP II is sound. A natural question arises whether soundness of one of these rules can be directly deduced from the soundness of the other rule. This can be accomplished by modifying the claims of Lemma **??** as follows.

Lemma 3. *Suppose that z is an integer variable that does not appear in p, B, t or S. Then*

(i) If $\models \{p \wedge B\}\ S\ \{p\}$ and $\models \{p \wedge B \wedge t = z\}\ S\ \{t < z\}$,
then $\models \{p \wedge B \wedge t = z\}\ S\ \{p \wedge t < z\}$.
(ii) If $\models \{p \wedge B \wedge t = z\}\ S\ \{p \wedge t < z\}$,
then $\models \{p \wedge B\}\ S\ \{p\}$ and $\models \{p \wedge B \wedge t = z\}\ S\ \{t < z\}$.

Proof. It is a direct consequence of the fact that the proof rules used in Lemma **??** are sound. Thus each time one of these rules is applied, truth of the correctness formulas is preserved. □

Suppose now that the LOOP II rule is sound. To prove the soundness of the LOOP III rule assume that its premises are true. Then by Lemma **??**(i) the premises of the LOOP II rule are true, so by its soundness the conclusion of both rules is true. The same argument shows that soundness of the LOOP III rule implies soundness of the LOOP II rule.

We conclude this discussion by two remarks. First, notice that the last premise of each of the LOOP rules II, III and IV can also be modified, by considering in each case the implication $p \wedge B \rightarrow t \geq 0$ instead of $p \rightarrow t \geq 0$. It is easy to see that such a modification does not affect provability in the considered proof systems II, III, and IV. Indeed, an application of the original rule with a bound function t is also a valid application of the modified rule and an application of the modified rule with a bound function t can be replaced by an application of the original rule with the bound function **if** B **then** t **else** 0 **fi**.

Finally, for the rules LOOP II–IV we assumed that the bound function t satisfies the restriction $var(t) \subseteq var(B) \cup var(S)$. This allows for the proof of admissibility of the $\exists$-INTRODUCTION rule in Theorem **??**, which in turn is used via Lemma **??** in the proof of Theorem **??** and thus in the proof of the equivalence result stated in Corollary **??**. In future we will investigate how to avoid this restriction. We see that the equivalence of the different proof systems for total correctness depends very subtly on the intricate interplay of the loop rules with the admissibility of standard auxiliary rules of Hoare's logic, which requires further investigation.

5 Representing Proofs

5.1 Proof outlines

Even though the LOOP rules for proving termination are equivalent in the sense of Theorem **??** or Corollary **??**, they lead to different proofs of total correctness of **while** programs. This clearly holds for the rule LOOP V which does not involve an explicit bound function but uses the invariant for the specification of a bound. But also the other rules lead to different proofs. The idea behind the LOOP III rule is to establish that p is a loop invariant and t is a bound function separately, while in the LOOP II rule both facts are established simultaneously. So the LOOP III rule looks more convenient when we want to strengthen a proof of partial correctness to a proof of total correctness: it suffices to establish two new premises concerned with the bound function t. In turn, the LOOP IV rule allows us to split the proof obligations even further, by identifying the property actually needed to be proved in terms of total correctness. This, as already mentioned, allows one to support modular reasoning.

However, matters change when we want to represent the proofs in the resulting proof systems in a convenient form. Given that these proofs deal with structured programs, their most natural representation consists of so-called ***proof outlines***, a notion introduced in [**?**]. Informally, it is a proof representation in the form of a program annotated by the assertions arising from the appropriate rule applications. Such a representation is possible thanks to the fact that the proof rules are syntax directed. Proof outlines were introduced in [**?**] in order to reason about correctness of parallel programs, where they served to establish so-called interference freedom among the proofs of the component programs. However, they are also very useful as a representation of correctness proofs of sequential programs and, when some obvious assertions are deleted, as a program documentation.

Now, given that the LOOP III and IV rules have more than one premise consisting of a correctness formula, it is difficult to employ a single proof outline to represent a proof involving any of these rules. This is not the case with the LOOP II and LOOP V rules. To illustrate this point recall first that the proof outlines are defined by induction on the program structure. We only focus on the crucial formation rule concerned with the **while** statement. For the proof system I the following formation rule was used in [**?**]:

$$\frac{\{p \wedge B\}\ S^*\ \{p\}}{\{\mathbf{inv} : p\}\ \mathbf{while}\ B\ \mathbf{do}\ \{p \wedge B\}\ S^*\ \{p\}\ \mathbf{od}\ \{p \wedge \neg B\}}$$

where S^* is the program S annotated with some assertions.

For the proof system II we introduce the following formation rule:

$$\frac{\{p \wedge B \wedge t = z\}\ S^*\ \{p \wedge t < z\},\quad p \rightarrow t \geq 0}{\{\mathbf{inv} : p\}\{\mathbf{bd} : t\}\ \mathbf{while}\ B\ \mathbf{do}\ \{p \wedge B \wedge t = z\}\ S^*\ \{p \wedge t < z\}\ \mathbf{od}\ \{p \wedge \neg B\}}$$

where S^* and S^{**} are annotations of the program S with some assertions, t is an integer expression and z is an integer variable not occurring in p, t, B or S^*.[5]

For a moment we defer the discussion of proof outlines for the proof system IV. One can easily prove by induction that each proof outline for the proof system I corresponds to a proof in this proof system. For example, if by the induction hypothesis the proof outline $\{p \wedge B\}\ S^*\ \{p\}$ corresponds to a proof of $\{p \wedge B\}\ S\ \{p\}$ in I, then the proof outline

$$\{\mathbf{inv} : p\}\ \mathbf{while}\ B\ \mathbf{do}\ \{p \wedge B\}\ S^*\ \{p\}\ \mathbf{od}\ \{p \wedge \neg B\}$$

corresponds to a proof of $\{p\}$ **while** B **do** S **od** $\{p \wedge \neg B\}$ in I, which is obtained by applying to $\{p \wedge B\}\ S\ \{p\}$ the LOOP I rule.

However, this property fails to hold for the proof outlines for the proof system III, because the second proof outline used in the premise of the formation rule is dropped. As a consequence, the proof outline for the **while** statement does not allow one to reconstruct the proof of $\{p\}$ **while** B **do** S **od** $\{p \wedge \neg B\}$ in the proof system III.

5.2 Proofs using the LOOP II rule

By contrast, each proof outline for the proof system II involving the LOOP II rule does correspond to a proof in this proof system, because all assertions used are retained. In other words, from each proof outline for the proof system II a proof in this system can be extracted.

To illustrate this point consider the following program S_N involving nested loops, suggested to us by Tobias Nipkow (private communication):

```
S_N ≡ while i < n do
          j := i;
          while 0 < j do
              j := j − 1
          od;
          i := i + 1
      od
```

where i, j, n are integer variables.

We would like to prove that it terminates for all initial states. To this end, we prove the correctness formula $\{\mathbf{true}\}\ S_N\ \{\mathbf{true}\}$ in the proof system II. In

[5] In [**?**], in contrast to [**?**], there is a typo and S^{**} is mentioned here instead of S^*.

Fig. **??**, we show the proof in the form of a proof outline, instantiating the corresponding formation rule for the **while** statement in proof system II with the following loop invariants and bound functions for the outer and the inner loop, respectively:

$$p \equiv \mathbf{true} \text{ and } t \equiv \mathit{max}(n-i,0),$$
$$p \equiv n-i = z_1 \wedge z_1 > 0 \text{ and } t \equiv \mathit{max}(j,0).$$

Note that in a proof outline adjacent assertions stand for implications according to an application of the CONSEQUENCE rule. For instance, the assertion in line 3 implies that of line 4. Assignments are treated by backward substitution according to the ASSIGNMENT axiom. For instance, the assignment $i := i+1$ in line 18 is dealt with by substituting i by $i+1$ in the assertion in line 19, yielding the assertion in line 17.

1 $\{\mathbf{inv} : \mathbf{true}\}$ $\{\mathbf{bd} : \mathit{max}(n-i,0)\}$
2 **while** $i < n$ **do**
3 $\{\mathbf{true} \wedge i < n \wedge \mathit{max}(n-i,0) = z_1\}$
4 $\{n-i = z_1 \wedge z_1 > 0\}$
5 $j := i;$
6 $\{n-i = z_1 \wedge z_1 > 0\}$
7 $\{\mathbf{inv} : n-i = z_1 \wedge z_1 > 0\}$ $\{\mathbf{bd} : \mathit{max}(j,0)\}$
8 **while** $0 < j$ **do**
9 $\{n-i = z_1 \wedge z_1 > 0 \wedge 0 < j \wedge \mathit{max}(j,0) = z_2\}$
10 $\{n-i = z_1 \wedge z_1 > 0 \wedge j = z_2 \wedge z_2 > 0\}$
11 $\{n-i = z_1 \wedge z_1 > 0 \wedge j-1 < z_2 \wedge z_2 > 0\}$
12 $j := j-1$
13 $\{n-i = z_1 \wedge z_1 > 0 \wedge j < z_2 \wedge z_2 > 0\}$
14 $\{n-i = z_1 \wedge z_1 > 0 \wedge \mathit{max}(j,0) < z_2\}$
15 **od**;
16 $\{n-i = z_1 \wedge z_1 > 0 \wedge \neg(0 < j)\}$
17 $\{n-(i+1) < z_1 \wedge z_1 > 0\}$
18 $i := i+1$
19 $\{n-i < z_1 \wedge z_1 > 0\}$
20 $\{\mathbf{true} \wedge \mathit{max}(n-i,0) < z_1\}$
21 **od**
22 $\{\mathbf{true} \wedge \neg(i < n)\}$
23 $\{\mathbf{true}\}$

Fig. 1. Proof outline for $\{\mathbf{true}\}$ S_N $\{\mathbf{true}\}$ in the proof system II. The line numbers have been added for reference only.

5.3 Proofs using the LOOP IV rule

Let us now move on to a discussion of the proofs involving the LOOP IV rule. This rule, just like the LOOP III rule, uses more than one correctness formula

as a premise. As a result, it shares with the LOOP III rule the problem that it is not clear how to faithfully represent correctness proofs using a single proof outline. Indeed, each of the first three premises calls for a separate proof outline.

But it is not difficult to see that this would give rise to largely overlapping proof outlines. So, instead, we propose an alternative approach in which we replace these overlapping proof outlines referring to the proof system IV by an interrelated *set* of proof outlines in the sense of *partial correctness*, so referring to the proof system I.

For a given correctness formula $\{p\}\ S\ \{q\}$ to be proved in the proof system IV this set is defined as follows. First, we have a proof outline $\{p\}\ S^*\ \{q\}$ that employs the first formation rule given above and thus represents a proof of partial correctness of $\{p\}\ S\ \{q\}$ in I.

Next, for each occurrence of a loop **while** B **do** S_0 **od** in S, we have a proof outline $\{p_0 \wedge B \wedge t = z\}\ S_0^*\ \{t < z\}$, which represents a proof of partial correctness of $\{p_0 \wedge B \wedge t = z\}\ S_0\ \{t < z\}$ in I. Here p_0 is the invariant associated with the occurrence of the loop **while** B **do** S_0 **od** in the above proof outline $\{p\}\ S^*\ \{q\}$ and t is some bound function t such that $p_0 \rightarrow t \geq 0$. We omit the proof that existence of such a set of proof outlines in the sense of partial correctness ensures a proof of the corresponding correctness formula in the proof system IV. Intuitively, the use of the above proof outlines for *each* loop occurrence in S ensures by structural induction the third premise of the LOOP IV rule, so $\{p \wedge B\}\ S\ \{\mathbf{true}\}$ in the sense of total correctness.

We illustrate how such a set of proof outlines can be used to establish the proof of the correctness formula $\{\mathbf{true}\}\ S_N\ \{\mathbf{true}\}$ for Nipkow's program S_N in the proof system IV. We skip the trivial proof outline $\{\mathbf{true}\}\ S_N^*\ \{\mathbf{true}\}$, which corresponds to a proof of partial correctness of $\{\mathbf{true}\}\ S_N\ \{\mathbf{true}\}$. Let S_0 denote the body of the outer loop of S_N. Given the bound function $max(n-i, 0)$, Fig. **??** shows the proof outline which corresponds to a proof of partial correctness of

$$\{\mathbf{true} \wedge i < n \wedge max(n-i, 0) = z\}\ S_0\ \{max(n-i, 0) < z\},$$

assuming that the trivial invariant **true** is associated with this loop in the proof outline $\{\mathbf{true}\}\ S_N^*\ \{\mathbf{true}\}$. Note that $\mathbf{true} \rightarrow max(n-i, 0) \geq 0$.

Finally, the LOOP IV rule requires us to prove termination of S_0. This boils down to establish the termination of the inner loop. To this end, we introduce the bound function $max(j, 0)$. Note that $\mathbf{true} \rightarrow max(j, 0) \geq 0$. Fig. **??** shows the proof outline which corresponds to a proof of partial correctness of

$$\{\mathbf{true} \wedge 0 < j \wedge max(j, 0) = z\}\ j := j - 1\ \{max(j, 0) < z\}.$$

ADDED

5.4 Proofs using the LOOP V rule

To conclude let us discuss proof outlines in presence of the LOOP V rule. To obtain a proof outline for $\{\mathbf{true}\}\ S_N\ \{\mathbf{true}\}$ in the proof system V it suffices to

$\{\mathbf{true} \wedge i < n \wedge max(n-i,0) = z\}$
$\{n - i = z \wedge z > 0\}$
$\quad j := i;$
$\{n - i = z \wedge z > 0\}$
$\{\mathbf{inv} : n - i = z \wedge z > 0\}$
$\quad\quad \mathbf{while}\ 0 < j\ \mathbf{do}$
$\quad\quad \{n - i = z \wedge z > 0\}$
$\quad\quad\quad j := j - 1$
$\quad\quad \{n - i = z \wedge z > 0\}$
$\mathbf{od};$
$\{n - i = z \wedge z > 0 \wedge \neg(0 < j)\}$
$\{n - (i+1) < z \wedge z > 0\}$
$\quad i := i + 1$
$\{n - i < z \wedge z > 0\}$
$\{max(n-i,0) < z\}$

Fig. 2. Proof outline for $\{\mathbf{true} \wedge i < n \wedge max(n-i,0) = z\}\ S_0\ \{max(n-i,0) < z\}$ in the proof system I.

$\{\mathbf{true} \wedge 0 < j \wedge max(j,0) = z\}$
$\{j - 1 < z \wedge z > 0\}$
$\quad j := j - 1$
$\{j < z \wedge z > 0\}$
$\{max(j,0) < z\}$

Fig. 3. Proof outline for $\{\mathbf{true} \wedge 0 < j \wedge max(j,0) = z\}\ j := j - 1\ \{max(j,0) < z\}$ in the proof system I.

modify the loop invariants in the proof outline for the proof system II given in Figure **??** as in the proof of Theorem **??**(vi), so by using, respectively,

$$p(z) \equiv max(n-i,0) = z,$$

$$p(z) \equiv n - i = z_1 \wedge z_1 > 0 \wedge max(j,0) = z,$$

and to drop the references to the bound functions.

END ADDED

6 Practical Applications

Research on program verification has entered practice most visibly by the use of assertions as annotations of programs and program interfaces that remain to be implemented. This can be seen in the paradigm of *design by contract*

introduced by Bertrand Meyer for his object-oriented programming language Eiffel [**?**]: program design starts with a specification in terms of assertions, the contract, against which the program is to be checked either statically by means of a proof or dynamically at runtime.

This paradigm has been adopted and extended to other programming languages, in particular to Java. The *Java Modeling Language* (JML) enriches Java with facilities for writing assertions (pre- and postconditions as well as class invariants) but also with a concept of abstract state space (using so-called model variables) [**?**][6]. For assertions, JML uses Java's Boolean expressions extended by universal and existential quantifiers. They are directly written into the Java source code in the form of comments starting with the symbols `//@` (so that the annotated source code may be processed by both an ordinary Java compiler and a specialized JML tool).

JML is designed to deal with the specification of Java classes, but here we focus on loops. JML provides the designated keywords `requires` for specifying the precondition, `ensures` for the postcondition, `loop_invariant`, and `loop_decreases` for the bound function. To enhance readability, the pre- and postcondition as well as the loop invariant may be split into several assertions, each one stated after a separate repeated keyword[7]. An annotated Java program corresponds to a proof outline as discussed here, but restricted to the essential assertions for each loop (pre- and postcondition, loop invariant and bound function).

Also for the programming language C, a standardized specification language for C programs, called ACSL and inspired by JML, has been designed, see for instance [**?**][8].

In this paper, we have shown that the rules LOOP II-V for proving termination of loops are equivalent. With respect to the number of premises LOOP II and LOOP V are clearly the simplest rules for proving termination. However, a proof using these rules in general require more complex assertions because of the accumulation of the specifications of the bound functions of inner loops. This can be seen in the proof outline given in Fig. 1, in which the assertions inside the inner loop refer to both z_1 and z_2, where z_1 is the variable freezing the value of the bound function $max(n-i, 0)$ of the outer loop and z_2 is the variable freezing the value of the bound function $max(j, 0)$ of the inner loop. The reason is that in this proof outline we need to establish that the value of the bound function of the outer loop is not affected by the inner loop and that the value of the other bound function decreases.

The LOOP IV rule allows for a separate proof of termination of each loop, which does not require the specification of bound functions for the inner loops. Thus we have a trade off between a single complex proof and a number of simpler proofs, where complexity is measured by the size of the assertions. What works best in practice depends on the particular program structure.

[6] See also `https://www.cs.ucf.edu/~leavens/JML/index.shtml`
[7] For an example, see `https://www.openjml.org/examples/binary-search.html`
[8] See also `https://frama-c.com/download/acsl-1.20.pdf`

A Proof System I

The proof system I consists of the following axioms and rules:

AXIOM SKIP

$$\{p\}\ skip\ \{p\}$$

AXIOM ASSIGNMENT

$$\{p[u := t]\}\ u := t\ \{p\}$$

RULE COMPOSITION

$$\frac{\{p\}\ S_1\ \{r\}, \{r\}\ S_2\ \{q\}}{\{p\}\ S_1;\ S_2\ \{q\}}$$

RULE CONDITIONAL

$$\frac{\{p \wedge B\}\ S_1\ \{q\}, \{p \wedge \neg B\}\ S_2\ \{q\}}{\{p\}\ \mathbf{if}\ B\ \mathbf{then}\ S_1\ \mathbf{else}\ S_2\ \mathbf{fi}\ \{q\}}$$

RULE LOOP I

$$\frac{\{p \wedge B\}\ S\ \{p\}}{\{p\}\ \mathbf{while}\ B\ \mathbf{do}\ S\ \mathbf{od}\ \{p \wedge \neg B\}}$$

RULE CONSEQUENCE

$$\frac{p \rightarrow p_1, \{p_1\}\ S\ \{q_1\}, q_1 \rightarrow q}{\{p\}\ S\ \{q\}}$$

Additionally, given an interpretation $\mathcal{I}$ for the underlying first-order language, we use as axioms all assertions that are true in $\mathcal{I}$. These assertions are used as premises in the CONSEQUENCE rule.

B Admissibility Auxiliary Rules

Theorem 3. *The auxiliary rules* $\exists$-INTRODUCTION*,* CONJUNCTION*, and* HYBRID CONJUNCTION*, are admissible in the following proof systems:*

(i) The $\exists$-INTRODUCTION *rule is admissible in the proof system* II*.*
(ii) The $\exists$-INTRODUCTION *rule is admissible in the proof system* III*.*
(iii) The CONJUNCTION *rule is admissible in the proof system* II*.*
(iv) The HYBRID CONJUNCTION *rule is admissible in the proof system* III*, that is,*
if $\vdash_{\mathrm{I}} \{p_1\}\ S\ \{q_1\}$ *and* $\vdash_{\mathrm{III}} \{p_2\}\ S\ \{q_2\}$*, then* $\vdash_{\mathrm{III}} \{p_1 \wedge p_2\}\ S\ \{q_1 \wedge q_2\}$*.*

To prove this theorem we first establish two lemmas that provide additional information about the proofs in the considered proof systems.

Lemma 4. *Let PR be one of the proof systems I, II, III, IV, or V. If PR* $\vdash \varphi$*, there exists a proof of* φ *in PR with exactly one final application of the CONSEQUENCE rule.*

Proof. Since implication $\rightarrow$ is reflexive, we can always add to a given proof in *PR* one final application of the CONSEQUENCE rule. Since implication is transitive, successive applications of the CONSEQUENCE rule in a proof in *PR* can be condensed into one application. □

Lemma 5. *Let PR be one of the proof systems I, II, III, IV, or V.*

(i) Suppose that $\vdash_{PR} \{p\}\ S_1;\ S_2\ \{q\}$*. Then for some assertion* r

$$\vdash_{PR} \{p\}\ S_1\ \{r\} \text{ and } \vdash_{PR} \{r\}\ S_2\ \{q\}.$$

(ii) Suppose that $\vdash_{PR} \{p\}$ **if** B **then** S_1 **else** S_2 **fi** $\{q\}$*. Then*

$$\vdash_{PR} \{p \wedge B\}\ S_1\ \{q\} \text{ and } \vdash_{PR} \{p \wedge \neg B\}\ S_2\ \{q\}.$$

Proof. (i) By Lemma **??**, the considered correctness formula was proved using the COMPOSITION rule followed by a single application of the CONSEQUENCE rule. So for some assertions p_1, r, q_1, we have

$$\vdash_{PR} \{p_1\}\ S_1\ \{r\} \text{ and } \vdash_{PR} \{r\}\ S_2\ \{q_1\}$$

and the implications $p \rightarrow p_1$ and $q_1 \rightarrow q$ hold. We now get the claim by the CONSEQUENCE rule.

(ii) The argument is analogous as in (ii). □

We now turn to the proof of Theorem **??**, repeating the four statements at the beginning of each proof part.

Proof. (i) The $\exists$-INTRODUCTION rule is admissible in the proof system II.

We proceed by induction on the structure of S and consider a proof of $\{p\}\ S\ \{q\}$ in the proof system II, where the last two steps involve an axiom or a rule of the proof system II for the top-level operator of the program S, followed by one final application of the CONSEQUENCE rule according to Lemma **??**. In all cases we assume that $x \notin var(S) \cup free(q)$.

- *Case* $S \equiv u := t$. Thus suppose $\vdash_{\mathrm{II}} \{p\}\ S\ \{q\}$. Then for some assertion q_1,

$$\vdash_{\mathrm{II}} \{q_1[u := t]\}\ S\ \{q_1\}$$

by the ASSIGNMENT axiom, and the implications $p \rightarrow q_1[u := t]$ and $q_1 \rightarrow q$ hold. We assumed that $x \notin var(S)$, so $x \not\equiv u$. By the assignment axiom, also

$$\{(\exists x : q_1)[u := t]\}\ S\ \{\exists x : q_1\}.$$

Note that the implications $\exists x : p \rightarrow \exists x : (q_1[u := t])$ and $\exists x : q_1 \rightarrow \exists x : q$ hold. Since $x \not\equiv u$ and $x \notin \mathit{free}(q)$, also the implications

$$\exists x : (q_1[u := t]) \rightarrow (\exists x : q_1)[u := t] \quad \text{and} \quad (\exists x : q) \rightarrow q$$

hold. So the CONSEQUENCE rule yields $\vdash_{\mathrm{II}} \{\exists x : p\}\ S\ \{q\}$, as desired.

• *Case* $S \equiv S_1;\ S_2$. Thus suppose $\vdash_{\mathrm{II}} \{p\}\ S\ \{q\}$. By Lemma **??**$(i)$ for some assertion r,

$$\vdash_{\mathrm{II}} \{p\}\ S_1\ \{r\} \text{ and } \vdash_{\mathrm{II}} \{r\}\ S_2\ \{q\}.$$

Since $r \rightarrow \exists x : r$, the CONSEQUENCE rule yields $\vdash_{\mathrm{II}} \{p\}\ S_1\ \{\exists x : r\}$. Since $x \notin \mathit{free}(\exists x : r)$, the induction hypothesis yields $\vdash_{\mathrm{II}} \{\exists x : p\}\ S_1\ \{\exists x : r\}$. Since $x \notin \mathit{free}(q)$, by the induction hypothesis, also $\vdash_{\mathrm{II}} \{\exists x : r\}\ S_2\ \{q\}$. Thus by the COMPOSITION rule, $\vdash_{\mathrm{II}} \{\exists x : p\}\ S\ \{q\}$, as desired.

• *Case* $S \equiv$ **if** B **then** S_1 **else** S_2 **fi**. Thus suppose $\vdash_{\mathrm{II}} \{p\}\ S\ \{q\}$. Then by Lemma **??**$(ii)$,

$$\vdash_{\mathrm{II}} \{p \wedge B\}\ S_1\ \{q\} \text{ and } \vdash_{\mathrm{II}} \{p \wedge \neg B\}\ S_2\ \{q\}.$$

By the induction hypothesis,

$$\vdash_{\mathrm{II}} \{\exists x : (p \wedge B)\}\ S_1\ \{q\} \text{ and } \vdash_{\mathrm{II}} \{\exists x : (p \wedge \neg B)\}\ S_2\ \{q\}.$$

Since $x \notin \mathit{var}(B)$, the CONSEQUENCE rule yields

$$\vdash_{\mathrm{II}} \{(\exists x : p) \wedge B\}\ S_1\ \{q\} \text{ and } \vdash_{\mathrm{II}} \{(\exists x : p) \wedge \neg B\}\ S_2\ \{q\}.$$

By the CONDITIONAL rule, $\vdash_{\mathrm{II}} \{\exists x : p\}\ S\ \{q\}$, as desired.

• *Case* $S \equiv$ **while** B **do** S_0 **od**. Thus suppose $\vdash_{\mathrm{II}} \{p\}\ S\ \{q\}$. By the assumption about t, we have $x \notin \mathit{var}(t)$ and without loss of generality we can assume $x \neq z$. Then for some assertion p_0 and an appropriate bound function t and variable z,

$$\vdash_{\mathrm{II}} \{p_0 \wedge B \wedge t = z\}\ S_0\ \{p_0 \wedge t < z\},$$

and the implications $p \rightarrow p_0$, $p_0 \rightarrow t \geq 0$, $(p_0 \wedge \neg B) \rightarrow q$ hold.

Since $(p_0 \wedge t < z) \rightarrow ((\exists x : p_0) \wedge t < z)$, the CONSEQUENCE rule yields

$$\vdash_{\mathrm{II}} \{p_0 \wedge B \wedge t = z\}\ S_0\ \{(\exists x : p_0) \wedge t < z\}.$$

By the induction hypothesis, the assumption about x, and the CONSEQUENCE rule

$$\vdash_{\mathrm{II}} \{(\exists x : p_0) \wedge B \wedge t = z\}\ S_0\ \{(\exists x : p_0) \wedge t < z\}.$$

So the LOOP II rule yields

$$\vdash_{\mathrm{II}} \{\exists x : p_0\}\ S\ \{(\exists x : p_0) \wedge \neg B\}.$$

Further, since $(p_0 \wedge \neg B) \rightarrow q$ holds and $x \notin (\mathit{free}(q) \cup \mathit{var}(B))$, also the implication $((\exists x : p_0) \wedge \neg B) \rightarrow q$ holds. So a final application of the CONSEQUENCE rule yields $\vdash_{\mathrm{II}} \{\exists x : p\}\ S\ \{q\}$, as desired.

(*ii*) The $\exists$-INTRODUCTION rule is admissible in the proof system III.

Again, we proceed by induction on the structure of S, but now consider a proof of $\{p\}\ S\ \{q\}$ in the proof system III, where the last two steps involve the axiom or rule of the proof system III for the top-level operator of the program S, followed by one final application of the CONSEQUENCE rule according to Lemma **??**.

Except for the **while** statement, all cases are analogous, with $\vdash_{\mathrm{II}}$ replaced by $\vdash_{\mathrm{III}}$. The case of the **while** statement differs from (*i*) only in that one now considers two correctness formulas in the premise of the LOOP III rule instead of one. As above, let $S \equiv$ **while** B **do** S_0 **od** and suppose $\vdash_{\mathrm{III}} \{p\}\ S\ \{q\}$. By the assumption about t, we have $x \notin var(t)$ and without loss of generality we can assume $x \neq z$. Then for some assertion p_0 and an appropriate bound function t and variable z,

$$\vdash_{\mathrm{III}} \{p_0 \wedge B\}\ S_0\ \{p_0\},$$

$$\vdash_{\mathrm{III}} \{p_0 \wedge B \wedge t = z\}\ S_0\ \{t < z\},$$

and the implications $p \to p_0$, $p_0 \to t \geq 0$, $(p_0 \wedge \neg B) \to q$ hold.

Since $p_0 \to \exists x : p_0$, the CONSEQUENCE rule yields

$$\vdash_{\mathrm{III}} \{p_0 \wedge B\}\ S_0\ \{\exists x : p_0\}.$$

By the induction hypothesis, the assumption about x, and the CONSEQUENCE rule both

$$\vdash_{\mathrm{III}} \{(\exists x : p_0) \wedge B\}\ S_0\ \{\exists x : p_0\}$$

and

$$\vdash_{\mathrm{III}} \{(\exists x : p_0) \wedge B \wedge t = z\}\ S_0\ \{t < z\}.$$

So the LOOP III rule yields

$$\vdash_{\mathrm{III}} \{\exists x : p_0\}\ S\ \{(\exists x : p_0) \wedge \neg B\}.$$

Further, since $(p_0 \wedge \neg B) \to q$ holds and $x \notin (\mathit{free}(q) \cup var(B))$, also the implication $((\exists x : p_0) \wedge \neg B) \to q$ holds. So a final application of the CONSEQUENCE rule yields $\vdash_{\mathrm{III}} \{\exists x : p\}\ S\ \{q\}$, as desired.

(*iii*) The CONJUNCTION rule is admissible in the proof system II.

We proceed by induction on the structure of S and for $i = 1, 2$ consider proofs of $\{p_i\}\ S\ \{q_i\}$ in the proof system II, where the last two steps involve the axiom or rule of the proof system II for the top-level operator of the program S, followed by one final application of the CONSEQUENCE rule according to Lemma **??**.

• *Case* $S \equiv u := t$. Thus suppose $\vdash_{\mathrm{II}} \{p_1\}\ S\ \{q_1\}$ and $\vdash_{\mathrm{II}} \{p_2\}\ S\ \{q_2\}$. Then for some assertions q_{01} and q_{02}, by the ASSIGNMENT axiom, both

$$\vdash_{\mathrm{II}} \{q_{01}[u := t]\}\ S\ \{q_{01}\} \quad \text{and} \quad \vdash_{\mathrm{II}} \{q_{02}[u := t]\}\ S\ \{q_{02}\},$$

and the implications $p_1 \to q_{01}[u := t]$, $q_{01} \to q_1$ and $p_2 \to q_{02}[u := t]$, $q_{02} \to q_2$ hold. The ASSIGNMENT axiom also yields

$$\{q_{01}[u := t] \wedge q_{02}[u := t]\}\ S\ \{q_{01} \wedge q_{02}\}.$$

By the implications $(p_1 \wedge p_2) \to (q_{01}[u := t] \wedge q_{02}[u := t])$ and $(q_{01} \wedge q_{02}) \to (q_1 \wedge q_2)$, the CONSEQUENCE rule yields

$$\vdash_{\mathrm{II}} \{p_1 \wedge p_2\}\ S\ \{q_1 \wedge q_2\},$$

as desired.

• *Case* $S \equiv S_1;\ S_2$. Thus suppose $\vdash_{\mathrm{II}} \{p_1\}\ S\ \{q_1\}$ and $\vdash_{\mathrm{II}} \{p_2\}\ S\ \{q_2\}$. Then by Lemma **??**$(i)$ for some assertions r_1 and r_2

$$\vdash_{\mathrm{II}} \{p_1\}\ S_1\ \{r_1\} \quad \text{and} \quad \vdash_{\mathrm{II}} \{r_1\}\ S_2\ \{q_1\},$$

$$\vdash_{\mathrm{II}} \{p_2\}\ S_1\ \{r_2\} \quad \text{and} \quad \vdash_{\mathrm{II}} \{r_2\}\ S_2\ \{q_2\}.$$

By the induction hypothesis,

$$\vdash_{\mathrm{II}} \{p_1 \wedge p_2\}\ S_1\ \{r_1 \wedge r_2\} \quad \text{and} \quad \vdash_{\mathrm{II}} \{r_1 \wedge r_2\}\ S_1\ \{q_1 \wedge q_2\},$$

so by the COMPOSITION rule,

$$\vdash_{\mathrm{II}} \{p_1 \wedge p_2\}\ S_1;\ S_2\ \{q_1 \wedge q_2\},$$

as desired.

• *Case* $S \equiv$ **if** B **then** S_1 **else** S_2 **fi**. Thus suppose $\vdash_{\mathrm{II}} \{p_1\}\ S\ \{q_1\}$ and $\vdash_{\mathrm{II}} \{p_2\}\ S\ \{q_2\}$. Then by Lemma **??**$(ii)$,

$$\vdash_{\mathrm{II}} \{p_1 \wedge B\}\ S_1\ \{q_1\} \quad \text{and} \quad \vdash_{\mathrm{II}} \{p_1 \wedge \neg B\}\ S_2\ \{q_1\},$$

$$\vdash_{\mathrm{II}} \{p_2 \wedge B\}\ S_1\ \{q_2\} \quad \text{and} \quad \vdash_{\mathrm{II}} \{p_2 \wedge \neg B\}\ S_2\ \{q_2\}.$$

By the induction hypothesis,

$$\vdash_{\mathrm{II}} \{p_1 \wedge p_2 \wedge B\}\ S_1\ \{q_1 \wedge q_2\}$$

and

$$\vdash_{\mathrm{II}} \{p_1 \wedge p_2 \wedge \neg B\}\ S_2\ \{q_1 \wedge q_2\}.$$

So by the CONDITIONAL rule,

$$\vdash_{\mathrm{II}} \{p_1 \wedge p_2\}\ \textbf{if}\ B\ \textbf{then}\ S_1\ \textbf{else}\ S_2\ \textbf{fi}\ \{q_1 \wedge q_2\},$$

as desired.

• *Case* $S \equiv$ **while** B **do** S_0 **od**. Suppose $\vdash_{\mathrm{II}} \{p_1\}\ S\ \{q_1\}$ and $\vdash_{\mathrm{II}} \{p_2\}\ S\ \{q_2\}$. Then for some assertions p_{01}, p_{02}, appropriate bound functions t_1, t_2 and variables z_1, z_2,

$$\vdash_{\mathrm{II}} \{p_{01} \wedge B \wedge t_1 = z_1\}\ S_0\ \{p_{01} \wedge t_1 < z_1\},$$

$$\vdash_{\mathrm{II}} \{p_{02} \wedge B \wedge t_2 = z_2\}\ S_0\ \{p_{02} \wedge t_2 < z_2\},$$

the implications $p_{01} \to t_1 \geq 0$, $p_1 \to p_{01}$, $(p_{01} \wedge \neg B) \to q_1$ and $p_{02} \to t_2 \geq 0$, $p_2 \to p_{02}$, $(p_{02} \wedge \neg B) \to q_2$ hold.

Without loss of generality we can assume that $z_2 \notin \{z_1\} \cup \mathit{free}(p_{01})$. By the induction hypothesis,

$$\vdash_{\mathrm{II}} \{p_{01} \wedge p_{02} \wedge B \wedge t_1 = z_1 \wedge t_2 = z_2\}\ S_0\ \{p_{01} \wedge p_{02} \wedge t_1 < z_1 \wedge t_2 < z_2\}.$$

It suffices to consider one bound function, say t_1. Formally, we show this as follows. By the CONSEQUENCE rule,

$$\vdash_{\mathrm{II}} \{p_{01} \wedge p_{02} \wedge B \wedge t_1 = z_1 \wedge t_2 = z_2\}\ S_0\ \{p_{01} \wedge p_{02} \wedge t_1 < z_1\}.$$

Now, an application of the $\exists$-INTRODUCTION rule, which is admissible in the proof system II according to part (ii) of this theorem, followed by an application of the CONSEQUENCE rule yields

$$\vdash_{\mathrm{II}} \{p_{01} \wedge p_{02} \wedge B \wedge t_1 = z_1 \wedge \exists z_2 : t_2 = z_2\}\ S_0\ \{p_{01} \wedge p_{02} \wedge t_1 < z_1\}.$$

A further application of the CONSEQUENCE rule yields

$$\vdash_{\mathrm{II}} \{p_{01} \wedge p_{02} \wedge B \wedge t_1 = z_1\}\ S_0\ \{p_{01} \wedge p_{02} \wedge t_1 < z_1\}.$$

Then by the LOOP II rule,

$$\vdash_{\mathrm{II}} \{p_{01} \wedge p_{02}\}\ \textbf{while}\ B\ \textbf{do}\ S_0\ \textbf{od}\ \{p_{01} \wedge p_{02} \wedge \neg B\}.$$

The implications above yield $(p_1 \wedge p_2) \to (p_{01} \wedge p_{02})$ and $(p_{01} \wedge p_{02} \wedge \neg B) \to (q_1 \wedge q_2)$. Thus by the CONSEQUENCE rule,

$$\vdash_{\mathrm{II}} \{p_1 \wedge p_2\}\ \textbf{while}\ B\ \textbf{do}\ S_0\ \textbf{od}\ \{q_1 \wedge q_2\},$$

as desired.

(iv) The HYBRID CONJUNCTION rule is admissible in the proof system III, that is, if $\vdash_{\mathrm{I}} \{p_1\}\ S\ \{q_1\}$ and $\vdash_{\mathrm{III}} \{p_2\}\ S\ \{q_2\}$, then $\vdash_{\mathrm{III}} \{p_1 \wedge p_2\}\ S\ \{q_1 \wedge q_2\}$.

The proof is analogous to the proof of (iii). The only case that is somewhat different is the one concerned with the **while** statement. So we only deal with

• *Case* $S \equiv$ **while** B **do** S_0 **od**. Suppose $\vdash_{\mathrm{I}} \{p_1\}\ S\ \{q_1\}$ and $\vdash_{\mathrm{III}} \{p_2\}\ S\ \{q_2\}$. Then for some assertions p_{01}, p_{02} and an appropriate bound function t and variable z,

$$\vdash_{\mathrm{I}} \{p_{01} \wedge B\}\ S_0\ \{p_{01}\},$$

$$\vdash_{\mathrm{III}} \{p_{02} \wedge B\}\ S_0\ \{p_{02}\},$$

$$\vdash_{\mathrm{III}} \{p_{02} \wedge B \wedge t = z\}\ S_0\ \{t < z\},$$

and the implications

$$p_1 \to p_{01},\ (p_{01} \wedge \neg B) \to q_1,\ p_{02} \to t \geq 0,\ p_2 \to p_{02},\ (p_{02} \wedge \neg B) \to q_2$$

hold. By the induction hypothesis,

$$\vdash_{\mathrm{III}} \{p_{01} \wedge p_{02} \wedge B\}\ S_0\ \{p_{01} \wedge p_{02}\},$$

and by the induction hypothesis combined with the CONSEQUENCE rule,

$$\vdash_{\mathrm{III}} \{p_{01} \wedge p_{02} \wedge B \wedge t = z\}\ S_0\ \{t < z\}.$$

So by the LOOP II rule,

$$\vdash_{\mathrm{III}} \{p_{01} \wedge p_{02}\}\ \mathbf{while}\ B\ \mathbf{do}\ S_0\ \mathbf{od}\ \{p_{01} \wedge p_{02} \wedge \neg B\}.$$

The implications above yield $(p_1 \wedge p_2) \rightarrow (p_{01} \wedge p_{02})$ and $(p_{01} \wedge p_{02} \wedge \neg B) \rightarrow (q_1 \wedge q_2)$. Thus by the CONSEQUENCE rule,

$$\vdash_{\mathrm{III}} \{p_1 \wedge p_2\}\ \mathbf{while}\ B\ \mathbf{do}\ S_0\ \mathbf{od}\ \{q_1 \wedge q_2\},$$

as desired. □

C Unrestricted Bound Functions

In [**?**], to prove termination, we used the LOOP III rule without the stipulation that $var(t) \subseteq var(B) \cup var(S)$. We call the rules LOOP II,III, IV without this stipulation *unrestricted*. We now prove the following result.

Theorem 4.
The unrestricted rules LOOP II, III, and IV, are admissible in the proof system II, III, and IV, respectively.

We begin by formulating a crucial lemma. Given a list of distinct variables $\bar{x}$ and a list of terms $\bar{t}$ of the same length, we denote by $[\bar{x} := \bar{t}]$ the *simultaneous substitution* of each variable in $\bar{x}$ by the corresponding term in $\bar{t}$. The simultaneous substitution applied to a term t or an assertion p, written as $t[\bar{x} := \bar{z}]$ (respectively, $p[\bar{x} := \bar{z}]$), is defined in the usual way by properly taking care of the bound variables (see, e.g., [**?**, Subsection 2.7]).

Below, given a list of distinct constants $\bar{c}$ and a list of terms $\bar{t}$ of the same length, we denote by $[\bar{c} := \bar{t}]$ the *simultaneous replacement* of each constant in $\bar{c}$ by the corresponding term in $\bar{t}$, defined in the expected way.

The following lemma formalizes the intuition that in the correctness proofs variables that are not used in the analyzed program play the same role as fresh constants. The first item concerns a simultaneous substitution, while the second one deals with a simultaneous replacement.

Lemma 6. *Let PR be one of the proof systems II, III or IV. Suppose that* $\vdash_{PR} \{p\}\ S\ \{q\}$.

Assume that $\bar{x}$ *and* $\bar{c}$ *are respectively a list of distinct variables and a list of distinct constants of the same length, none of which appears in* S. *Then*

(i) $\vdash_{PR} \{p[\bar{x} := \bar{c}]\}\ S\ \{q[\bar{x} := \bar{c}]\}$,
(ii) $\vdash_{PR} \{p[\bar{c} := \bar{x}]\}\ S\ \{q[\bar{c} := \bar{x}]\}$.

Proof. (Sketch)
The proof proceeds by induction on the structure of S and is straightforward. We illustrate it by presenting just one case for (ii), when $S \equiv$ **while** B **do** S_0 **od** and the proof system II. Let

$$\vdash_{\mathrm{II}} \{p_0 \wedge B \wedge t = z\}\ S_0\ \{p_0 \wedge t < z\},$$

for some assertion p_0, bound function t and variable z such that the implications $p \to p_0$, $p_0 \to t \geq 0$ and $(p_0 \wedge \neg B) \to q$ hold.

Denote $p[\bar{c} := \bar{x}]$ by p' and similarly with p_0, B, q and t. By the induction hypothesis the above statement holds when we replace each assertion or term by its primed version and the primed versions of all three implications hold by logic. Note that by the assumption about the constants in $\bar{c}$ we have $B' \equiv B$. Hence by the LOOP II rule

$$\vdash_{\mathrm{II}} \{p_0'\}\ \textbf{while}\ B\ \textbf{do}\ S\ \textbf{od}\ \{p_0' \wedge \neg B\},$$

so by the CONSEQUENCE rule

$$\vdash_{\mathrm{II}} \{p'\}\ \textbf{while}\ B\ \textbf{do}\ S\ \textbf{od}\ \{q'\},$$

as desired.

Below we shall use this lemma twice, first to simultaneously substitute specific variables by (fresh) constants and then to replace these constants by the original variables.

Proof of Theorem ??. We treat the case of the unrestricted LOOP II rule. Suppose

$$\vdash_{\mathrm{II}} \{p \wedge B \wedge t = z\}\ S\ \{p \wedge t < z\},$$

and that $p \to t \geq 0$ holds, where t is an integer expression such that $var(t) \not\subseteq var(B) \cup var(S)$ and z is an integer variable that does not appear in p, B, S, t or q.

Let $\bar{x}$ be a list formed by the variables from $var(t)$ that are not elements of $var(B) \cup var(S)$ and let $\bar{c}$ be a list of distinct constants of the same length as $\bar{x}$, none of which appears in p, B, t or S. Denote $p[\bar{x} := \bar{c}]$ by p' and $t[\bar{x} := \bar{c}]$ by t'.

Then we have by Lemma **??**(i)

$$\vdash_{\mathrm{II}} \{p' \wedge B \wedge t' = z\}\ S\ \{p' \wedge t' < z\},$$

and

$$p' \to t' \geq 0.$$

Note that $var(t') \subseteq var(B) \cup var(S)$. So by an application of the LOOP II rule we obtain

$$\vdash_{\mathrm{II}} \{p'\}\ \textbf{while}\ B\ \textbf{do}\ S\ \textbf{od}\ \{p' \wedge \neg\ B\},$$

from which we get by Lemma **??**(ii)

$$\vdash_{\mathrm{II}} \{p\}\ \mathbf{while}\ B\ \mathbf{do}\ S\ \mathbf{od}\ \{p \wedge \neg\, B\},$$

since $p'[\bar{c} := \bar{x}] \equiv p$ and $B[\bar{c} := \bar{x}] \equiv B$. $\square$