7	A	В	C																	
					ļļ.								 						ļ	
	a,	61	9													~				
	a.	62	C																	
		14	7										 							
	,												 						ļ	
DLAB	Cal	الم	B					4	•	_										
- MB	7.1.1		U					H	Ae	ا).	-)			A	7		C	•		
		a	- 64			-	\dashv						a	- 1			C			
												1						 		
		a	₩,	2_									a	Л			C	2		
	ļ			ļ										•					-	
	<u> </u>				-	-														
死	7 4g (r)	м 2	Tan Co	-1		A		Ω			9									
	י עצר		ME (4	- 4	4		<u>e</u>			1		•						ļ	
				-	(2,		6	1	-	2		 					ļ		
											•								ļ	•
						9		6	1	-	G		 					ļ	ļ	
				-		ay		6	•	-	P		 						ļ	
							1		·	1	,	•	 							•
						a		1	2		e	L	 							
	ļ												 					ļ	ļ	
乐	g(r)	NI	~ / C C	1 -	4	/							 					ļ		
/ YA	タレノ	17. 00	ACU'	/ 7															ļ	
~~/·		0		21	A			a					 					ļ	ļ	
10	e),	151,	K ₂ .	$\geq \ell$	Z,	K	4	ھِ	30	U			 							-
										1			 							•
Hp.	(r)	DE Y	0. (r)	: 1								 							-
	++			j	1								 							
T_{α}	(٢)	MI	2 (0	1 C	Z Y	+							 						ļ	٠
174	•		16 C.	17									 							
	 <u> </u> 				-		a						 						ļ	
7	S Ty	(0)	M	丸。	10		?						 					ļi		-
•		401/		14	 	1	9						 							-
				ļļ					7/				 			ļ		ļ		
				-									 ******			ļ		ļ	ļ	
													 		ļ			<u> </u>		
					ļļ.,						ļ		 			ļ	ļ	ļ	ļ	

www.up.pt UPORTO

$r(u) = \pi_{R_1}(r) \bowtie \pi_{R_2}(r), \mathcal{R}_1 R_2 = \mathcal{U}$ $t \in \mathfrak{I} \Rightarrow$ $t \in \mathcal{R}_1 \ni \in \pi_{R_1}(r) \land t \in \mathcal{R}_2 \vdash \in \pi_{R_2}(r) \Rightarrow$ $t \in \mathcal{R}_1 \downarrow \in \mathcal{R}_2 \supset \in \pi_{R_1}(r) \bowtie \pi_{R_2}(r) \Rightarrow$ $t \in \mathcal{R}_{R_1}(r) \bowtie \pi_{R_2}(r) \subseteq r$ $r \bowtie \mathcal{R}_1 \subseteq \mathcal{R}_2 \subseteq \mathcal{R}_3 \subseteq \mathcal{R}_4 \subseteq$	7/1	<i>)</i> .	S (ton	(~ \	r	M	#			' ,	. 9		<u> </u>	12	. 6	2.		- 7) [<u> </u>	
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$		7/		K	1	• /				K	2	. '	/ 5				1	2	ļ	*	7			ļ	
$ \begin{array}{cccccccccccccccccccccccccccccccccccc$	4c	4	=					-											ļ			ļ	.		
$\begin{array}{cccccccccccccccccccccccccccccccccccc$,				,	ــــــــــــــــــــــــــــــــــــــ	,					- 4) .					4			ļ			
$\begin{array}{cccccccccccccccccccccccccccccccccccc$		t	しバ	yJ	E	: 1	42	L	()							યૃ	1	\in	12	12	ر (1	/	=	€
$t \in \mathcal{T}_{R_{1}}(r) \bowtie \mathcal{T}_{R_{2}}(r) \bigcirc$ $\mathcal{T}_{R_{1}}(r) \bowtie \mathcal{T}_{R_{2}}(r) \subseteq r ?$ $r \triangleq B \subseteq c$ $a_{1} \notin c_{1}$ $a_{1} \notin c_{1}$ $a_{1} \notin c_{2}$ $a_{1} \notin c_{1}$ $a_{1} \notin c_{2}$ $a_{1} \notin c_{1}$ $a_{1} \notin c_{2}$		1			T T				7		1					1			7						
$t \in \mathcal{T}_{R_{1}}(r) \bowtie \mathcal{T}_{R_{2}}(r) \bigcirc$ $\mathcal{T}_{R_{1}}(r) \bowtie \mathcal{T}_{R_{2}}(r) \subseteq r ?$ $r \triangleq B \subseteq c$ $a_{1} \notin c_{1}$ $a_{1} \notin c_{1}$ $a_{1} \notin c_{2}$ $a_{1} \notin c_{1}$ $a_{1} \notin c_{2}$ $a_{1} \notin c_{1}$ $a_{1} \notin c_{2}$		t	U!	41	₹		R_2	J		€	•	ル	124	(~)	t	4	ħ	12,	(r)	_	>	
$T_{P_{\alpha}}(r) \times t_{P_{\alpha}}(r) \leq r$? $r \wedge A \otimes C \wedge C$		1		7	1			ليب																Ţ	
$T_{P_{\alpha}}(r) \times t_{P_{\alpha}}(r) \leq r$? $r \wedge A \otimes C \wedge C$					5												-		ļ				ļ	ļ	
$T_{P_{\alpha}}(r) \bowtie t_{P_{\alpha}}(r) \leq r$? $r \triangleq B \cdot C$ $q \in G \cdot G$ $q \in G $		Ł	- e	- 9	47	. ((1)	M	d	TL,) _	ſr	• }			1	-					ļ	ļ	ļ	
TOAB(T) M TAC(T) A B C TAB(T) M TAC(T) A B C An 6, 42 TAB(T) M TAC(T) A B C					17	•					-2		1			1			ļ					ļ	
TOAB(T) M TAC(T) A B C TAB(T) M TAC(T) A B C An 6, 42 TAB(T) M TAC(T) A B C			-					-									-		-				ļ	ļ	ļ
TOAB(T) M TAC(T) A B C TAB(T) M TAC(T) A B C An 6, 42 TAB(T) M TAC(T) A B C	T.	1	(1)	N		大	a		~)	<	3	Y	`	?)		1	-	ļ				ļ	ļ	
TAB(C) A B JAC(1) A C A1 61 C2 TAB(C) M TAC(C) A B C A1 61 C2		` `					2		7																
TAB(C) A B JAC(1) A C A1 61 C2 TAB(C) M TAC(C) A B C A1 61 C2	r		4	•	5	3		C									-								ļ
TAB(C) A B JAC(1) A C A1 61 C2 TAB(C) M TAC(C) A B C A1 61 C2					-	,																ļ			
TAB(C) A B JAC(1) A C A1 61 C2 TAB(C) M TAC(C) A B C A1 61 C2			CL,		E	1	(-	ļ			ļ	ļ		ļ
TAB(C) A B JAC(1) A C A1 61 C2 TAB(C) M TAC(C) A B C A1 61 C2			a.		6			<u> </u>										+							ļ
TAB(C) M TAC(O) A B C			I					2	,								-	-	ļ						
TAB(C) M TAC(O) A B C			-					ļ									-	-	ļ			ļ		ļ	
TAB(C) M TAC(O) A B C	4																		ļ						
TAB(T) M TAC(T) A B C	JOAB	(\cap)			A		-8								H	4	6	(1		_	4	ļ	(_	
12, C2 12, C2 13, C2 14, C3 14, C4 14, C4 14, C4			1				_																		******
Ang (r) M Dac (r) A C C		ļ		(41		U	1												Q	1		4	1	
Ang (r) M Dac (r) A C C								ļ												a				7	ļ
191 4 G																								2_	•
191 4 G			-	r.			.,										-		ļ						
191 4 G	h.	2 (/	')	M	I		7	n			Δ			ß				ļ							
	- 1.8					- 1	<u> </u>				-	•		7		7	-	•	ļ						
			 			.		1			a	1	4	6.		C.	-						L		
								ļ	1		^	•				1									
RAB(C) A Tac(C) = r			 					ļ			U	4	•	6	1	Ç	2	-					ļ		
MAB(r) A Rec(r) = r																	ļ	ļ							
	MAI	3 (r) D	4.	The same		() :	-	r	•							-							

www.up.pt U.PORTO

```
Tixy (r) & Tix(u)(r)=
          Fxy(r) N Fx(ary)(r) Sr
(tt)(te Txy(r) + Tx(u,y)(r); > ter)
 te Txy (r) M Tx (n,y)(r) >
 t[xy] e Toxy (r) A t[x(u)y)] e tx(u)y)(r)=
(34 er)(4 Cx y J = + Cx y J) 1
(3 fzer)(fz[x(u)y)] = t[x(u)y)] =
tier a ticxyj= tcxyj a
t2 er 1 t2[x(uxy)] = t[x(uxy)]
t_1(x) = t(x)
              Λ
                       SCX7= SCX7=4CX7
(2[x] = t [x]
rfxyy
                          4CYJ=6CYJ=6CY
れて×Jニセン×J=セン×J
 6 [4] = t[4]
                              t<sub>2</sub> = t ) ⇒ ter
t<sub>2</sub> ∈ ( ) ===
 tz [x(uxy)] = t[x(uxy)]
  ouver cyrap:
 They (r) HEX(nexy)(r)=r
```

U.PORTO

WW.up.ot

```
Dow:
                   Axx (r) H Ex(exy)(r)=
FXY (r) M FX (vy) (r) S T
t∈ txy(r) & tx(u)y)(r) >
 texyj € texy(r) A tex(ary)J € tex(ary)(r) =>
(746r)(th[xy7=+[xy3) A
(7 t2 er) ( t2 [x(a)y)] = t2x(a,y)])
 tiev a ticxyj = tcxyj a
 t2 Gr 1 t2[x(u)]] = t[x(u)y)] =
 ticx] = tecx] = tcx]
 ケドメララソ
 (Juer) ( MIXYI = 4, [XY] A
           ~ [x(~\y)] = & tx:(~\y)])
 ueramexyj= f, cxyja
         u[x(u,y)] = t2[x(ne(y)]
 4(Cxy) = t, Cxy) ) = 4(xy) = t(xy)
 f[x y] = ti [xy]
 u(x(u(y)) = f_2(x(u(y)))
 t [x(4)y)] = fz[x(u)y)]
```

u[xy]= t[xy] $M[X(u \mid y)] = t[X(u \mid y)]$ (=) xxx(r) & xx(uxy)(r)=r (Yu, ver)(u(x) = v(x) => (Fter) (Ntxy) = Ecxy1 n $\mathscr{P}[\times(u \mid y)] = t \, \mathbb{I} \times (u \mid y) \, \mathcal{I}))$ u,ver, MCX] = V [X] UET > MIXYJEILY(1) N[x(uxy)] & fix(uxy)(r) v∈r⇒ u[x]=v[x] NE(XY) O [X(M)Y)) E STXX (r) M FX(M)Y)(r)2 Txy(r) A tx(uxy)(r) = M(XX) or (X (M) X)] er ter $t[XY] = \alpha[XY]$ t [x(axy)] = v[x(axy)]

U. PORTO

www.up.ot

FABCD		W(AB,	Bc, cd)
a, 8, e, d,					
Qu 82 62 d2					
TAB(+) A 8	. A _{Bc}	(-)	B	C	
4, 4,	austropy and the		THE RESERVE THE PERSON NAMED IN	CA	
a, 6,			6,	4	
			6,	. 12	
Twenty CD					
G da					
(C, d					
Ray(r) M Juga(r)	A 8	C			
	a 61	A STATE OF THE PARTY OF THE PAR			
	an bi		1 1 1		
	1 a2 62	, c ₂	•		
ins (~) H Froc (~) b	4 Fac (1)	A	- B	C D	
			A STATE OF THE PARTY OF THE PAR	en dy	
				c, d,	
				c _k d	
(, U UX , = U					
FX (X1,, XK)	akko	JUX.	(r).k	147	

OL