

# Remember to record your attendance

1. Download the University of Bristol app
2. Sign in with your university user-name and password
3. Select 'Record attendance'  
(or tap on the notification)
4. If prompted, enable location services
5. Tap 'Check in' for this class, then 'Finish'
6. Your attendance confirmation will be displayed

Or go to [check-in.bristol.ac.uk](http://check-in.bristol.ac.uk) for web check-in

For more information or help, search for 'record attendance' on [www.bristol.ac.uk](http://www.bristol.ac.uk)




Apple App Store




Google Play Store

# PHYS40071: Complex Disordered Matter




Nigel Wilding, Francesco Turci and Adrian Barnes



# Welcome!

# Course structure

TB 1 | 20 Credits | 3 Lectures per week + Problem Class



## 📌 **Unifying Concepts** (Weeks 1-5: Nigel Wilding)

- Phase transitions and critical phenomena
- Metastability and dynamics
- Stochastic processes and fluctuations

## 🧩 **Complex Disordered Systems** (Weeks 7-11: Francesco Turci)

- Colloids and interactions
- Polymers & Surfactants
- Liquid Crystals, Gels, Glasses
- Active Matter

## 🔬 **Experimental Systems & Techniques** (Weeks 5 and 11: Adrian Barnes)

- Calorimetry
- Microscopy
- Diffraction

**Teaching Block 1**

<b>Weeks</b>	<b>Commencing</b>	<b>Section</b>	<b>Lecturer</b>	<b>Assessment</b>
1 - 4	22nd Sep	Unifying Concepts	Prof. Nigel Wilding	Coursework 1: Released Monday 13th October (Week 4) 12:30, due Monday 27th October, 09:30 (Week 6). Marks and feedback returned by Fri 14th November.
5	20th Oct	Unifying Concept and Experimental Techniques	Prof. Nigel Wilding and Dr. Adrian Barnes	
6	27st Oct	<i>Consolidation Week</i>	-	
7 - 10	3rd Nov	Complex Disordered Systems	Dr. Francesco Turci	Coursework 2: Released Thursday 13th Nov 12:30 (Week 8), due Thursday 27th November, 09:30. Marks and feedback returned Friday 12th December.
11	1st Dec	Complex Disordered Systems and Experimental Techniques	Dr. Francesco Turci and Dr. Adrian Barnes	
12	8th Dec	<i>Revision Week</i>	-	
	<b>15th Dec</b>	<b>Assessment Period (1 week)</b>	-	<b>1.5 hour examination</b>

Each teaching week, we shall meet as follows:

Tuesdays 10:00, Room: 3.34 (Physics)	Lecture
Wednesday 11:00, Room: 3.21 (Berry theatre)	Lecture
Thursdays 12:00, Room: Fry Building G.13	Problems class
Fridays 14:00, Room: 3.21 (Berry theatre)	Lecture

# Assessment Breakdown

💻 **Computational investigations** - 1 Assignment (30%)


🤔 **Problem set** (20%)

📖 **90 minute exam (Dec)** – Covering all lecture material (50%)

📌 **Total: 100% Assessment** – Equally balanced between coursework and final examination.

As this is a new course, you will be provided with a mock examination paper

# Delivery and format (Unifying Concepts)




- Detailed e-notes (see Blackboard) can be viewed on a variety of devices.
- 'Traditional' lectures (Tues, Wed, Fri) in which I use slides to summarise and explain the lecture content. Questions welcome (within reason...)
- Try to read ahead in the notes, then come to lectures, listen to my explanations and then reread the notes.



- Rewriting the notes or slides to express your own understanding, or annotating a pdf copy can help wire the material into your own way of thinking.
- Problem class (Thurs) where you can try problem sets and seek help. I will go over some problems with the class. **No classes week 6.**

# 1. Introduction to phase behaviour


- A wide variety of physical systems undergo rearrangements of their internal constituents in response to changes in thermodynamic conditions.
- Two classic examples of systems displaying such phase transitions are the ferromagnet and fluid systems.



- As the temperature  $T$  of a ferromagnet is increased, its magnetic moment is observed to decrease smoothly, until at a certain **critical temperature**, it vanishes altogether

We define the magnetisation to be the **order parameter** of this phase transition.

- Similarly, we can induce a change of state from liquid to gas in a fluid simply by raising the temperature.



- Typically the liquid-gas transition is abrupt. However, abruptness can be reduced by applying pressure.
- At one particular  $p, T$  the jump in density at the transition vanishes. These conditions correspond to the critical point of the fluid.

- For  $H_2O$ :  $T_c = 374$  K,  $P_c = 218$  atm
- For  $CO_2$ :  $T_c = 304$  K,  $P_c = 73$  atm
- Density difference  $\rho_{liquid} - \rho_{gas}$  is the order parameter for the liquid-gas phase transition.

- Close to its critical point, a system exhibits a variety of remarkable effects known as **critical phenomena**
- Examples are the divergence (infinite values) of thermal response functions such as the specific heat and the fluid compressibility or magnetic susceptibility
- Origin of the singularities in these quantities tracable to large-length-scale co-operative effects between the microscopic constituents of the system.

- Illustration:  $CO_2$  in a pressurised container at  $P_c$ .
- As approach  $T_c = 31^\circ C$  from above, critical opalescence occurs.
- For  $T < T_c$  a liquid-gas meniscus forms



- Understanding the physics of critical points is important because often one observes **quantitatively identical** critical phenomena in a range of apparently quite disparate physical systems.
- This implies a profound underlying similarity among physical systems at criticality, regardless of many aspects of their distinctive microscopic nature.
- These ideas have found formal expression in the celebrated **universality hypothesis** (see later)

## 2. Background concepts

- Let us denote the order parameter as  $Q$ .
- $Q$  provides a quantitative measure of the difference between the phases coalescing at the critical point:
- Reminder: For the fluid,  $Q = \rho_{liq} - \rho_{gas}$ . For the ferromagnet  $Q = m$ , the magnetisation.
  
- But why should a system exhibit a phase transition at all?
- Statistical Mechanics provides the answer!

- Probability  $p_a$  that a physical system at temperature  $T$  will have a particular microscopic arrangement ('configuration' or 'state'), labelled  $a$ , of energy  $E_a$  is

$$p_a = \frac{1}{Z} e^{-E_a/k_B T}$$

- Prefactor  $Z^{-1}$  is a  $T$ -dependent constant - the partition function (recall year 2 thermal physics). Since the system must always have some specific arrangement, the sum of the probabilities  $p_a$  must be unity, implying that

$$Z = \sum_a e^{-E_a/k_B T}$$

where the sum extends over all possible microscopic arrangements.

- Expectation (ie. average) value of an observable  $O$  is given by averaging  $O$  over all the arrangements  $a$ , weighting each contribution by  $p_a$

$$\bar{O} \equiv \langle O \rangle = \frac{1}{Z} \sum_a O_a e^{-E_a/k_B T}$$

- The order parameter is similarly a thermal average over configurations:

$$Q = \frac{1}{Z} \sum_a Q_a e^{-E_a/k_B T} \quad \dagger$$

- Consider the ferromagnetic case  $Q = m$ . For  $T$  very small, system will be overwhelmingly likely to be in its minimum energy arrangement (ground state) having magnetisation +1, or -1.
- For  $T$  large, enhanced prob. of ground state arrangements is insufficient to offset the fact that the sum in eq.  $\dagger$  contains a vastly greater number of arrangements in which  $Q_a$  has some intermediate value.
- In fact arrangements which have essentially zero magnetisation (equal populations of up and down spins) are by far the most numerous
- At high temperature, these disordered arrangements dominate the sum in eq.  $\dagger$  and  $Q \approx 0$ .

- Thus  $T$  dependence of  $Q$  results from a competition between energy-of-arrangements weighting (or simply 'energy') and the 'number of arrangements' weighting (or 'entropy').
- The critical point is that  $T$  at which the system is forced to choose amongst a number of macroscopically different sets of microscopic arrangements i.e. the finite  $Q$  arrangements and the zero  $Q$  arrangements.
- Partition function provides bridge between stat. mech and thermodynamics via

$$F = -k_B T \ln Z$$

where  $F$  is the Helmholtz free energy (see year 2, and preparatory reading).

- All thermodynamic observables, eg. Order parameter  $Q$ , specific heat  $C_H$ , susceptibility  $\chi$  and compressibility  $\kappa$ , are obtainable as appropriate derivatives of the free energy. Eg:

$$\begin{array}{c}
 Z = \sum_a e^{-E_a/k_B T} \\
 \downarrow \\
 F = -k_B T \ln Z \\
 \swarrow \qquad \searrow \\
 \bar{E} = - \left( \frac{\partial(\beta F)}{\partial \beta} \right)_H \qquad \bar{M} = - \left( \frac{\partial F}{\partial H} \right)_T \quad (M = mN) \\
 \downarrow \qquad \qquad \qquad \downarrow \\
 C_H = \left( \frac{\partial \bar{E}}{\partial T} \right)_H \qquad \chi_T = \left( \frac{\partial m}{\partial H} \right)_T
 \end{array}$$

(Here  $\beta \equiv (k_B T)^{-1}$  )

# Correlations

- Consider first **spatial** correlations.
- Two-point correlation function measures statistical relation between fluctuations at two spatial points
- For scalar field  $\phi(\vec{R})$  eg. local density or local magnetisation, can define

$$C(r) = \langle \phi(\vec{R})\phi(\vec{R} + \vec{r}) \rangle - \langle \phi(\vec{R}) \rangle^2 \quad r = |\vec{r}| \text{ is spatial separation}$$

and the average is over configurations

Including this term gives a quantity that decays to zero


- In homogeneous/isotropic systems,  $C(r)$  depends only on  $r$  and not the reference point  $\vec{R}$

# Behaviour of spatial correlations

- Correlations can be short ranged or long ranged:


- Short-range:  $C(r) \sim e^{-r/\xi}$ , fast decay
- Long-range (criticality): power-law decay  
 $C(r) \sim r^{-x}$

- **Correlation length  $\xi$**  sets spatial scale.




## Example: pair correlation function of a fluid (a.k.a radial distribution function)

- For a fluid  $\phi(\vec{R}) = \rho(\vec{R})$ , the local number density
- $g(r) = \text{probability of finding particle at distance } r = |\vec{r}| \text{ relative to ideal gas.}$
- It is related to density correlations via:  $g(r) = \frac{1}{\rho^2} \langle \rho(0)\rho(r) \rangle = 1 + C(r)/\rho^2$




- At large  $r$  correlation die out, ie.  
$$g(r) \rightarrow 1 \quad \text{as } r \rightarrow \infty$$



- Peaks in  $g(r)$  indicate ordering (e.g. shells in liquids).

# Structure Factor $S(k)$

- Fourier transform of correlations:  $S(k) = \int d^3r e^{i\vec{k}\cdot\vec{r}} C(r)$   
 $k = \frac{2\pi}{\lambda}$  is the scattering wave vector (often also written as  $q$ )
- Measured in scattering experiments (see lectures by A. Barnes). Connects real-space correlations with momentum-space
- Peaks in  $S(k)$  = characteristic length scales.
- Near criticality: divergence at small  $k$  (long-range fluctuations)
- For short ranged correlations, decay of  $S(k)$  has a Lorentzian form.
- For long ranged correlations, decay has a power law form (see notes)



# Temporal correlations


- Consider a thermodynamic variable  $x$  with zero mean that fluctuates over time.
- Temporal correlations are characterized by the two-time correlation function:

$$\langle x(\tau)x(\tau + t) \rangle$$

- In **equilibrium**:  $\langle x(\tau)x(\tau + t) \rangle = M_{xx}(t)$  i.e only depends on  $t$  not on  $\tau$
- Typically (away from criticality):

$$M_{xx}(t) \sim \exp(-t/t_c)$$

ie. memory of fluctuations fades exponentially with correlation time  $t_c$



# 3. The approach to criticality

- Approach to criticality is characterised by the divergence of various thermodynamic observables.
- Eg. in a ferromagnet near  $T_c$ ,  $C_H$  and  $\chi_T$  are singular functions, diverging as some power of the **reduced temperature**  $t \equiv (T - T_c)/T_c$

$$\chi \equiv \frac{\partial m}{\partial h} \propto t^{-\gamma} \quad C_H \equiv \frac{\partial E}{\partial T} \propto t^{-\alpha}$$

- Recall that the correlation length  $\xi$ , measuring the distance over which fluctuations of the magnetic moments are correlated. This diverges with an exponent  $\nu$ .

$$\xi \propto t^{-\nu} \quad (T > T_c, H = 0)$$


- Similar power law behaviour is found for the order parameter  $Q$  which vanishes in a singular fashion (it has infinite gradient) as  $T \rightarrow T_c^-$

$$m \propto t^\beta \quad (T < T_c, H = 0)$$

And as a function of magnetic field:

$$m \propto h^{1/\delta} \quad (T = T_c, H > 0)$$

with  $h \equiv (H - H_c)/H_c$  the reduced magnetic field.



- $\gamma, \alpha, \nu, \beta$  are known as **critical exponents**. They control the rate at which the associated observables change on the approach to criticality.

- Remarkably, similar power laws occurs in many qualitatively distinct systems near their critical point (eg, fluid, polymer solutions, magnets, electrolytes, traffic jams, sandpiles, financial markets).
- To obtain the corresponding power law relationships simply substitute the analogous thermodynamic quantities in to the above equations. eg.

$$\rho_{liq} - \rho_{gas} \sim t^\beta; \quad \kappa \sim t^{-\gamma}$$

- Even more remarkable is the experimental observation that the values of the critical exponents for a whole range of fluids and magnets (and indeed many other systems with critical points) are **identical**.
- This is the celebrated phenomenon of universality. It implies a deep similarity between systems at their critical points.


# 4. Ising model

- We can use simple models to probe the properties of the critical region.
- Simplest is the 2d spin- $\frac{1}{2}$  Ising model, which comprises a lattice of  $N$  magnetic moments or 'spins' on an infinite plane.
- Each spin can take two values, +1 ('up' spins) or -1 ('down' spins) and interacts with its nearest neighbours via the Hamiltonian

$$\mathcal{H}_I = - J \sum_{\langle ij \rangle} s_i s_j - H \sum_i s_i \quad J > 0$$

- The order parameter is simply the average magnetisation:  $m = \frac{1}{N} \langle \sum_i s_i \rangle$


- At low temperatures for which there is little thermal disorder, there is a preponderance of aligned spins and hence a net spontaneous magnetic moment  $|m| \approx 1$
- As  $T$  is raised, thermal disorder increases until at  $T_c$ , entropy drives the system through a continuous phase transition to a disordered spin arrangement with  $|m| = 0$ .
- These trends are visible in configurational snapshots from computer simulations of the 2D Ising model.



$T = 1.2T_c$



$T = T_c$




$T = 0.95T_c$

- Although each spin interacts only with its nearest neighbours, the phase transition occurs due to cooperative effects among a large number of spins.
- In the neighbourhood of the transition temperature these cooperative effects engender fluctuations that can extend over all length-scales from the lattice spacing up to the correlation length.

- Despite its simplicity, critical point universality implies that critical exponents of Ising model are same as those of real magnets.
- Ising model therefore provides a simple, yet quantitatively accurate representation of the critical properties of a whole range of real magnetic (and indeed fluid) systems.
- This universal feature of the model is largely responsible for its ubiquity in the field of critical phenomena.

# Exact Solutions: the 1D Ising chain

- Why is the 2D Ising model the simplest to exhibit a phase transition? What happens in 1D?
- In fact there is no phase transition in 1D for  $T > 0$ .
- Consider the ground state of a 1D Ising chain and a state with various "domain walls" dividing spin-up and spin-down regions



- Transform from a spin representation to a domain wall representation.

- Domain walls can occur on the bonds of the lattice of which there are  $N - 1$ . If a wall is present, the energy cost is  $\Delta = 2J$ .
- A configuration can be specified by stating whether or not there is a domain wall on each bond.
- Presence of a domain wall on one bond doesn't affect likelihood of domain wall on neighbouring bond (as long as  $H = 0$ ). Thus domain walls are **independent**
- Partition functions from independent contributions multiply.
- Hence the partition function of system is  $Z = Z_1^{N-1}$ , where for a single domain wall,

$$Z_1 = e^{\beta J} + e^{\beta(J-\Delta)} = e^{\beta J}(1 + e^{-\beta\Delta}) \quad \beta \equiv \frac{1}{k_B T}$$

So free energy density

$$\beta f \equiv \beta F/(N - 1) = -\ln Z_1 = -\beta J - \ln(1 + e^{-\beta\Delta})$$

- Second term arises from the entropy of the domain wall population and since it is negative for all  $T > 0$ , the free energy is lowered by having domain walls, ie the system is always disordered.

# More general 1D spin systems

- For a 1-d assembly of  $N$  spins each having  $m$  discrete energy states, and in the presence of a magnetic field  $H$ , possible to get free energy via the **transfer matrix method**.
- Let us start by assuming that the assembly has cyclic boundary conditions, then the total energy of configuration  $\{s\}$  is

$$\begin{aligned}\mathcal{H}(\{s\}) &= - \sum_{i=1}^N (Js_i s_{i+1} + Hs_i) \\ &= - \sum_{i=1}^N (Js_i s_{i+1} + H(s_i + s_{i+1})/2) \\ &= \sum_{i=1}^N E(s_i, s_{i+1})\end{aligned}$$

where we have defined  $E(s_i, s_{i+1}) = - Js_i s_{i+1} - H(s_i + s_{i+1})/2$ .

Partition function may be written

$$\begin{aligned}
 Z_N &= \sum_{\{s\}} \exp(-\beta \mathcal{H}(\{s\})) \\
 &= \sum_{\{s\}} \exp(-\beta [E(s_1, s_2) + E(s_2, s_3) + \dots + E(s_N, s_1)]) \\
 &= \sum_{\{s\}} \exp(-\beta E(s_1, s_2)) \exp(-\beta E(s_2, s_3)) \dots \exp(-\beta E(s_N, s_1)) \\
 &= \sum_{i,j,\dots,l=1}^m V_{ij} V_{jk} \dots V_{li} \quad \ddagger
 \end{aligned}$$

where the  $V_{ij} = \exp(-\beta E_{ij})$  are elements of an  $m \times m$  matrix  $\mathbf{V}$ , known as the transfer matrix.

- Transpires that the sum over the product of matrix elements in  $\ddagger$  is the trace of  $\mathbf{V}^N$ , given by the sum of its eigenvalues:-

$$Z_N = \lambda_1^N + \lambda_2^N + \dots + \lambda_m^N$$

- As  $N \rightarrow \infty$ , largest eigenvalue  $\lambda_1$  dominates since  $(\lambda_2/\lambda_1)^N$  vanishes. Consequently  $Z_N = \lambda_1^N$
- Specializing to the case of the simple Ising model, the transfer matrix takes the form

$$\mathbf{V}(H) = \begin{pmatrix} e^{\beta(J+H)} & e^{-\beta J} \\ e^{-\beta J} & e^{\beta(J-H)} \end{pmatrix}$$

- This matrix has two eigenvalues which can be readily calculated in the usual fashion. They are

$$\lambda_{\pm} = e^{\beta J} \cosh(\beta H) \pm \sqrt{e^{2\beta J} \sinh^2 \beta H + e^{-2\beta J}}$$

- Hence the free energy per spin  $f = -k_B T \ln \lambda_+$  is

$$f = -k_B T \ln \left[ e^{\beta J} \cosh(\beta H) + \sqrt{e^{2\beta J} \sinh^2 \beta H + e^{-2\beta J}} \right]$$

# 5. Mean field theory

- The critical behaviour of most model systems cannot be found analytically.
- A few exceptions: eg. 2D Ising model (but not the 3D) have been solved  
 $(\beta = \frac{1}{8}, \nu = 1, \gamma = \frac{7}{4}, T_c = -2J/\ln(\sqrt{2} - 1) \approx 2.269J)$
- But such solutions provide little insight into the essential nature of criticality.
- When an exact solution is elusive, can try to make simplifying assumptions to calculate critical behaviour. Mean field theory is such an approximation scheme.

- Look for a mean field expression for the free energy of the Ising model. Write

$$s_i = \langle s_i \rangle + (s_i - \langle s_i \rangle) = m + (s_i - m) = m + \delta s_i$$

- Then

$$\begin{aligned} \mathcal{H}_I &= -J \sum_{\langle i,j \rangle} [m + (s_i - m)][m + (s_j - m)] - H \sum_i s_i \\ &= -J \sum_{\langle i,j \rangle} [m^2 + m(s_i - m) + m(s_j - m) + \delta s_i \delta s_j] - H \sum_i s_i \\ &= -J \sum_i (qms_i - qm^2/2) - H \sum_i s_i - J \sum_{\langle i,j \rangle} \delta s_i \delta s_j \end{aligned}$$

where the sum  $\sum_{\langle i,j \rangle}$  over bonds of a quantity which independent of  $s_j$  is just  $q/2$  times that quantity, with  $q$  the lattice coordination.

- Now the mean field approximation is to ignore the last term giving

$$\mathcal{H}_{mf} = - \sum_i H_{mf} s_i + NqJm^2/2 \quad \text{where } H_{mf} \equiv Jqm + H$$

- It follows that the partition function is

$$Z = Z(1)^N = e^{-\beta qJm^2N/2} [2 \cosh(\beta(qJm + H))]^N$$

so the free energy is

$$F(m) = NJqm^2/2 - Nk_B T \ln[2 \cosh(\beta(qJm + H))]$$


- From which the magnetisation follows as

$$m = -\frac{1}{N} \frac{\partial F}{\partial H} = \tanh(\beta(qJm + H))$$

- To find  $m(H, T)$ , we must numerically solve this last equation self consistently.

# Spontaneous symmetry breaking

- Mean field method reveals what is happening in the Ising model near the critical temperature  $T_c$
- Plot  $\beta F(m)/N$  vs  $T$  for  $H = 0$ :
- For  $H = 0$ ,  $F(m)$  is symmetric in  $m$ . At high  $T$ , entropy dominates  $\rightarrow$  single minimum in  $F(m)$  at  $m = 0$ .



- As  $T$  is lowered, there comes a point ( $T = T_c$ ) where the curvature of  $F(m)$  at the origin changes sign; ie.

$$\frac{\partial^2 F}{\partial m^2} = 0$$

- At lower temperature: two minima at nonzero  $m = \pm m^*$ , where the equilibrium magnetisation  $m^*$  is the positive root (calculated explicitly below) of


$$m^* = \tanh(\beta J q m^*) = \tanh\left(\frac{m^* T_c}{T}\right)$$

- $m^* = 0$  which remains a root of this equation, is clearly an unstable point for  $T < T_c$  (since  $F$  has a maximum there).
- This is an example of spontaneous symmetry breaking: Pair of ferromagnetic states (spins mostly up, or spins mostly down) which - by symmetry - have the same free energy, lower than the unmagnetized state.

# Phase diagram


- The resulting zero-field magnetisation curve:

- Sudden change of behaviour at  $T_c$  (continuous phase transition).
- For  $T < T_c$ , arbitrary which of the two roots  $\pm m^*$  is chosen; typically it will be different in different parts of the sample (giving macroscopic ``magnetic domains'').



- Picture is qualitatively modified by a magnetic field  $H$ . Then there is always a finite magnetization, even for  $T \gg T_c$  (no phase transition).

- Sit below  $T_c$  with  $H > 0$  and gradually reduce  $H$  so that it becomes negative. Observe very sudden change of behaviour at  $H = 0$ : the equilibrium state jumps discontinuously from  $m = m^*$  to  $m = -m^*$ .



- This is called a **first order phase transition**.

**First order transition:** magnetisation (or similar order parameter) depends discontinuously on other variables such as  $H$  or  $T$

**Continuous transition (criticality):** Change of functional form, but no discontinuity in  $m$ ; typically, however,  $(\partial m / \partial T)_H$  (or similar) is either discontinuous, or diverges.

- We say that the phase diagram of the magnet in the  $H, T$  plane shows a line of first order phase transitions, terminating at a continuous transition, which is the critical point.

# Mean field predictions for critical exponents

- Look for a solution to  $m = \tanh(\beta J q m)$  in zero field ( $H = 0$ ) where  $m$  is small ( $\ll 1$ ). Taylor expanding the  $\tanh$  function yields

$$m = \frac{mT_c}{T} - \frac{1}{3} \left( \frac{mT_c}{T} \right)^3 + O(m^5)$$

- Then  $m = 0$  is one solution. The other solution is given by


$$m^2 = 3 \left( \frac{T}{T_c} \right)^3 \left( \frac{T_c}{T} - 1 \right)$$

- Now, for  $T$  close to  $T_c$  (i.e. small  $m$ ), and writing  $t = (T - T_c)/T_c$ , one finds

$$m^2 \simeq -3t$$

i.e.

$$\begin{aligned} m &= 0 && \text{for } T > T_c \quad \text{since otherwise } m \text{ imaginary} \\ m &= \pm \sqrt{-3t} && \text{for } T < T_c \quad \text{real} \end{aligned}$$



- In a small finite field we can make the Taylor expansion

$$m = \frac{mT_c}{T} - \frac{1}{3} \left( \frac{mT_c}{T} \right)^3 + \frac{H}{k_B T}$$

- Consider now the isothermal susceptibility

$$\begin{aligned} \chi &\equiv \left( \frac{\partial m}{\partial H} \right)_T \\ &= \frac{T_c}{T} \chi - \left( \frac{T_c}{T} \right)^3 \chi m^2 + \frac{1}{k_B T} \end{aligned}$$

- Then

$$\chi \left[ 1 - \frac{T_c}{T} + \left( \frac{T_c}{T} \right)^3 m^2 \right] = \frac{1}{k_B T}$$


- Hence near  $T_c$

$$\chi = \frac{1}{k_B T_c} \left( \frac{1}{t + m^2} \right)$$

And recalling that  $m^2 \simeq -3t$  for  $T < T_c$

$$\chi = (k_B T_c t)^{-1} \text{ for } T > T_c$$

$$\chi = (-2k_B T_c t)^{-1} \text{ for } T \leq T_c$$



where one has to take the non-zero value for  $m$  below  $T_c$  to ensure positive  $\chi$ , i.e. thermodynamic stability.


- This result implies that (within the mean field approximation) the critical exponent  $\gamma = 1$ .

# Landau theory

- Landau theory is more general type of mean field theory. Not based on a particular microscopic model. We say that it is a **coarse-grained** theory.
- Starting point is the Helmholtz free energy, which is written as a truncated power series expansion of the order parameter.
- For systems with a symmetrical order parameter (such as a ferromagnet where the energy is invariant under  $m \rightarrow -m$ ) this takes the form

$$F(m) = F_0 + a_2 m^2 + a_4 m^4$$

- Equilibrium  $m$  is that for which  $F(m)$  is minimum.
- Plots of the Landau free energy (for various  $a_2$ , with  $a_4 > 0$ ) show how it gives rise to a critical point



- Thermodynamics tells us that the system adopts the state of lowest free energy.
- Thus for  $a_2 > 0$ , the system will have  $m = 0$ , i.e. will be in the disordered (or paramagnetic) phase.
- For  $a_2 < 0$ , minimum of  $F$  occurs at two symmetric minima at  $m = \pm m_0$ , i.e., the ordered phase is the stable one.

- $a_2 = 0$  corresponds to the critical point which marks the border between the ordered and disordered phases.
- Thus clearly  $a_2$  controls the deviation from the critical temperature, i.e.  $a_2 = \tilde{a}_2 t$
- Can now attempt to calculate critical exponents. First find equilibrium magnetisation, corresponding to the minimum of the Landau free energy:

$$\frac{dF}{dm} = 2\tilde{a}_2 tm + 4a_4 m^3 = 0$$

$$\Rightarrow m \propto (-t)^{1/2}, \quad \text{so } \beta = 1/2, \text{ (mean field result).}$$

- We can also calculate the effect of a small field  $H$  if we sit at  $T_c$ . Since  $a_2 = 0$ , we have

$$F(m) = F_0 + a_4 m^4 - Hm$$

$$\frac{\partial F}{\partial m} = 0 \Rightarrow m(H, T_c) = \left( \frac{H}{4a_4} \right)^{1/3}$$

or  $H \sim m^\delta$  ie.  $\delta = 3$  which defines a second critical exponent.

- Thirdly, magnetic susceptibility at zero field

$$\chi = \left( \frac{\partial m}{\partial H} \right)_{T,V} \sim |T - T_c|^{-\gamma}$$

**Exercise:** Show that  $\gamma = 1$

- Finally heat capacity (per site or per unit volume)  $C_H$ , for  $H = 0$ :

$$C_H \sim |T - T_c|^{-\alpha}$$

**Exercise:** Show that  $\alpha = 0$

# Shortcomings of mean field theories

- In real ferromagnets, as well as in more sophisticated theories, the exponents  $\beta$  and  $\gamma$  are not the simple fraction and integers found here.
- This failure of mean field theory to predict the correct exponents is of course traceable to their neglect of correlations.

	Mean Field	$d = 1$	$d = 2$	$d = 3$
Critical temperature $k_B T/qJ$	1	0	0.5673	0.75
Order parameter exponent $\beta$	$\frac{1}{2}$	-	$\frac{1}{8}$	$0.325 \pm 0.001$
Susceptibility exponent $\gamma$	1	$\infty$	$\frac{7}{4}$	$1.24 \pm 0.001$
Correlation length exponent $\nu$	$\frac{1}{2}$	$\infty$	1	$0.63 \pm 0.001$

# 6. Static Scaling Hypothesis

- The static scaling hypothesis provides a basis for power law behaviour. Moreover, it predicts the existence of so-called **scaling phenomena** in near-critical systems.
- The hypothesis asserts that: near criticality, the free energy is a so-called generalised homogeneous function of the thermodynamic fields.
- A function of two variables  $g(u, v)$  is called a generalised homogeneous function if it has the property

$$g(\lambda^a u, \lambda^b v) = \lambda g(u, v) \quad \text{for all } \lambda$$

where the parameters  $a$  and  $b$  (known as scaling parameters) are constants

- Example functions:
- $g(u, v) = u^3 + v^2$  with  $a = 1/3, b = 1/2$
- $g(u, v) = u^4v^5$  with  $a = 1/4, b = 1/5$

- For such functions one can always implement a **change of scale**, to reduce the dependence on two variables (e.g.  $t$  and  $h$ ) to dependence on one new variable.
- The arbitrary scale factor  $\lambda$  can be chosen as  $\lambda^a = u^{-1}$  giving

$$g(u, v) = u^{1/a} g\left(1, \frac{v}{u^{b/a}}\right)$$

- Thus  $g(u, v)$  satisfies a simple power law in **one** variable, provided  $v/u^{b/a} = C$ . Note, however, that this relationship specifies neither the function  $g$  nor the parameters  $a$  and  $b$ .
- Scaling hypothesis asserts that in the critical region, the free energy  $F$  is a generalised homogeneous function of thermodynamic fields
- Thus for the ferromagnet (fields  $t$  and  $h$ ):

$$F(\lambda^a t, \lambda^b h) = \lambda F(t, h)$$

- Without loss of generality, we can set  $\lambda^a = t^{-1}$ , implying  $\lambda = t^{-1/a}$  and  $\lambda^b = t^{-b/a}$ .
- Then

$$F(t, h) = t^{1/a} F(1, t^{-b/a} h)$$

where our choice of  $\lambda$  ensures that the r.h.s is now a function of a single variable  $t^{-b/a}h$

- An expression for the magnetisation can be obtained simply by taking the field derivative of  $F$

$$m(t, h) = -t^{(1-b)/a} m(1, t^{-b/a} h)$$

- In zero applied field  $h = 0$ , this reduces to

$$m(t, 0) = (-t)^{(1-b)/a} m(1, 0)$$

where the r.h.s. is a power law in  $t$

- Can now identify the exponent  $\beta$  in terms of the scaling parameters  $a$  and  $b$ .

$$\beta = \frac{1 - b}{a}$$

- By differentiating the free energy, other relations between scaling parameters and critical exponents may be deduced.
- Such calculations (try as an exercise!) yield the results  
 $\delta = b/(1 - b)$ ,  $\gamma = (2b - 1)/a$ ,  $\alpha = (2a - 1)/a$
- Relationships between the critical exponents follow by eliminating the scaling parameters from these equations. The principal results (known as ``scaling laws'') are:-

$$\alpha + \beta(\delta + 1) = 2$$

$$\alpha + 2\beta + \gamma = 2$$

- Thus only two critical exponents need be specified, for all others to be deduced.

# Experimental Verification of Scaling


- Experiments confirm the scaling hypothesis.
- Rewriting the above expression for  $m(t, h)$  in terms of the exponents  $\beta$  and  $\delta$ , one finds

$$\frac{m(t, h)}{t^\beta} = m\left(1, \frac{h}{t^{\beta\delta}}\right)$$

where the r.h.s. is a function of the single scaled variable  $\tilde{H} \equiv t^{-\beta\delta}h(t, M)$ .

- For some magnet, measure  $m$  vs  $h$  for various fixed temperatures and construct  $m - h$  isotherms.
- Plotting the data against the scaling variables  $\tilde{M} = t^{-\beta}m(t, h)$  and  $\tilde{H} = t^{-\beta\delta}h(t, M)$  one finds scaling, i.e. all isotherms collapse onto a single curve, one for  $t > 0$ , and another for  $t < 0$ .


Magnetisation of CrBr<sub>3</sub> in the critical region plotted in scaled form



- Similar results are found using the scaled equation of state of simple fluid systems such as He<sup>3</sup> or Xe.

# Computer simulation


- Simulation widely used to study critical point phenomena,
- But computational constraints restrict one to dealing with systems of finite-size.
- Cannot access the regime of truly long ranged fluctuations that characterize the near-critical regime.
- As a consequence, the critical singularities in  $C_V, \chi_T$ , order parameter, etc. appear rounded and shifted in a simulation study of a system of linear extent  $L$ .
- The peak position does not provide an accurate estimate of the critical temperature.



- Although the degree of rounding and shifting reduces with system size, it still makes it hard access to the largest system sizes which would provide accurate estimates of critical parameters.
- To deal with this, finite-size scaling methods have been developed.
- Finite-size scaling allows extraction of bulk critical properties from simulations of finite size (see later)

# 7. Universality and Renormalisation Group Theory

- Critical region is characterised by correlated microstructure on **all** length-scales up to and including the correlation length.
- Can only be accurately characterised by a very large number of variables.
- To obtain a fuller understanding of the critical region, must take account of existence of structure on all length-scales.



# The critical point: A many length scale problem


- A near critical system can be characterised by three important length scales, namely
  - (a) The correlation length,  $\xi$ , ie the size of correlated microstructure.
  - (b) Minimum length scale  $L_{min}$ , i.e. the smallest length in the microscopics of the problem, e.g. lattice spacing of a magnet or the particle size in a fluid.
  - (c) Macroscopic size  $L_{max}$  eg. size of the system.
- The authentic critical region is defined by a window condition:

$$L_{\max} \gg \xi \gg L_{\min}$$

- This regime is hard to tackle because it is characterised by configurational structure on all scales between  $L_{min}$  and  $\xi$  (it is fractal). Moreover structure on different length scales are correlated with one another.

# Philosophy and Methodology of the RG

- Central idea: a stepwise elimination of the degrees of freedom of the system on successively larger length-scales.
- Introduce a fourth length scale  $L$ , which in contrast to the other three, characterises the **description** of the system.
- $L$  typifies the size of the smallest resolvable detail in a description of the system's microstructure.
- Ising model snapshots contain **all** details of each configuration: the resolution length  $L$  coincides with the lattice spacing i.e.  $L = L_{min}$ .
- But explicit form of the small scale microstructure is irrelevant to the behaviour of  $\xi$ . Microstructure is 'noise'.
- To eliminate it, select a larger value of  $L$ , the resolution (or 'coarse-graining') length



- There are many ways of implementing this ‘coarse-graining’ procedure.
- Adopt a simple strategy in which we divide our sample into blocks of side  $L$ , each of which contains  $L^d$  sites.
- The centres of the blocks define a lattice of points indexed by  $I = 1, 2, \dots, N/L^d$ . We associate with each block lattice point centre,  $I$ , a coarse-grained or block variable  $S_I(L)$  defined as the spatial average of the local variables it contains:

$$S_I(L) = L^{-d} \sum_i^I s_i$$


- The set of coarse grained coordinates  $\{S(L)\}$  are the basic ingredients of a picture of the system having spatial resolution of order  $L$ .

- Such a coarse graining operation is easily implemented on a computer.
- But while the underlying Ising spins can only take two possible values, the block variables  $S_I(L)$  have  $L^d + 1$  possible values.
- Thus, need a more elaborate colour convention to represent block spins.  
Adopt a grey scale.
- In presenting results implement:
  - A length scaling: the lattice spacing on each blocked lattice is scaled to that of the original lattice. Can display correspondingly larger portions of the physical system.
  - A variable scaling: adjust the scale ('contrast') of the block variable so as to match block variable spectrum the full range of grey shades.
- We shall consider the results of coarse-graining configurations typical of three different temperatures:  $T > T_c$ ,  $T = T_c$ , and  $T < T_c$

$$T > T_c$$


- Consider first a system marginally above  $T_c$ , having  $\xi \approx 6$ . Apply coarse graining with block sizes  $L = 4$  and  $L = 8$ .

$$L = 1$$




- Coarse-graining amplifies the consequences of the small deviation of  $T$  from  $T_c$ .

$$L = 4$$



- As  $L$  is increased, the ratio of the size of the largest configurational features ( $\xi$ ) to the size of the smallest ( $L$ ) is reduced.  $\xi/L$  provides a natural measure of how 'critical' is a configuration.


$$L = 8$$




- Thus coarse-graining generates a representation of the system that is effectively **less critical** the larger the  $L$  value.
- Limit is an effectively fully disordered arrangement
- When viewed on length scales  $L$  larger than  $\xi$ , the correlated microstructure is no longer apparent; each coarse-grained variable is independent.

$$T < T_c$$


- A similar trend is apparent for  $T < T_c$ .
- Again, take  $\xi \approx 6$ . Coarse-graining with  $L = 4$  and  $L = 8$  again generates representations which are effectively less critical .
- This time the coarse-graining smoothes out the microstructure which makes the order incomplete.
- The limit point of this procedure is a homogeneously ordered arrangement.



$$L = 1$$




$$L = 4$$





$$L = 8$$

- Consider the situation at the critical point.
- Since  $\xi$  is as large as the system itself, coarse graining does not produce less critical representations.
- In each figure, one sees structure over all length scales between the lower limit set by  $L$  and the upper limit set by the display size.
- A limiting trend is nevertheless apparent.



- Although the  $L = 4$  pattern differs qualitatively from  $L = L_{min}$ , the  $L = 4$  and  $L = 8$  patterns display qualitatively similar features.
- Thus patterns formed by the ordering variable at criticality look statistically the same when viewed on all sufficiently large length scales (fractal like).
- A statistical analysis of the spectrum of  $L = 4$  configurations shows that it is almost identical to that of the  $L = 8$  configurations.





- **Summary:** Under the coarse-graining operation there is an evolution or flow of the system's configuration spectrum.
- The flow tends to a limit, or **fixed point**, such that the pattern spectrum does not change under further coarse-graining.
- These scale-invariant limits have a trivial character for  $T > T_c$ , (a perfectly disordered arrangement) and  $T < T_c$ , (a perfectly ordered arrangement).
- The hallmark of the critical point is the existence of a scale-invariant limit which is neither fully ordered nor fully disordered but which possesses structure on **all** length scales.