Complex Disordered Systems

Static and dynamic correlations

Francesco Turci

Today

- **Characterisation of colloids**
  - Static (pair) correlations
  - Diffusive dynamics

Pairwise interactions

True colloidal systems can be complex:

- charges
- shape deformability/swelling
- instabilities of the components

Their theoretical description focuses on **essential ingredients**.

A simplifying assumption is that the potential energy of a collection of colloids is purely **pairwise**


$$U_N\left(\mathbf{r}^N\right) = \sum_{i=1}^{N-1} \sum_{j=i+1}^{N} V\left(r_{ij}\right) = \frac{1}{2} \sum_{i \neq j} V\left(r_{ij}\right)$$


where $V(r_{ij})$ is the pairwise, radial interaction potential that only depends on the distance between particle centres $r_{ij} = |\mathbf{r}_i - \mathbf{r}_j|$.

Aside (optional): Not pairwise?

Some examples of non-pairwise situations:

- **Atomistic models**: Electronic densities that fluctuate due to quantum effects are intrinsically many-body
- **True colloid-polymer mixtures**: if the polymers are not ideal, their spatial distribution and their forces depend on the positions of more than two colloids





C_{60} isosurface via density functional theory (Wikipedia) and configuration of non-ideal colloid-polymer mixture, Rohrbach et al JCP (2022)

Pairwise interactions and pair correlations


- For **pairwise** systems, all the thermodynamic properties are solely encoded in the **pairwise correlations**.

You have already seen such correlations in other contexts (Ising model, lattice gas).

The key pair correlation is the real space **radial distribution function**:

$$g(r) = \frac{1}{4\pi r^2 \rho N} \left\langle \sum_{i=1}^{N} \sum_{j \neq i} \delta\left(|\mathbf{r}_i - \mathbf{r}_j| - r\right) \right\rangle$$

Radial distribution function



The radial distribution function, algorithmic interpretation: at a distance r, we count the number of particle centres within a slice of width dr and then normalise.

Radial distribution function

- The radial distribution function $g(r)$ is related to the average local density at a distance r from a reference particle:
- To formally derive this, consider a system of N particles in volume V with average density $\rho = N/V$. The local density at a distance r from a reference particle is the number of particles in a shell of radius r divided by the shell volume $4\pi r^2 dr$. Averaging over the N reference particles,

$$\langle \rho(r) \rangle = \frac{1}{4\pi r^2 N} \left\langle \sum_{i=1}^{N} \sum_{j \neq i} \delta\left(|\mathbf{r}_i - \mathbf{r}_j| - r\right) \right\rangle$$

Normalising by the bulk density ρ gives the radial distribution function:

$$g(r) = \frac{1}{4\pi r^2 \rho N} \left\langle \sum_{i=1}^{N} \sum_{j \neq i} \delta\left(|\mathbf{r}_i - \mathbf{r}_j| - r\right) \right\rangle = \frac{\langle \rho(r) \rangle}{\rho}$$

where $\langle \rho(r) \rangle$ is the average density at distance r from a particle, and ρ is the bulk density.

Radial distribution function

- For an **ideal gas**, $g(r) = 1$ everywhere (no correlations).
- For **hard spheres**, $g(r) = 0$ for $r < \sigma$ (no overlap), and $g(r)$ shows oscillations at higher r due to packing effects.

> **Note**
>
> Hard spheres in 3D are not exactly solvable: we only have approximate theories (e.g. Percus-Yevick, below) and simulations. In 1D (hard rods) they are exactly solvable.

- The $g(r)$ is the central object of (equilibrium) liquid state theory
- Correlations between two particles 1 and 2 in a fluid can have two possible origins
  - a **direct** correlation between the two particles, mediated by direct interactions (e.g. collisions) between 1 and 2.
  - an **indirect correlation**, mediated by other particles in the fluid


Aside (optional): direct and indirect correlations

- The $g(r)$ contains **both**.
- We can assume the existence of a correlation function that instead only includes the **direct** part and call it $c(r)$. Then a hierarchy exists

$$h\left(r_{12}\right)=c\left(r_{12}\right)+\rho \int c\left(r_{13}\right) h\left(r_{32}\right) d\mathbf{r}_3$$

where $h(r)=g(r)-1$. This is the **Ornstein-Zernike integral equation** between particles 1,2 and 3.

Both $h(r)$ and $c(r)$ are unknown in principle. One needs an additional condition to close the equation: these are **physical closures** leading to approximations.



Meaning of the Ornstein-Zernike relation, adapted from Santos, Springer(2016)


Aside (optional): Percus-Yevick approximation

A common closure is the **Percus-Yevick approximation**.

$$c(r) = [1 + h(r)] \left[1 - e^{\beta V(r)}\right]$$

This closure interpolates between the hard-core exclusion (for $V(r) \to \infty$ inside the core, $g(r) = 0$) and the ideal gas limit (for $V(r) = 0$, $c(r) = 0$, so for hard spheres $c(r) = 0$ outside the core).

The result is (semi-)analytical. For hard spheres, it allows one to calculate the g(r) accurately in a wide range of packing fractions



PY approximation vs simulated data for hard spheres.

Equation of states from $g(r)$

The knowledge of the pair correlations also allows one to extract equation of states and significantly improve on the virial expansion, as the $g(r)$ contains packing effects!

$$P = \rho k_B T - \frac{2\pi\rho^2}{3}\int_0^\infty r^3 \frac{dV(r)}{dr} g(r) dr$$

Note

For hard spheres, $V(r)$ jumps at contact, so write $g(r) = y(r)\, e^{-\beta V(r)}$ with $y(r)$ continuous. Since $e^{-\beta V(r)}$ is a step at $r = \sigma$,

$$\beta \frac{dV}{dr}\, g(r) = -y(r) \frac{de^{-\beta V}}{dr} = -g(\sigma^+)\, \delta(r - \sigma)$$

Substituting into the pressure equation:

$$P = \rho k_B T + \frac{2\pi\rho^2}{3} k_B T \sigma^3 g(\sigma^+)$$

Therefore, for hard spheres, the equation of state only requires knowledge of the contact value $g(\sigma^+)$.

We will see this in a problem class.

Structure factor

The real-space correlations and relations have their reciprocal (*Fourier* space) counterparts.

The structure factor is defined as

$$S(k) = 1 + \rho \int [g(r) - 1] e^{-i\mathbf{k}\cdot\mathbf{r}} d\mathbf{r}$$

Isotropicity in 3D simplifies the expression to

$$S(k) = 1 + 4\pi\rho \int_0^\infty r^2 [g(r) - 1] \frac{\sin(kr)}{kr} dr$$


- Radial distribution functions $\rightarrow$ easily accessed in **confocal experiments** (large colloids)
- Structure factors $\rightarrow$ immediately accessed via **scattering experiments**

Structure factor

Interestingly, reciprocal space simplifies integral equations such as Ornstein-Zernike

$$\tilde{h}(k) = \frac{\tilde{c}(k)}{1 - \rho\tilde{c}(k)}$$

allowing one to read the direct correlation function directly from scattering data, as $S(k) = 1 + \rho\tilde{h}(k)$.




Structure factors (left) and radial distribution functions (right) for Gallium at extreme conditions, Drewitt et al PRL (2020)

Aside (optional): higher order correlations

Nothing prevents us from quantifying higher order correlations. In general these are non-trivial and controlled by the indirect part of the correlations.

Standard ways to identify **n-particle** correlations is to measure **structural motifs**.



Many body correlations in hard spheres, seminal work from Josh Robinson, PhD student in Bristol a few years ago, Robinson et al PRL (2019)

This is particularly important for dense systems (e.g. glasses, see future lecture).

Dynamics: single vs collective displacements

The thermal motion of individual colloids is often well modeled via the **Langevin equation**

$$m\frac{d\vec{v}}{dt} = -\gamma\vec{v} + \vec{\eta}(t)$$

Coarse-graining over small volumes (and short times) gives us access to the local density $\rho(\mathbf{r}, t)$ which obeys a **diffusion equation**

$$\frac{\partial\rho(\mathbf{r}, t)}{\partial t} = D\nabla^2\rho(\mathbf{r}, t)$$

The diffusivity is linked to the microscopic Langevin parameters via **fluctuation-dissipation** (Einstein's relation)

$$D = \frac{k_B T}{\gamma}$$

(where the friction is the dissipation and the random force is the fluctuations).

Diffusion equation: macroscopic picture

It is possible to (re)-derive the diffusion equation solely from macroscopic requirements. We have a macroscopic density $\rho(r,t)$ and a macroscopic flux $\vec{J}(r,t)$

- **continuity**: there is no mass loss in the dynamics

$$\frac{\partial \rho(\mathbf{r},t)}{\partial t} + \nabla \cdot \mathbf{J}(\mathbf{r},t) = 0$$

where the divergence $\nabla \cdot \mathbf{J}(\mathbf{r},t)$ represents the net outflow of particles from a given region due to the flux $\mathbf{J}$.

- **Fick's law** : we assume the flow is a gradient field of the density, flowing from high to low density

$$\mathbf{J}(\mathbf{r},t) = -D\nabla\rho(\mathbf{r},t)$$

It is immediate to combine the two to obtain the diffusion equation

$$\frac{\partial \rho(\mathbf{r},t)}{\partial t} = D\nabla^2\rho(\mathbf{r},t)$$

Measuring diffusion coefficients

The distribution $\rho(\mathbf{r}, t)$ encodes all the information about the motion of an ensemble.

Often, less is sufficient to characterise the diffusion. The second moment of the displacements (the mean squared displacement) provides this information. In d dimensions

$$\left\langle \left| \mathbf{r}(t) - \mathbf{r}(t_0) \right|^2 \right\rangle = 2dDt$$

Hence D can be extracted from linear fits to the MSD.

Stokes-Einstein relation: fluctuation dissipation for colloids

You have seen the **Einstein relation** for a Brownian particle with friction coefficient γ (Langevin dynamics):

$$D = \frac{k_B T}{\gamma}$$

(equivalently, in sedimentation equilibrium the drift flux nF/γ balances the Fickian flux $-D\,\partial n/\partial x$ of the Boltzmann profile).

For a sphere, **Stokes' law** $\gamma = 6\pi\eta R$ yields simply

$$D = \frac{k_B T}{6\pi\eta R}$$

Water, $R = 0.5\,\mu$m: $D \approx 0.5\,\mu\mathrm{m}^2/\mathrm{s}$, so a colloid needs $\tau_B = R^2/D \approx 0.5$ s to move its own radius (an atom: ~ 10 ps). Colloidal liquids and crystals can be **filmed** under a microscope.

This is the **Stokes-Einstein** relation, a balance between thermal fluctuations and viscous dissipation:

$$\text{Fluctuations: } k_B T \quad \longrightarrow \quad \text{Observed motion: } D \quad \longrightarrow \quad \text{Dissipation: } 6\pi\eta R.$$

Aside (optional): random walkers in 2d, example

Python Code ⟳ Start Over ▷ Run Code


```
1 n_steps, n_walkers, dim = 10000, 500, 2
2 steps = np.random.randint(-1, 2, size=(n_walkers, n_steps, dim))
3 positions = np.cumsum(steps, axis=1)  # cumulative sum along steps
4 msd = positions.var(axis=0).sum(axis=1)  # MSD at each timestep
5 plot_tj(positions,msd,walker=0,loglog=False)
```

Hidden assumptions

> **Note**
>
> - We have been considering ensembles of particles microscopically governed by the Langevin equation
> - We did not comment on the inter-particle interactions and assumed that the properties of the thermal bath determine the diffusivity D. In reality, interactions between particles can lead to deviations from simple diffusion, especially at higher concentrations, where collective effects and hydrodynamic interactions become important.


Dilute vs dense diffusion

- Consider hard spheres in a fluid.
- At low volume fractions ϕ they rarely collide and they perform overdamped (Brownian) dynamics. They will therefore diffuse as $\mathrm{MSD}(t) = 6D_0 t$
- Things change when we increase the packing.
  - Short times: $D_s < D_0$, free motion inside the **cage** formed by the neighbours, slowed only by **hydrodynamic** interactions (without them $D_s = D_0$).
  - Long times: $D_L \ll D_0$ cage breaking. Interactions are both hard core and hydrodynamic



Dilute vs dense diffusion

- At **very high** volume fractions the cage mechanism becomes more complex: particles are trapped for long times and only occasionally break cages:
  - cage-jump dynamics $\rightarrow$ glassy dynamics (see future lectures)
  - rare events $\rightarrow$ non-Gaussian (exponential tails) $\neq$ Gaussian/diffusive statistics
  - Breakdown of the **Stokes-Einstein** relation
    - the viscosity increases much faster than the decrease in diffusion $D\gamma/T \neq \text{constant}$
    - This means that $D\eta \neq \text{constant}$ and since the viscosity $\eta \sim \tau$ (a characteristic relaxation time) $D\tau \neq \text{constant}$



Stokes-Einstein breakdown in a theoretical model of glassy behaviour

Summary: colloids

Topic	Key result
Colloidal scale	$\mathrm{Pe}_g = \Delta m\, gR/k_BT \lesssim 1$: Brownian motion beats sedimentation
Van der Waals	$W_{vdW}(h) = -\frac{A_H}{6} f(h/R)$, attractive, $\approx -A_H R/12D$ near contact
Double layer	$W_{DR}(h) = B\frac{R}{\lambda_B} e^{-(h-2R)/\lambda_D}$ (gap $h-2R$), salt shortens λ_D
DLVO	$W_{vdW} + W_{DR}$: salt switches between stable and aggregating
Depletion (AO)	$W_{AO}(r) = -k_BT\rho_p^r V_{\text{overlap}}(r)$: entropic, $-(1+\frac{3R}{2\delta})\phi_p^r k_BT$ at contact
Hard spheres	single control parameter $\phi = \pi\sigma^3\rho/6$; entropy only
Equation of state	dilute fluid $Z_{\text{comp}} = 1+4\phi$; crystal (cell model) $Z_{\text{comp}} = \left[1-(\phi/\phi_{cp})^{1/3}\right]^{-1}$
Phase diagram	$\phi_f = 0.494$, $\phi_m = 0.545$, $\phi_{cp} = 0.74$: the crystal has **higher** entropy
Structure	$g(r)$ from imaging, $S(k)$ from scattering; hard spheres $Z_{\text{comp}} = 1+4\phi\, g(\sigma^+)$
Diffusion	$\langle \Delta r^2 \rangle = 2dDt$; dense: $D_0 > D_s > D_L$ (cages)
Stokes-Einstein	$D = \dfrac{k_BT}{6\pi\eta R}$: fluctuation $\leftrightarrow$ dissipation

Full summary with links in the notes