Complex Disordered Systems

Francesco Turci

Today

- Entropy matters
- Introduction to colloids

Entropy matters

Entropy matters



"I blame entropy."


Blame it on entropy (New Yorker)

More is different

Philip Anderson (Science 1972)

> **Complexity emerges when many simple units interact**: studying the individual components does not explain the emergence of collective properties

- Simple examples:
  - **phases of matter**
  - magnetism, superconductivity
  - emergence of order from disorder
- Also beyond physics:
  - formation of biological molecules
  - ecological networks
  - economic structures (markets, currencies etc.)




Complex mixtures, spintronic magnetisation, ecological networks and Hong Kong stock market

Beyond the usual phases of matter


We sometimes oversimplify...

extreme dilution



gases

extreme order



perfect crystals

Extreme examples of phases of matter

Beyond the usual phases of matter

We sometimes oversimplify...



Complex Fluids or *Soft Matter*

© Edinburg Complex Fluids

Complex fluids and soft condensed matter

- **Soft condensed matter** includes assemblies of colloids, polymers, surfactants, and biological macromolecules and much more.
  - Term due to Pierre-Gilles de Gennes (Nobel 1991)
- Often, many of these are also referred to as **complex fluids**.
- These materials are **easily deformed** and show complex, disordered structures.
- Their properties are determined by a balance between **energy and entropy**.
- **Thermal fluctuations** play a major role in their behavior.
- Understanding them requires **statistical mechanics** .

Thermal fluctuations and entropy

- Helmholtz Free energy per particle

$$f = u - Ts$$

- Entropy counts the arrangements: $s = \dfrac{k_B}{N} \ln \Omega = k_B \ln \omega$, with $\omega = \Omega^{1/N}$ arrangements per particle (geometric mean). Because of the logarithm, $\omega \sim 10$–1000 gives $s \approx 2$–$7\, k_B$, so Ts is a few $k_B T$.
- In soft matter the **interaction energies** per particle are also a few $k_B T$:

$$\boxed{\Delta u \sim k_B T}$$


  so thermal fluctuations can break and rearrange the structure.
- The two terms in f are comparable: **entropy can decide the structure**.

Thermal fluctuations: single particle in a double well

Temp. 1

Tilt 0

Reset simulation



Systems and definitions

Elementary constituents and energy scales

- Soft matter systems are made of **many parts**
- The assembly of these many parts can be **easily deformed.**
- Interactions between these parts are **weak** compared to thermal or mechanical forces.


- **Hard condensed matter:**
- Basic units: atoms
- Strong interactions (0.1–10 eV)
- Covalent/ionic bonds
- Focus on low temperatures

- **Soft matter:**
- Basic units: molecular aggregates
- Weak interactions (0.001–0.2 eV)
- Van der Waals, hydrogen bonds
- $1k_BT \approx 0.025eV$ (room temperature)

Coarse graining

- Soft matter interactions are mainly **electrostatic**.
- Atomistic details are often unimportant for macroscopic properties.
- **Coarse-graining** simplifies models by focusing on key features:
  - deliberate selection of what matters
  - systematic integration of a number of degrees of freedom

- Example: The oxDNA model represents DNA as a chain of coarse grained units, much larger than the atoms.




Atomistic DNA structure

Coarse graining

- Soft matter interactions are mainly **electrostatic**.
- Atomistic details are often unimportant for macroscopic properties.
- **Coarse-graining** simplifies models by focusing on key features:
  - deliberate selection of what matters
  - systematic integration of a number of degrees of freedom

- Example: The oxDNA model represents DNA as a chain of coarse grained units, much larger than the atoms.







oxDNA model: (a) Base structure on one strand; (b) planarity of the bonding; (c) an example of the resulting double strand.

Classes of systems

In our exploration of soft matter we will focus on five main classes of systems which display different physics:

- colloidal dispersions
- polymeric systems
- liquid crystals
- arrested systems
- active matter






Colloidal assemblies, polymers, surfactants, liquid crystals, glasses and bacteria.

Colloidal dispersions


- **Colloidal dispersions**: small particles (nano–micrometer) suspended in a solvent.
- **Spherical colloids** are common, but many shapes and interactions exist.
- Behave as **“big atoms”**: show Brownian motion, phase transitions, and can form ordered structures.
- **Larger size** and **slower dynamics** make them ideal for direct observation of phenomena like:
  - crystallisation
  - glass formation
  - gel formation



(A) Solvent collisions cause random forces on particles. (B) Particles execute random walks on larger timescales. (C) Hard spheres crystallize at high density (right) due to reduced available space compared to fluid phase (left). From Manohran, *Science* (2015)

Polymeric systems


- **Polymers** are long-chain macromolecules made of repeating monomers.
- Their properties result from a balance of **entropy** and **energy**.
- Two main types: **synthetic polymers** (e.g., plastics) and **biopolymers** (e.g., DNA, proteins).
- **Entanglement**: chains cannot cross, leading to unique mechanical behavior.



Polymer entanglement, from Likhtman and Ponmurugan, *Macromolecules* (2014)

Liquid crystals


- **Liquid crystals** form when anisotropic soft matter units (e.g., rod-like or disk-shaped molecules) pack densely, leading to partial order—intermediate between liquids and crystals.
- **Continuum free energy theories** describe liquid crystals by considering the symmetry of their order parameters.
- Liquid crystals are crucial in technologies like **liquid crystal displays (LCDs)**.



Texture of nematic liquid crystals, from https://doi.org/10.3986/alternator.2020.38

Arrested systems


- **Arrested systems** are trapped in disordered, non-equilibrium states.
- Lack of long-range order leads to **slow structural relaxation** and prevents reaching the global energy minimum.
- Examples: **glasses** and **gels**
- These systems exhibit **rigidity, elasticity, and plasticity**: disordered (semi)-solids.
- Relaxation times are **longer than observable timescales**, making equilibrium statistical mechanics insufficient.



A gel (left) and a glass (right) from molecular dynamics with local structure highlighted in colours

Active matter

- **Active matter**: systems of units that **consume energy** to move or exert forces.
- **Inherently out of equilibrium** due to continuous energy input.
- Includes **biological systems** (bacteria, cells, flocks) and **synthetic systems** (self-propelled colloids).
- Shows **emergent collective behaviors**: swarming, clustering, pattern formation.
- Studied using **hydrodynamic theories** and **agent-based models**.



Example of active systems, from Cichos et al. *Nature Machine Intelligence* (2020)


Introduction to Colloids


Kinds of colloids

- Colloids = **dispersed phase** + **dispersion medium**







Opal, paint, smoke, ice-cream, milk, fog, expanded polystyrene and foam.

Kinds of colloids

Dispersed phase	Dispersion medium	
	Solid	**Liquid**
Solid	*Solid suspension:* pigmented plastics, stained glass, ruby glass, opal, pearl	*Sol, colloidal suspension:* metal sol, toothpaste, paint, ink, clay slurries, mud
Liquid	*Solid emulsion:* bituminous road paving, ice cream	*Emulsion:* milk, mayonnaise, butter, pharmaceutical creams
Gas	*Solid foam:* zeolites, expanded polystyrene, ‘silica gel’	*Foam:* froths, soap foam, fire-extinguisher foam

IUPAC definition

The *International Union of Pure and Applied Chemistry* (IUPAC) defines

> **colloidal**: The term refers to a state of subdivision, implying that the molecules or polymolecular particles dispersed in a medium have at least in one direction a dimension roughly between 1 nm and 1 μm , or that in a system discontinuities are found at distances of that order.

Note

The definition has little to do with the **chemistry** of the polymolecules (i.e. aggregates), but essentially is determined by the **size**.

- *Why the size?*

Colloidal scale

- Particle of size R (micrometric) in medium with constant collisions at **thermal energy** $\sim 1k_BT$
- Compare the energy with the potential energy of settling over a length R
- Obtain a **nondimensional number** the **gravitational Péclet number**

$$\mathrm{Pe}_g = \frac{\Delta m g R}{k_B T}$$

Example

For a **milk protein particle** (casein micelle) in water:


- $R \sim 0.5\,\mu$m (estimate)
- $\Delta m = \frac{4}{3}\pi R^3 \Delta\rho \sim \frac{4}{3}\pi(0.5\times10^{-6})^3 \times 200 \sim 1\times10^{-16}$ kg (assuming $\Delta\rho \sim 200$ kg/m³)
- $g = 9.8$ m/s²
- $k_BT \sim 4\times10^{-21}$ J (room temperature)

$$\mathrm{Pe}_g = \frac{1\times10^{-16}\times 9.8\times 0.5\times10^{-6}}{4\times10^{-21}} \sim 0.1$$

Since $\mathrm{Pe}_g < 1$, **thermal motion dominates gravity** → particles remain suspended → stable colloid!


Observation time and stability

- Any measurement in physics has an **observation time** t_{obs}
- Thermal systems have some degree of memory and hence an intrinsic timescale τ (**relaxation time**)
- When $\tau \ll t_{\mathrm{obs}}$ we can take time-averages and consider the system in a stable **steady state**.
- In the absence of net currents, this is an **equilibrium state**
- A colloidal dispersion is stable if it is able to remain dispersed and Brownian for a time much longer than t_{obs}.
- Instability leads to **aggregation**



Destabilisation

- Milk is an emulsion where droplets of fat are dispersed in water and stabilised by proteins.
- If lemon juice is added, the dispersion medium (water) changes, the pH drops, and the emulsion is *destabilised*.
- The interactions change → separation of curds (solid) and whey (liquid).



Curds and whey resulting from the destabilisation of milk, a colloidal emulsion. (Wikimedia)

Which interactions?

- At the colloidal scale: **gravity** and **electrostatics** (eventually **magnetostatics**).
- Electrostatic forces are the main **microscopic** forces: colloids are typically charged
- **Separation of length and timescales**: colloids are much larger and slower than the atomic or molecular constituents of the surrounding fluid.
- Colloid–colloid interactions result from the collective effect (sum or average) of many microscopic interactions, evaluated over times much longer than microscopic relaxation times.
- When we focus on colloidal scales, we **integrate out** the microscopic degrees of freedom (*coarse-graining*)
- Coarse-grained interactions are then obtained and are called **effective interactions**.

- **Fundamental forces:**
- gravity
- electro (magneto) static forces

- **Effective forces**
- Van der Waals forces
- electrostatic repulsion (DLVO)
- depletion forces
- hydrophobic/hydrophilic interactions
- steric forces