

A Phase II, Open-label Study of Rovanimab plus Carboplatin in Previously Treated Non-Small Cell Lung Cancer

Protocol ONC-210, Version 2.0

FICTIONAL EXAMPLE PROTOCOL - FOR SOFTWARE DEMONSTRATION ONLY. All names and details are invented.

Primary endpoint: objective response rate by RECIST 1.1.

5. Study Treatment

Rovanimab vials must be stored at 2 to 8 C and must not be frozen.

Treatment must be held if the absolute neutrophil count is below $1.0 \times 10^9/L$ on the day of dosing.

Dosing a subject whose absolute neutrophil count is below $1.0 \times 10^9/L$ is an important protocol deviation.

7. Assessments

Tumour assessments (CT scans) are performed every 6 weeks plus or minus 7 days.

A tumour assessment outside its window, or missed, is an important protocol deviation.

Pharmacokinetic samples are collected before and at the end of the Cycle 1 Day 1 infusion.

9. Adverse Events

Disease progression is an efficacy outcome and must not be reported as an adverse event or serious adverse event.

Hospitalization due only to disease progression is not a serious adverse event.

All serious adverse events must be reported to the sponsor within 24 hours of the site becoming aware of the event.

Infusion-related reactions of grade 3 or higher are adverse events of special interest and must be reported within 24 hours.

11. Informed Consent

When a new version of the informed consent form is approved, ongoing subjects must be re-consented at their next visit.