

A Phase III, Observer-blind Study of an Adjuvanted RSV Vaccine (RSVa-2) in Adults Aged 60 and Over

Protocol VAX-118, Version 1.2

FICTIONAL EXAMPLE PROTOCOL - FOR SOFTWARE DEMONSTRATION ONLY. All names and details are invented.

Primary endpoint: RSV neutralizing antibody titres at Day 29.

6. Vaccine Handling

Vaccine must be stored at 2 to 8 C at all times.

Any temperature excursion must be reported to the sponsor within 1 calendar day, and affected doses must be quarantined and must not be administered until the sponsor approves their use.

Administering a dose exposed to an unapproved temperature excursion is an important protocol deviation.

7. Schedule

The Day 29 visit must occur within plus or minus 3 days.

Blood samples for immunogenicity are taken before vaccination on Day 1 and at Day 29.

A missed Day 29 immunogenicity sample is an important protocol deviation.

8. Reactogenicity

Subjects record solicited reactions in an electronic diary for 7 days after vaccination.

Diary completion on fewer than 5 of the 7 days requires retraining of the subject and is a minor deviation.

9. Safety Reporting

All serious adverse events must be reported to the sponsor within 24 hours of the site becoming aware of the event.

Guillain-Barre syndrome is an adverse event of special interest and must be reported within 24 hours.