

A Phase III, Randomized, Double-blind, Placebo-controlled Study of Zelvatinib in Adults with Moderate-to-Severe Plaque Psoriasis

Protocol ZLV-301, Amendment 2 (Version 3.0)

FICTIONAL EXAMPLE PROTOCOL - FOR SOFTWARE DEMONSTRATION ONLY. All names and details are invented.

Primary endpoint: proportion of subjects achieving PASI 75 at Week 12.

The Week 12 PASI assessment is the primary efficacy assessment.

6. Study Population

Subjects who do not meet all eligibility criteria must not be randomized.

Randomization of an ineligible subject is an important protocol deviation.

7. Study Drug

Zelvatinib tablets must be stored at 15 to 25 C.

Any temperature excursion must be reported to the sponsor within 1 business day, and affected kits must be quarantined and not dispensed until the sponsor approves their use.

Study drug must be interrupted if ALT or AST rises above 3 times the upper limit of normal.

Returned tablets must be counted at every dispensing visit.

8. Schedule of Assessments

Visits from Week 2 to Week 12 must occur within plus or minus 3 days of the scheduled day; visits from Week 16 onward within plus or minus 7 days.

A missed Week 12 PASI assessment is an important protocol deviation.

10. Safety Reporting

Worsening of psoriasis is recorded as an adverse event only if it is more severe than the subject's usual fluctuation.

A hospitalization for an elective procedure that was planned before the subject signed the informed consent form is not a serious adverse event.

All serious adverse events must be reported to the sponsor within 2 business days of the site becoming aware of the event.

Any pregnancy in a subject must be reported to the sponsor within 24 hours of the site becoming aware of it.