

LIST OF FIGURES

	<i>Page</i>
Figure 2.1	Pseudo-adiabatic diagram for dewpoint reduction to 1 000 hPa at zero height.....11
Figure 2.2	Determination of maximum dewpoint in a storm. Representative dewpoint for this weather map is the average of values in boxes.12
Figure 2.3	Enveloping maximum persisting 12-hour dewpoints at a station.....13
Figure 2.4	Maximum persisting 12-hour 1 000-hPa dewpoints for August (Environmental Data Service, 1968).....13
Figure 2.5	Schematic illustrating concept for alternative moisture adjustment for intervening barrier to air flow (after Hart, 1982)16
Figure 2.6	Transposition limits (heavy dashed line) of 9–13 July 1951 storm. Locations of synoptically similar summer storms marked by X. Light lines indicate maximum persisting 12-hour 1 000-hPa dewpoints (°C) for July.19
Figure 2.7	Example of storm transposition. Long dashed lines indicate maximum persisting 12-hour 1 000-hPa dewpoints (°C) for the same time of year the storm occurred or within 15 days according to common practice (section 2.3.1).21
Figure 2.8	Isohyets (cm) for the 6-hour afternoon storms of 16 June 1965 (solid lines) and 17 June 1965 (dashed lines) in eastern Colorado (Riedel and others, 1969)24
Figure 2.9	Isohyets (cm) resulting from the combination of patterns of the 6-hour storms of 16–17 June 1965 shown in Figure 2.8 (Riedel and others, 1969).....25
Figure 2.10	Depth–duration envelope of transposed maximized storm values for 2 000 km ²26
Figure 2.11	Depth–area envelope of transposed maximized 24-hour storm rainfall values26
Figure 2.12	Enveloping DAD curves of probable maximum precipitation for a hypothetical basin26
Figure 2.13	Seasonal variation of probable maximum precipitation in the Upper Tigris River basin in Iraq.....29
Figure 2.14	Maximization of a pattern storm by the sliding technique for a 5 000 km ² basin, the point of first contact occurs on the 72-hour curves between about 2 000 and 4 500 km ²31
Figure 2.15	Example of enveloping DAD curves of PMP and within-storm rainfall depths31
Figure 2.16	Critical isohyetal pattern over 3 000 km ² basin.....32
Figure 2.17	Isohyetal profile constructed from data in Table 2.332
Figure 2.18	Observed rainfalls expressed as a percentage of PMP in the United States, east of the 105th meridian \geq 50 per cent of all-season PMP for the 24-hour duration and an area size of 518 km ² (Riedel and Schreiner, 1980).....35
Figure 3.1	Simplified inflow and outflow wind profiles over a mountain barrier (United States Weather Bureau, 1961a)40
Figure 3.10	Decrease of maximum precipitable water with duration (after United States Weather Bureau, 1958)51
Figure 3.11	Seasonal envelope of maximum observed dewpoints at Los Angeles, California (after United States Weather Bureau, 1961a).....52
Figure 3.12	Maximum persisting 12-hour 1 000-hPa dewpoints (°C) for February (United States Weather Bureau, 1961a)52
Figure 3.13	Maximum P/M ratios for combination with orographic storms (after United States Weather Bureau, 1961a)53
Figure 3.14	Ratios of 6-hour and 72-hour precipitation to 24-hour precipitation (United States Weather Bureau, 1961a)53
Figure 3.15	Effective elevation and barrier heights (30.5 m) in northern California (square delineates Blue Canyon orographic model test area; United States Weather Bureau, 1961a).....54
Figure 3.16	Six-hour 518 km ² convergence PMP (inches) for January and February (square delineates Blue Canyon orographic model test area; United States Weather Bureau, 1961a).....55
Figure 3.17	Variation of convergence PMP index with basin size and duration for December (United States Weather Bureau, 1961a)56