

	<i>Page</i>
Figure 5.34 Orographic factor (T/C) map for the state of Colorado east of the Continental Divide (Miller and others, 1984b)	105
Figure 5.35 Storm intensity factor (M) for the state of Montana east of the Continental Divide (Miller and others, 1984b)	106
Figure 5.36 Twenty-four-hour 25.9-km ² PMP estimates (inches) for the northern portion of the state of New Mexico east of the Continental Divide	106
Figure 5.37 Schematic diagram of the sub-regional system used in developing depth–area–duration relations (Miller and others, 1984b)	107
Figure 5.38 Depth–area–duration relation for orographic region of Missouri and Yellowstone River basins (Miller and others, 1984b)	108
Figure 5.39 Twenty-four-hour 25.9 km ² orographic PMP (inches) for southern Arizona, south-western New Mexico and south-eastern California (Hansen and others, 1977)	109
Figure 5.40 Variation of orographic PMP with basin size for Colorado River and Great Basin drainages (Hansen and others, 1977)	109
Figure 5.41 Adopted durational variation in orographic PMP for Colorado River and Great Basin drainages (Hansen and others, 1977)	110
Figure 5.42 Isolines on convergence 100-year 24-hours rainfall (cm) for central United States (Fenn, 1985)	110
Figure 5.43 Depth–area relations for local storm PMP Pacific Northwest states	113
Figure 5.44 Idealized isohyetal pattern for local storm PMP areas up to 1 300 km ² (Hansen and others, 1977)	114
Figure 5.45 One-hour 2.6-km ² local storm PMP in inches for elevations to 1 830 m	114
Figure 5.46 Temporal distribution relation for Mud Mountain Dam	116
Figure 5.47 Twenty-four-hour PMP isoline map for Auburn watershed (inches)	118
Figure 5.48 California DAD region map	119
Figure 5.49 Curve of relationship between areal mean PMP and duration for Auburn watershed	120
Figure 5.50 Temporal distribution of 72-hour PMP for Auburn watershed	121
Figure 5.51 California Wash watershed	121
Figure 5.52 Isoline map of PMP for California for local storms with 1-hour duration and storm area of 2.6 km ²	122
Figure 5.53 Corrected percentage of moisture content in air columns for different watershed elevations	122
Figure 5.54 Coefficients of correcting PMP for different durations	123
Figure 5.55 Isoline map of watershed correction types	124
Figure 5.56 Coefficients of area reduction corresponding to Type C	124
Figure 5.57 Curve of duration–area PMP relationship for the watershed	124
Figure 5.58 Generalized isohyetal map of PMP	125
Figure 5.59 Montana storm isohyets (inches) transposed to Cheesman basin	126
Figure 5.60 Analysis of isohyetal orientations for selected major storms in United States east of the 105th meridian adopted as recommended orientation for PMP within $\pm 40^\circ$ (Hansen and others, 1982)	85
Figure 5.61 Cheesman 100-year 24-hour rainfall (inches)	126
Figure 5.62 Grid of rainfall depth (inches) of Montana storm transposed over the Cheesman basin multiplied by the ratio of the grids of Figures 5.61 to 5.60	127
Figure 5.63 Zones for use with the generalized short-duration method (Australian Bureau of Meteorology, 2003)	128
Figure 5.64 Generalized short-duration method depth–duration–area curves (Australian Bureau of Meteorology, 2003)	128
Figure 5.65 Reduction factor for geographic variation from extreme moisture (Australian Bureau of Meteorology, 2003)	130
Figure 5.66 Temporal distribution for use with PMP estimates derived using the generalized short-duration method (Australian Bureau of Meteorology, 2003)	130
Figure 5.67 Generalized short-duration method spatial distribution	130
Figure 5.68 Monthly percentage moisture adjustment for southern Australia (south of 30° S); areal limit is 500 km ²	132
Figure 5.69 Boundaries between PMP methods	133
Figure 5.70 An example of a set of standard area polygons as applied to the 6-day storm, 18–23 January 1974	134