

Figure 3.18	Typical orographic rainfall pattern for southwesterly winds. Isolines indicate ratios of orographic to non-orographic rainfall (Schwarz, 1965)	58
Figure 3.19	Basin subdivisions for check of topographic effects on basin-wide precipitation volume (Schwarz, 1963).....	58
Figure 3.2	Single-layer laminar flow wind model	41
Figure 3.20	March 24-hour 25 900 km ² PMP (cm; after Schwarz, 1965).....	59
Figure 3.21	Seasonal variation of PMP for 55 426 km ² as percentage of March PMP (Schwarz, 1965).....	59
Figure 3.22	Depth–area ratios (55 426/ 20 688 km ²) for 24-hour rainfall (Schwarz, 1965).....	60
Figure 3.23	Depth–duration relations in percentage of 24-hour rainfall (Schwarz, 1965)	61
Figure 3.24	Six-hour PMP storm pattern (mm) for maximum 6-hour increment for total basin (55 426 km ²) (Schwarz, 1965); isohyet values are applicable to the March PMP.....	61
Figure 3.25	Nomogram for obtaining isohyet values for maximum 6-hour rainfall increment in pattern storms	62
Figure 3.26	PMP storm pattern for maximum 6-hour increment for sub-basin (21 000 km ² ; Schwarz, 1965); isohyet values are applicable to March	63
Figure 3.3	Multiple-layer laminar flow wind model	41
Figure 3.4	Blue Canyon orographic laminar-flow model test area in California (United States Weather Bureau, 1966)	42
Figure 3.5	Precipitation station elevations relative to adopted ground profile for test area of Figure 3.4 (United States Weather Bureau, 1966)	42
Figure 3.6	Air streamlines and precipitation trajectories for test case.....	43
Figure 3.7	Maximum 1-hour wind profile and supporting data (United States Weather Bureau, 1961 <i>b</i>).....	48
Figure 3.8	Variation of maximum 6-hour wind speed with time (after United States Weather Bureau, 1961 <i>a</i>).....	49
Figure 3.9	Six-hour orographic PMP (inches) for January (square delineates Blue Canyon orographic model test area; United States Weather Bureau, 1961 <i>a</i>)	50
Figure 4.1	Km as a function of rainfall duration and mean of annual series (Hershfield, 1965).....	65
Figure 4.2	Adjustment of mean of annual series for maximum observed rainfall (Hershfield, 1961 <i>b</i>)	67
Figure 4.3	Adjustment of standard deviation of annual series for maximum observed rainfall (Hershfield, 1961 <i>b</i>).....	67
Figure 4.4	Adjustment of mean and standard deviation of annual series for length of record (Hershfield, 1961 <i>b</i>)	67
Figure 4.5	Adjustment of fixed-interval precipitation amounts for number of observational units within the interval (Weiss, 1964)	68
Figure 4.6	Examples of (A) isohyetal pattern centred over basin as would be the case for storm-centred depth–area curves, and (B) two possible occurrences of isohyetal patterns over a geographically fixed area as would be the case in development of curves for a geographically fixed area (Miller and others, 1973).....	69
Figure 4.7	Depth–area, or area-reduction, curves for western United States (United States Weather Bureau, 1960)	69
Figure 4.8	Maximum depth–duration curve (Huff, 1967).....	70
Figure 5.1	Contrast in (a) PMP (Schwarz, 1963) and (b) 25-year rainfall patterns (United States Weather Bureau, 1962), both for 24-hour at a point, Island of Hawaii; particularly note differences on north-western coast	80
Figure 5.2	Depth-duration interpolation diagrams from study for western United States (United States Weather Bureau, 1960)	81
Figure 5.3	All-season envelope of 24-hour 25.9 km ² PMP (cm; Schreiner and Riedel, 1978); stippled areas are regions where orographic effects have not been considered.....	82
Figure 5.4	Examples of temporal sequences of 6-hour precipitation in major storms in eastern and central United States (Hansen and others, 1982).....	83
Figure 5.5	Standard isohyetal pattern recommended for spatial distribution of PMP for the United States east of the 105th meridian (Hansen and others, 1982)	84
Figure 5.60	Montana 100-year 24-hour rainfall (inches)	126