



UNIVERSITÀ DEGLI STUDI DI PADOVA

Dipartimento di Fisica e Astronomia

Corso di Laurea Triennale in Fisica

Tesi di Laurea

**Inverse Beta Decay events selection in JUNO using
Machine Learning algorithms**

Relatore

prof. Alberto Garfagnini

Correlatori

dott. XXXXXX

dott. XXXXXX

Laureando

Candidate

Badge Number

Anno Accademico YYYY/YYYY

Discussion date

Abstract

The Jiangmen Underground Neutrino Observatory (JUNO) will be the largest liquid scintillator-based neutrino detectors in the World, for the next decade. Thanks to its large active mass (20 kt) and state of the art performances (3% effective energy resolution at 1 MeV), it will be able to perform important measurements in neutrino physics. The proposed thesis will study the application of different Machine Learning inspired algorithms for the discrimination of signal events (interactions of anti-neutrinos coming from the nearby nuclear power plants) from background events.

Contents

Contents	ii
1 Introduction	1
1.1 Neutrinos Oscillation	1
1.2 The JUNO experiment	2
2 Framework	5
2.1 Machine Learning	5
References	7

Chapter 1

Introduction

1.1 Neutrinos Oscillation

The Standard Model of elementary particle interactions provides an accurate description of strong, weak, and electromagnetic interactions, but it treats these interactions as distinct and unrelated. Within this framework, neutrinos are assumed to be massless, but this assumption has been called into question by physicists. Neutrino oscillations, which occur when neutrinos change from one flavor to another, are a potential indication of neutrino mass.

The term "neutrino oscillations" refers to this phenomena and it involve the conversion of a neutrino of a particular flavor to another as it propagates through space.

Each known flavor eigenstate, $(\nu_e, \nu_\mu, \nu_\tau)$, linked to three respective charged leptons (e, μ, τ) via the charged current interactions can be considered a complex combination of neutrino mass eigenstates as follow:

$$\begin{pmatrix} \nu_e \\ \nu_\mu \\ \nu_\tau \end{pmatrix} = U_{\text{PMNS}} \begin{pmatrix} \nu_1 \\ \nu_2 \\ \nu_3 \end{pmatrix}$$

in wich ν_i are the three mass eigensates, that have 3 masses m_i ($i = 1, 2, 3$), which are non-degenerate, with $m_i \neq m_j$ for $i \neq j$.

The matrix U_{PMNS} , called Pontecorvo-Maki-Nakagawa-Sakata (PMNS) matrix, is composed of three rotation matrices, R_{23} , R_{13} , and R_{12} , each corresponding to a different mixing angle, θ_{23} , θ_{13} , and θ_{12} , respectively and a parameter δ_{CP} called the Dirac CP-violating phase. For this case, the Majorana CP phases are η_i ($i = 1, 2$), which are only physically possible if neutrinos are Majorana-type particles and do not participate in neutrino oscillations. Therefore, U can be expressed as:

$$U_{\text{PMNS}} = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13}e^{-i\delta_{CP}} \\ 0 & 1 & 0 \\ -s_{13}e^{i\delta_{CP}} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} e^{i\eta_1} & 0 & 0 \\ 0 & e^{i\eta_2} & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

The theoretical framework for neutrino oscillations involves the calculation of the oscillation probability as a function of the distance traveled by the neutrino, the neutrino mixing

matrix, and the difference in squared masses between the three neutrino mass states, Δm_{ij}^2 . Specifically, two nuclear power reactors 53 km away from the detector, which mostly produce anti-electron neutrinos $\bar{\nu}_e$ with energy below 10 MeV, are the principal sources of neutrinos for the JUNO experiment. So, it is necessary for the JUNO experiment to calculate the survival probability $P(\bar{\nu}_e \rightarrow \bar{\nu}_e)$ of electron antineutrinos.

$$P(\bar{\nu}_e \rightarrow \bar{\nu}_e) = 1 - \sin^2 2\theta_{12} c_{13}^4 \sin^2 \left(\frac{\Delta m_{21}^2 L}{4\mathcal{E}} \right) - \sin^2 2\theta_{13} \left[c_{12}^2 \sin^2 \left(\frac{\Delta m_{31}^2 L}{4\mathcal{E}} \right) + s_{12}^2 \sin^2 \left(\frac{\Delta m_{32}^2 L}{4\mathcal{E}} \right) \right]$$

where $s_{ij} \equiv \sin \theta_{ij}$, $c_{ij} \equiv \cos \theta_{ij}$, \mathcal{E} is the neutrino energy, L the travelled distance and $\Delta m_{ij}^2 \equiv m_i^2 - m_j^2$.

Past experiments have already given estimates for Δm_{21}^2 , $|\Delta m_{31}^2|$ and the 3 mixing angles.




Figure 1.1: JUNO's reactor antineutrino energy spectrum is shown with and without the effect of neutrino oscillation. The gray dashed curve includes only the term in the disappearance probability modulated by $\sin^2(2\theta_{12})$, while the blue and red curves use the full oscillation probability for normal and inverted mass orderings. Spectral features driven by oscillation parameters are illustrated, highlighting the rich information available in JUNO's high-resolution measurement of the oscillated spectrum.

The aim of JUNO is then to improve these results, and especially fix the sign of Δm_{31}^2 by discriminating between two possibilities: *Normal Ordering* (**NO**), where $|\Delta m_{31}^2| = |\Delta m_{32}^2| + |\Delta m_{21}^2|$, and if $m_1 < m_2 < m_3$ we have the so called *Inverted Ordering* (**IO**), where $|\Delta m_{31}^2| = |\Delta m_{32}^2| - |\Delta m_{21}^2|$, and $m_3 < m_1 < m_2$. In fact, depending on the sign of Δm_{31}^2 , the plot of 1.1 is minimally different.

1.2 The JUNO experiment

The Jiangmen Underground Neutrino Observatory (JUNO) is an experiment that is currently under construction in Southern China. It is a multipurpose experiment that will use a 20 kton liquid scintillator target to study neutrinos from a variety of natural sources as well as from nuclear reactors. The experiment is being built with the aim of achieving a high sensitivity to the neutrino mass ordering, which is one of the most important unresolved issues in neutrino physics.




Figure 1.2

JUNO is expected to detect a large number of antineutrinos from nuclear reactors, with most of them originating from the Taishan and Yangjiang nuclear power plants (NPPs). These two plants are located at a baseline of about 52.5 km from the JUNO detector, which was optimized for the best sensitivity to the neutrino mass ordering, and have a combined nominal thermal power of $26.6 \text{ GW}_t h$. However, in order to precisely measure the unoscillated reactor antineutrino spectrum shape, a dedicated small satellite detector called TAO will be placed at about 30 m from one of the Taishan reactors. TAO will serve as a data-driven input to constrain the spectra of the other cores.

A schematic illustrating the location of both JUNO and TAO is shown in Fig. 1.2.

In JUNO's location, the energy spectrum will be distorted by two types of oscillations. The first is a slow (low frequency) oscillation driven by Δm_{21}^2 and modulated by $\sin^2 \theta_{12}$, while the second is a fast (high frequency) oscillation driven by Δm_{31}^2 and modulated by $\sin^2 \theta_{13}$, as shown in Fig. 2. JUNO will be the first experiment to observe these two oscillation modes simultaneously. Fitting the data spectrum against the predicted spectrum distorted by standard neutrino oscillations enables measuring the oscillation parameters.

It is worth noting that the measurement of the neutrino mass ordering is not addressed in the current publication. Nonetheless, the measurement of the energy spectrum in JUNO will provide valuable information on the oscillation parameters and will shed light on some of the most important questions in neutrino physics.

In addition to studying reactor antineutrinos, JUNO will also investigate neutrinos from natural sources such as the Sun, supernovae, and atmospheric neutrinos. The experiment is expected to make important contributions to a variety of topics in neutrino physics, including the measurement of the absolute neutrino mass scale, the search for sterile neutrinos, and the determination of the neutrino mass ordering.

To achieve its scientific goals, JUNO employs a large liquid scintillator detector with a fiducial mass of 20 kton. The detector is located in a cylindrical cavity excavated under a hill in Jiangmen City, Guangdong Province, China, at a depth of 700 m.w.e. to reduce background radiation from cosmic rays. The scintillator is doped with a small amount of gadolinium to enhance its sensitivity to antineutrinos via the inverse beta decay (IBD) process.

The reactor antineutrinos in JUNO will primarily come from two nuclear power plants, Taishan and Yangjiang, located at a distance of approximately 52.5 km from the detector. To accurately measure the unoscillated reactor antineutrino spectrum, JUNO will use a small

satellite detector called TAO, which will be positioned about 30 m from one of the Taishan reactors. The data from TAO will be used to constrain the spectra of the other cores.

JUNO's location and the characteristics of the liquid scintillator will allow the experiment to observe both slow and fast oscillation modes of reactor antineutrinos, making it the first experiment to do so. By fitting the observed spectrum against the predicted spectrum distorted by standard neutrino oscillations, JUNO will be able to measure the oscillation parameters Δm_{31}^2 , Δm_{21}^2 , $\sin^2 \theta_{12}$, and $\sin^2 \theta_{13}$. The oscillated spectrum in JUNO will also depend on the neutrino mass ordering, providing sensitivity to this parameter.

The liquid scintillator used in JUNO is a combination of LAB (linear alkyl benzene) and PPO (2,5-diphenyloxazole) doped with a small amount of bis-MSB (1,4-bis(2-methylstyryl) benzene). When a neutrino interacts with the scintillator, it can produce charged particles such as electrons, protons, and alpha particles that travel through the scintillator and excite the scintillation molecules. This excitation results in the emission of photons with a wavelength of around 430 nm. These photons are detected by 20,000 20-inch photomultiplier tubes (PMTs) distributed in a 3-dimensional arrangement inside the detector.

The PMTs detect the light and convert it into an electrical signal. The signals from all the PMTs are then combined to reconstruct the position and energy of the original neutrino interaction. This technique allows JUNO to measure the energy of the incoming neutrino to high precision, which is crucial for studying neutrino oscillation.

Moreover, the scintillator's composition and the detector's design are optimized to reduce background noise from other sources of radiation, such as cosmic rays and natural radioactivity. By carefully controlling these backgrounds, JUNO aims to achieve a signal-to-background ratio of better than 1:10,000, which is essential for observing the subtle effects of neutrino oscillation.

1.3 Inverse Beta Decay

Chapter 2

Framework

2.1 Machine Learning

References

- [Kaj16] Takaaki Kajita. “Nobel Lecture: Discovery of atmospheric neutrino oscillations”. In: *Reviews of Modern Physics* 88.3 (July 2016). DOI: 10.1103/revmodphys.88.030501.