

1 Mathematical modeling of neuron's electric activity

In Section 1.1, the concept of *excitability* of neurons has been introduced: neurons are able to generate and transmit **electrical impulses** as reaction to external stimuli. These impulses are a direct consequence of rapid changes in intracellular and extracellular *ionic concentrations* of main elements such as Na^+ , Cl^- , K^+ , Ca^{2+} . The change in such concentrations determines a change in the potential crossed across the cell's membrane, and the formation of an **action potential**.

In this chapter, mathematical models to describe the behaviour of such quantities are presented. After having introduced the main assumptions of the model, the equivalent circuit formulation for cellular electrophysiology will be presented, along with different models to describe its electrical components. Finally, the main model to describe the propagation of the action potential between different neurons will be under study, namely the **cable equation model**. These models consist **ordinary differential equations (ODEs)** or **partial differential equations (PDEs)**, to be numerically solved through appropriate techniques.

1.1 Electrical activity in neurons




Figure 1: *Schematic model of a cellular membrane*

From a biological point of view, the different ionic concentrations, assumed inside and outside the neuron, give rise to a *passive transport* of ions, which moves accordingly to the *electrochemical gradient* (i.e. movement of positive charges from high to low potentials). The passage of ion species across the membrane can be allowed or denied by **ionic channels**, each of one designed for a specific ion. The passive transport is not the only mechanism of ionic transport. Indeed, thanks to special **ionic pumps**, the cell is able to perform an *active transport*, moving ions against the electrochemical gradient. In order to do this, the cell needs to consume energy, in form of *ATP*.

Let us consider a geometrical setting representing the relevant components of a neuron for its electrical activity: an intracellular region Ω_{in} and an extracellular

region Ω_{out} . The two regions are separated by the cell's membrane, presenting internal and external surfaces Σ_{in} and Σ_{out} . Finally, define the normal versors to Σ_{in} and Σ_{out} as \mathbf{n}_{in} and $\mathbf{n}_{out} = -\mathbf{n}_{in}$.

Under this setting, we define the **total ionic current density** of ion α as

$$J_{\alpha,i}^{TOT} = \mathbf{J}_{\alpha,i}^{TOT} \cdot \mathbf{n}_i = J_{\alpha,i}^{cond} + J_{\alpha,i}^{cap} \quad i = in, out \quad (1)$$

The total current density is formed by two contributions:

- **Conduction current** J_{α}^{cond} : the current formed by the conduction of ions along the ionic channels. We assume no loss of current inside the channel, so that $J_{\alpha,in}^{cond} = J_{\alpha,out}^{cond}$
- **Capacitive current** J_{α}^{cap} : the current formed by the charge accumulation in Σ_{in} and Σ_{out} . The charge accumulation makes the membrane behave like a *capacitor*. Thus, we assume

$$J_{\alpha,i}^{cap} = \frac{\partial \sigma_{\alpha,i}}{\partial t} \quad i = in, out \quad (2)$$

where $\sigma_{\alpha,i}$ is the **surface charge density**. Also in this case we require *wall's neutrality*

$$\sigma_{\alpha,in} = \sigma_{\alpha,out} \quad (3)$$

The neutrality conditions on conduction and capacitive current implies current conservation

$$J_{\alpha,in}^{TOT} = J_{\alpha,out}^{TOT} \quad (4)$$

Considering two points $\mathbf{x}' \in \Omega_{in}$ and $\mathbf{x}'' \in \Omega_{out}$ and a time interval $[0, T]$, we can consider the electric potentials $\psi^{in}(t) = \psi(\mathbf{x}', t)$ and $\psi^{out}(t) = \psi(\mathbf{x}'', t)$, assuming the potentials constant respectively in Ω_{in} and Ω_{out} . The **membrane potential** is then defined as

$$\psi_m(t) = \psi^{in}(t) - \psi^{out}(t) \quad t \in [0, T] \quad (5)$$

The existence of the membrane potential is consequence of the Faraday's equation (known as well as Maxwell's third equation)

$$\nabla \wedge \mathbf{E} = -\frac{\partial \mathbf{B}}{\partial t} \quad (6)$$

where \mathbf{E} and \mathbf{B} are the electric and magnetic field, respectively. In the microscopic cellular environment, it happens in practice that $|\frac{\partial \mathbf{B}}{\partial t}| \ll |\nabla \wedge \mathbf{E}|$, being the magnetic contribution negligible with respect to the electric one. This implies from (2) that $\nabla \wedge \mathbf{E} = 0 \implies \exists \psi s.t. \mathbf{E} = -\nabla \psi$.

If R_c and t_m are, respectively, the cell's radius and the membrane's thickness, in practice we have that their ratio $\eta_c := \frac{t_m}{R_c} \ll 1$. This implies that from a modeling point of view we can assume $t_m \sim 0$, i.e. we can neglect the thickness of the membrane and assume $\Sigma_{in} \sim \Sigma_{out} \sim \Sigma$. In practice, for example, realistic values of such quantities can be $R_c \sim 10^{-6}m$, $t_m \sim 10^{-9}m$ and thus $\eta_c \sim 10^{-3}$.

If ϵ_m is the membrane's *dielectric constant*, we can define the **specific capacitance** of the membrane as $c_m = \frac{\epsilon_m}{t_m}$. Assuming then that the surface charge densities have the form $\sigma_{in}(\mathbf{x}', t) = c_m \psi_m(\mathbf{x}, t)$ and $\sigma_{out}(\mathbf{x}'', t) = -c_m \psi_m(\mathbf{x}, t)$, where $\mathbf{x} \in \Sigma$. The capacitive current density (2) assumes the form

$$J_{\alpha,i}^{cap}(\mathbf{x}, t) = c_m \frac{\partial \psi_m(\mathbf{x}, t)}{\partial t} \quad t \in [0, T], \mathbf{x} \in \Sigma \quad (7)$$

1.2 ODE local models




Figure 2: *Equivalent circuit of the total current density as in eq. (1)*

Let us now consider a specific $\mathbf{x} \in \Sigma$. Then, equation (1) expresses a balance law for the current densities $J_{\alpha,i}^{TOT}, i = in, out$, which can be divided in two contributed (capactive and conductive), where the capacitive current assumes the form $J_{\alpha,i}^{cap}(t) = c_m \frac{\partial \psi_m(t)}{\partial t}$.

As for the conductive current, we consider a general form