



Sufficient Conditions for the Lipschitz Regularity of Mean-Field Optimal Controls

Benoît Bonnet

(in collaboration with F. Rossi)



Groupe de Travail de Calcul des Variations

March 29, 2021

Outline of the talk

Multi-agent systems and mean-field optimal control

Lipschitz feedbacks in classical OCPs

Lipschitz feedbacks in mean-field OCPs

Example in 1D – Maximisation of the variance

Conclusion and open problems

Outline of the talk

Multi-agent systems and mean-field optimal control

Lipschitz feedbacks in classical OCPs

Lipschitz feedbacks in mean-field OCPs

Example in 1D – Maximisation of the variance

Conclusion and open problems

Multi-agent systems – *Some illustrations*

Multi-agent system \rightsquigarrow **large** ensemble of **interacting** things



Introduction – *Pattern formation*

Central observation (pattern formation)

Simple **microscopic** rules \rightsquigarrow complex **macroscopic** structures!

- ◇ **Consensus** (everybody tends to agree on something) :
 \hookrightarrow *aggregation models in biology, herds analysis, etc...*
- ◇ **Flocking** (everybody goes in the same direction) :
 \hookrightarrow *flocks of birds, opinion models, etc..*
- ◇ **Shock-waves & mills** (periodic motions in the system) :
 \hookrightarrow *Unidimensional traffic networks, schools of fishes, etc...*

Multi-agent control (main themes)

1. First, **model** agent dynamics and analyse pattern formations,
2. Then, **stir** systems towards/away from relevant patterns.

Introduction – *Pattern formation*

Central observation (pattern formation)

Simple **microscopic** rules \rightsquigarrow complex **macroscopic** structures!

- ◇ **Consensus** (everybody tends to agree on something) :
 \hookrightarrow *aggregation models in biology, herds analysis, etc...*
- ◇ **Flocking** (everybody goes in the same direction) :
 \hookrightarrow *flocks of birds, opinion models, etc..*
- ◇ **Shock-waves & mills** (periodic motions in the system) :
 \hookrightarrow *Unidimensional traffic networks, schools of fishes, etc...*

Multi-agent control (main themes)

1. First, **model** agent dynamics and analyse pattern formations,
2. Then, **stir** systems towards/away from relevant patterns.

Introduction – *Pattern formation*

Central observation (pattern formation)

Simple **microscopic** rules \rightsquigarrow complex **macroscopic** structures!

- ◇ **Consensus** (everybody tends to agree on something) :
 \hookrightarrow *aggregation models in biology, herds analysis, etc...*
- ◇ **Flocking** (everybody goes in the same direction) :
 \hookrightarrow *flocks of birds, opinion models, etc..*
- ◇ **Shock-waves & mills** (periodic motions in the system) :
 \hookrightarrow *Unidimensional traffic networks, schools of fishes, etc...*

Multi-agent control (main themes)

1. First, **model** agent dynamics and analyse pattern formations,
2. Then, **stir** systems towards/away from relevant patterns.

Introduction – *Pattern formation*

Central observation (pattern formation)

Simple **microscopic** rules \rightsquigarrow complex **macroscopic** structures!

- ◇ **Consensus** (everybody tends to agree on something) :
 \hookrightarrow *aggregation models in biology, herds analysis, etc...*
- ◇ **Flocking** (everybody goes in the same direction) :
 \hookrightarrow *flocks of birds, opinion models, etc..*
- ◇ **Shock-waves & mills** (periodic motions in the system) :
 \hookrightarrow *Unidimensional traffic networks, schools of fishes, etc...*

Multi-agent control (main themes)

1. First, **model** agent dynamics and analyse pattern formations.
2. Then, **steer** systems towards/away from relevant patterns.

Introduction – *Pattern formation*

Central observation (pattern formation)

Simple **microscopic** rules \rightsquigarrow complex **macroscopic** structures!

- ◇ **Consensus** (everybody tends to agree on something) :
 \hookrightarrow *aggregation models in biology, herds analysis, etc...*
- ◇ **Flocking** (everybody goes in the same direction) :
 \hookrightarrow *flocks of birds, opinion models, etc..*
- ◇ **Shock-waves & mills** (periodic motions in the system) :
 \hookrightarrow *Unidimensional traffic networks, schools of fishes, etc...*

Multi-agent control (main themes)

1. First, **model** agent dynamics and **analyse** pattern formations,
2. Then, **stir** systems **towards/away** from relevant patterns.

Introduction – *Pattern formation*

Central observation (pattern formation)

Simple **microscopic** rules \rightsquigarrow complex **macroscopic** structures!

- ◇ **Consensus** (everybody tends to agree on something) :
 \hookrightarrow *aggregation models in biology, herds analysis, etc...*
- ◇ **Flocking** (everybody goes in the same direction) :
 \hookrightarrow *flocks of birds, opinion models, etc..*
- ◇ **Shock-waves & mills** (periodic motions in the system) :
 \hookrightarrow *Unidimensional traffic networks, schools of fishes, etc...*

Multi-agent control (main themes)

1. First, **model** agent dynamics and **analyse** pattern formations,
2. Then, **stir** systems **towards/away** from relevant patterns.

Introduction – *Pattern formation*

Central observation (pattern formation)

Simple **microscopic** rules \rightsquigarrow complex **macroscopic** structures!

- ◇ **Consensus** (everybody tends to agree on something) :
 \hookrightarrow *aggregation models in biology, herds analysis, etc...*
- ◇ **Flocking** (everybody goes in the same direction) :
 \hookrightarrow *flocks of birds, opinion models, etc..*
- ◇ **Shock-waves & mills** (periodic motions in the system) :
 \hookrightarrow *Unidimensional traffic networks, schools of fishes, etc...*

Multi-agent control (main themes)

1. First, **model** agent dynamics and **analyse** pattern formations,
2. Then, **stir** systems **towards/away** from relevant patterns.

Multi-agent control – *Finite dimensional setting*

Controlled multi-agent dynamics \rightsquigarrow **systems of ODEs**

$$\dot{x}_i(t) = \mathbf{v}_N(t, \mathbf{x}_N(t), x_i(t)) + u_i(t),$$

where

- ◇ $\mathbf{x}_N(\cdot) = (x_1(\cdot), \dots, x_N(\cdot)) \in \text{Lip}([0, T], (\mathbb{R}^d)^N)$ are the **states**,
- ◇ $\mathbf{v}_N : [0, T] \times (\mathbb{R}^d)^N \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a **non-local** drift,
- ◇ $u(\cdot) = (u_1(\cdot), \dots, u_N(\cdot)) \in L^\infty([0, T], U^N)$ are the **controls**.

\leftrightarrow Controllability, explicit synthesis, **optimal control** !

$$\left\{ \begin{array}{l} \min_{u(\cdot)} \left[\frac{1}{N} \sum_{i=1}^N \int_0^T \left(L_N(t, \mathbf{x}_N(t), x_i(t)) + \psi(u_i(t)) \right) dt + \varphi_N(\mathbf{x}_N(T)) \right] \\ \text{s.t.} \left\{ \begin{array}{l} \dot{x}_i(t) = \mathbf{v}_N(t, \mathbf{x}_N(t), x_i(t)) + u_i(t), \\ x_i(0) = x_i^0. \end{array} \right. \end{array} \right.$$

Multi-agent control – *Finite dimensional setting*

Controlled multi-agent dynamics \rightsquigarrow **systems of ODEs**

$$\dot{x}_i(t) = \mathbf{v}_N(t, \mathbf{x}_N(t), x_i(t)) + u_i(t),$$

where

- ◇ $\mathbf{x}_N(\cdot) = (x_1(\cdot), \dots, x_N(\cdot)) \in \text{Lip}([0, T], (\mathbb{R}^d)^N)$ are the **states**,
- ◇ $\mathbf{v}_N : [0, T] \times (\mathbb{R}^d)^N \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a **non-local** drift,
- ◇ $\mathbf{u}(\cdot) = (u_1(\cdot), \dots, u_N(\cdot)) \in L^\infty([0, T], U^N)$ are the **controls**.

\hookrightarrow Controllability, explicit synthesis, **optimal control** !

$$\begin{cases} \min_{\mathbf{u}(\cdot)} \left[\frac{1}{N} \sum_{i=1}^N \int_0^T \left(L_N(t, \mathbf{x}_N(t), x_i(t)) + \psi(u_i(t)) \right) dt + \varphi_N(\mathbf{x}_N(T)) \right] \\ \text{s.t.} \begin{cases} \dot{x}_i(t) = \mathbf{v}_N(t, \mathbf{x}_N(t), x_i(t)) + u_i(t), \\ x_i(0) = x_i^0. \end{cases} \end{cases}$$

Multi-agent control – *Finite dimensional setting*

Controlled multi-agent dynamics \rightsquigarrow **systems of ODEs**

$$\dot{x}_i(t) = \mathbf{v}_N(t, \mathbf{x}_N(t), x_i(t)) + u_i(t),$$

where

- ◇ $\mathbf{x}_N(\cdot) = (x_1(\cdot), \dots, x_N(\cdot)) \in \text{Lip}([0, T], (\mathbb{R}^d)^N)$ are the **states**,
- ◇ $\mathbf{v}_N : [0, T] \times (\mathbb{R}^d)^N \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a **non-local** drift,
- ◇ $\mathbf{u}(\cdot) = (u_1(\cdot), \dots, u_N(\cdot)) \in L^\infty([0, T], U^N)$ are the **controls**.

\hookrightarrow Controllability, explicit synthesis, **optimal control** !

$$\begin{cases} \min_{\mathbf{u}(\cdot)} \left[\frac{1}{N} \sum_{i=1}^N \int_0^T \left(L_N(t, \mathbf{x}_N(t), x_i(t)) + \psi(u_i(t)) \right) dt + \varphi_N(\mathbf{x}_N(T)) \right] \\ \text{s.t.} \begin{cases} \dot{x}_i(t) = \mathbf{v}_N(t, \mathbf{x}_N(t), x_i(t)) + u_i(t), \\ x_i(0) = x_i^0. \end{cases} \end{cases}$$

Multi-agent control – *Finite dimensional setting*

Controlled multi-agent dynamics \rightsquigarrow **systems of ODEs**

$$\dot{x}_i(t) = \mathbf{v}_N(t, \mathbf{x}_N(t), x_i(t)) + u_i(t),$$

where

- ◇ $\mathbf{x}_N(\cdot) = (x_1(\cdot), \dots, x_N(\cdot)) \in \text{Lip}([0, T], (\mathbb{R}^d)^N)$ are the **states**,
- ◇ $\mathbf{v}_N : [0, T] \times (\mathbb{R}^d)^N \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a **non-local** drift,
- ◇ $\mathbf{u}(\cdot) = (u_1(\cdot), \dots, u_N(\cdot)) \in L^\infty([0, T], U^N)$ are the **controls**.

\hookrightarrow Controllability, explicit synthesis, **optimal control** !

$$\begin{cases} \min_{\mathbf{u}(\cdot)} \left[\frac{1}{N} \sum_{i=1}^N \int_0^T \left(L_N(t, \mathbf{x}_N(t), x_i(t)) + \psi(u_i(t)) \right) dt + \varphi_N(\mathbf{x}_N(T)) \right] \\ \text{s.t.} \begin{cases} \dot{x}_i(t) = \mathbf{v}_N(t, \mathbf{x}_N(t), x_i(t)) + u_i(t), \\ x_i(0) = x_i^0. \end{cases} \end{cases}$$

Multi-agent control – *Finite dimensional setting*

Controlled multi-agent dynamics \rightsquigarrow **systems of ODEs**

$$\dot{x}_i(t) = \mathbf{v}_N(t, \mathbf{x}_N(t), x_i(t)) + u_i(t),$$

where

- ◇ $\mathbf{x}_N(\cdot) = (x_1(\cdot), \dots, x_N(\cdot)) \in \text{Lip}([0, T], (\mathbb{R}^d)^N)$ are the **states**,
- ◇ $\mathbf{v}_N : [0, T] \times (\mathbb{R}^d)^N \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a **non-local** drift,
- ◇ $\mathbf{u}(\cdot) = (u_1(\cdot), \dots, u_N(\cdot)) \in L^\infty([0, T], U^N)$ are the **controls**.

\hookrightarrow Controllability, explicit synthesis, **optimal control** !

$$\begin{cases} \min_{\mathbf{u}(\cdot)} \left[\frac{1}{N} \sum_{i=1}^N \int_0^T \left(L_N(t, \mathbf{x}_N(t), x_i(t)) + \psi(u_i(t)) \right) dt + \varphi_N(\mathbf{x}_N(T)) \right] \\ \text{s.t.} \begin{cases} \dot{x}_i(t) = \mathbf{v}_N(t, \mathbf{x}_N(t), x_i(t)) + u_i(t), \\ x_i(0) = x_i^0. \end{cases} \end{cases}$$

Multi-agent control – *Finite dimensional setting*

Controlled multi-agent dynamics \rightsquigarrow **systems of ODEs**

$$\dot{x}_i(t) = \mathbf{v}_N(t, \mathbf{x}_N(t), x_i(t)) + u_i(t),$$

where

- ◇ $\mathbf{x}_N(\cdot) = (x_1(\cdot), \dots, x_N(\cdot)) \in \text{Lip}([0, T], (\mathbb{R}^d)^N)$ are the **states**,
- ◇ $\mathbf{v}_N : [0, T] \times (\mathbb{R}^d)^N \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a **non-local** drift,
- ◇ $\mathbf{u}(\cdot) = (u_1(\cdot), \dots, u_N(\cdot)) \in L^\infty([0, T], U^N)$ are the **controls**.

\hookrightarrow Controllability, explicit synthesis, **optimal control** !


$$\begin{cases} \min_{\mathbf{u}(\cdot)} \left[\frac{1}{N} \sum_{i=1}^N \int_0^T \left(L_N(t, \mathbf{x}_N(t), x_i(t)) + \psi(u_i(t)) \right) dt + \varphi_N(\mathbf{x}_N(T)) \right] \\ \text{s.t.} \begin{cases} \dot{x}_i(t) = \mathbf{v}_N(t, \mathbf{x}_N(t), x_i(t)) + u_i(t), \\ x_i(0) = x_i^0. \end{cases} \end{cases}$$

Multi-agent control – *Dimension and mean-field limits*

First observation (dimensionality-related problems)

The number N of agents is usually **very large** \rightsquigarrow numerical issues

Idea: Approximation procedure using **mean-field** limits!



System of ODEs on N agents $(x_1, \dots, x_N) \in (\mathbb{R}^d)^N$

$$\left(\mu_N = \frac{1}{N} \sum_{i=1}^N \delta_{x_i} \right)$$


Single PDE on the density of agents $\mu : \mathbb{R}^d \rightarrow \mathbb{R}$.

Multi-agent control – *Dimension and mean-field limits*

First observation (dimensionality-related problems)

The number N of agents is usually **very large** \rightsquigarrow **numerical issues**

Idea: Approximation procedure using **mean-field** limits!



System of ODEs on N agents $(x_1, \dots, x_N) \in (\mathbb{R}^d)^N$

$$\left(\mu_N = \frac{1}{N} \sum_{i=1}^N \delta_{x_i} \right)$$


Single PDE on the density of agents $\mu : \mathbb{R}^d \rightarrow \mathbb{R}$.

Multi-agent control – *Dimension and mean-field limits*

First observation (dimensionality-related problems)

The number N of agents is usually **very large** \rightsquigarrow **numerical issues**

Idea: Approximation procedure using **mean-field** limits!



System of ODEs on N agents $(x_1, \dots, x_N) \in (\mathbb{R}^d)^N$

$$\left(\mu_N = \frac{1}{N} \sum_{i=1}^N \delta_{x_i} \right)$$


Single PDE on the density of agents $\mu : \mathbb{R}^d \rightarrow \mathbb{R}$.

Multi-agent control – *Dimension and mean-field limits*

First observation (dimensionality-related problems)

The number N of agents is usually **very large** \rightsquigarrow **numerical issues**

Idea: Approximation procedure using **mean-field** limits!



System of ODEs on N agents $(x_1, \dots, x_N) \in (\mathbb{R}^d)^N$

$$\left(\mu_N = \frac{1}{N} \sum_{i=1}^N \delta_{x_i} \right)$$

Single PDE on the density of agents $\mu : \mathbb{R}^d \rightarrow \mathbb{R}$.


Multi-agent control – *Towards mean-field optimal control*

Deterministic mean-field dynamics \rightsquigarrow **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}_x \left((v(t, \mu(t), \cdot) + u(t, \cdot)) \mu(t) \right) = 0,$$

where

- ◇ $\mu(t) \in \mathcal{P}(\mathbb{R}^d)$ is a **probability measure** \rightsquigarrow micro-macro ok!
- ◇ $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a **non-local** velocity field,
- ◇ the controls $u : [0, T] \times \mathbb{R}^d \rightarrow U$ are **vector fields**




Multi-agent control – *Towards mean-field optimal control*

Deterministic mean-field dynamics \rightsquigarrow **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}_x \left((v(t, \mu(t), \cdot) + u(t, \cdot)) \mu(t) \right) = 0,$$

where

- ◇ $\mu(t) \in \mathcal{P}(\mathbb{R}^d)$ is a **probability measure** \rightsquigarrow micro-macro ok!
- ◇ $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a **non-local** velocity field,
- ◇ the controls $u : [0, T] \times \mathbb{R}^d \rightarrow U$ are **vector fields**




Multi-agent control – *Towards mean-field optimal control*

Deterministic mean-field dynamics \rightsquigarrow **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}_x \left((v(t, \mu(t), \cdot) + u(t, \cdot)) \mu(t) \right) = 0,$$

where

- ◇ $\mu(t) \in \mathcal{P}(\mathbb{R}^d)$ is a **probability measure** \rightsquigarrow micro-macro ok!
- ◇ $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a **non-local** velocity field,
- ◇ the controls $u : [0, T] \times \mathbb{R}^d \rightarrow U$ are **vector fields**




Multi-agent control – *Towards mean-field optimal control*

Deterministic mean-field dynamics \rightsquigarrow **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}_x \left((v(t, \mu(t), \cdot) + u(t, \cdot)) \mu(t) \right) = 0,$$

where

- ◇ $\mu(t) \in \mathcal{P}(\mathbb{R}^d)$ is a **probability measure** \rightsquigarrow micro-macro ok!
- ◇ $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a **non-local** velocity field,
- ◇ the controls $u : [0, T] \times \mathbb{R}^d \rightarrow U$ are **vector fields**




Multi-agent control – *Towards mean-field optimal control*

Deterministic mean-field dynamics \rightsquigarrow **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}_x \left((v(t, \mu(t), \cdot) + u(t, \cdot)) \mu(t) \right) = 0,$$

where

- ◇ $\mu(t) \in \mathcal{P}(\mathbb{R}^d)$ is a **probability measure** \rightsquigarrow micro-macro ok!
- ◇ $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is a **non-local** velocity field,
- ◇ the controls $u : [0, T] \times \mathbb{R}^d \rightarrow U$ are **vector fields**



Multi-agent control – Mean-field optimal control problems

Idea: From (OCP_N) \rightsquigarrow **formally** associate an OCP on $\mu(\cdot)$

General mean-field optimal control problem (**MFOCP**)

$$\begin{cases} \min_{u(\cdot, \cdot)} \left[\int_0^T \int_{\mathbb{R}^d} \left(L(t, \mu(t), x) + \psi(u(t, x)) \right) d\mu(t)(x) dt + \varphi(\mu(T)) \right] \\ \text{s.t.} \begin{cases} \partial_t \mu(t) + \text{div}_x \left((v(t, \mu(t), \cdot) + u(t, \cdot)) \mu(t) \right) = 0, \\ \mu(0) = \mu^0, \end{cases} \end{cases}$$

where the dynamics and cost functionals satisfy

$$\begin{cases} v(t, \mu[\mathbf{x}], x) = \mathbf{v}_N(t, \mathbf{x}, x), & \varphi(\mu[\mathbf{x}]) = \varphi_N(\mathbf{x}), \\ L(t, \mu[\mathbf{x}], x) = \mathbf{L}_N(t, \mathbf{x}, x), \end{cases}$$

for every $\mathbf{x} := (x_1, \dots, x_N) \in (\mathbb{R}^d)^N$ and $\mu[\mathbf{x}] := \frac{1}{N} \sum_{i=1}^N \delta_{x_i}$.

Multi-agent control – Mean-field optimal control problems

Idea: From $(\text{OCP}_N) \rightsquigarrow$ **formally** associate an OCP on $\mu(\cdot)$

General mean-field optimal control problem (**MFOCP**)

$$\begin{cases} \min_{u(\cdot, \cdot)} \left[\int_0^T \int_{\mathbb{R}^d} \left(L(t, \mu(t), x) + \psi(u(t, x)) \right) d\mu(t)(x) dt + \varphi(\mu(T)) \right] \\ \text{s.t.} \begin{cases} \partial_t \mu(t) + \operatorname{div}_x \left((v(t, \mu(t), \cdot) + u(t, \cdot)) \mu(t) \right) = 0, \\ \mu(0) = \mu^0, \end{cases} \end{cases}$$

where the dynamics and cost functionals satisfy

$$\begin{cases} v(t, \mu[x], x) = \mathbf{v}_N(t, \mathbf{x}, x), & \varphi(\mu[x]) = \varphi_N(\mathbf{x}), \\ L(t, \mu[x], x) = \mathbf{L}_N(t, \mathbf{x}, x), \end{cases}$$

for every $\mathbf{x} := (x_1, \dots, x_N) \in (\mathbb{R}^d)^N$ and $\mu[\mathbf{x}] := \frac{1}{N} \sum_{i=1}^N \delta_{x_i}$.

Multi-agent control – Mean-field optimal control problems

Idea: From $(\text{OCP}_N) \rightsquigarrow$ **formally** associate an OCP on $\mu(\cdot)$

General mean-field optimal control problem (**MFOCP**)


$$\begin{cases} \min_{u(\cdot, \cdot)} \left[\int_0^T \int_{\mathbb{R}^d} \left(L(t, \mu(t), x) + \psi(u(t, x)) \right) d\mu(t)(x) dt + \varphi(\mu(T)) \right] \\ \text{s.t.} \begin{cases} \partial_t \mu(t) + \text{div}_x \left((v(t, \mu(t), \cdot) + u(t, \cdot)) \mu(t) \right) = 0, \\ \mu(0) = \mu^0, \end{cases} \end{cases}$$

where the dynamics and cost functionals satisfy


$$\begin{cases} v(t, \mu[\mathbf{x}], x) = \mathbf{v}_N(t, \mathbf{x}, x), & \varphi(\mu[\mathbf{x}]) = \varphi_N(\mathbf{x}), \\ L(t, \mu[\mathbf{x}], x) = \mathbf{L}_N(t, \mathbf{x}, x), \end{cases}$$

for every $\mathbf{x} := (x_1, \dots, x_N) \in (\mathbb{R}^d)^N$ and $\mu[\mathbf{x}] := \frac{1}{N} \sum_{i=1}^N \delta_{x_i}$.


Multi-agent control – *Summing up the whole scheme*




Multi-agent control – *Summing up the whole scheme*




Multi-agent control – *Summing up the whole scheme*




Multi-agent control – *Summing up the whole scheme*



Multi-agent control – *Summing up the whole scheme*



Multi-agent control – *Summing up the whole scheme*



Multi-agent systems – *Micro-macro correspondence*

Mean-field controls (back to discrete problems)

(MFOCP) on measures $\mu(\cdot) \implies \text{sol. } u^*(\cdot, \cdot)$

Plug in $\mu_N(\cdot) = \frac{1}{N} \sum_{i=1}^N \delta_{x_i(\cdot)}$

discrete classical trajectories $(x_1(\cdot), \dots, x_N(\cdot))$

Stirring $(x_1(\cdot), \dots, x_N(\cdot))$ with the mean-field control $u^*(\cdot, \cdot)$ yields

$$\dot{x}_i(t) = v_N(t, x_N(t), x_i(t)) + u^*(t, x_i(t))$$

May be ill-defined !

Well-posedness of continuity equations (general idea) [DpL'89 & A'04]

Arbitrary $\mu^0 \in \mathcal{P}(\mathbb{R}^d) \iff$ **Cauchy-Lipschitz** regularity on $u^*(t, \cdot)$

Multi-agent systems – *Micro-macro correspondence*

Mean-field controls (back to discrete problems)

(MFOCP) on measures $\mu(\cdot) \implies$ sol. $u^*(\cdot, \cdot)$

Plug in $\mu_N(\cdot) = \frac{1}{N} \sum_{i=1}^N \delta_{x_i(\cdot)}$

discrete classical trajectories $(x_1(\cdot), \dots, x_N(\cdot))$

Stirring $(x_1(\cdot), \dots, x_N(\cdot))$ with the mean-field control $u^*(\cdot, \cdot)$ yields

$$\dot{x}_i(t) = v_N(t, x_N(t), x_i(t)) + u^*(t, x_i(t))$$

May be ill-defined !

Well-posedness of continuity equations (general idea) [DpL'89 & A'04]

Arbitrary $\mu^0 \in \mathcal{P}(\mathbb{R}^d) \iff$ **Cauchy-Lipschitz** regularity on $u^*(t, \cdot)$

Multi-agent systems – *Micro-macro correspondence*

Mean-field controls (back to discrete problems)

(**MFOCP**) on measures $\mu(\cdot) \implies$ sol. $u^*(\cdot, \cdot)$

Plug in $\mu_N(\cdot) = \frac{1}{N} \sum_{i=1}^N \delta_{x_i(\cdot)}$

discrete classical trajectories $(x_1(\cdot), \dots, x_N(\cdot))$

Stirring $(x_1(\cdot), \dots, x_N(\cdot))$ with the mean-field control $u^*(\cdot, \cdot)$ yields

$$\dot{x}_i(t) = v_N(t, x_N(t), x_i(t)) + u^*(t, x_i(t))$$

May be ill-defined !

Well-posedness of continuity equations (general idea) [DpL'89 & A'04]

Arbitrary $\mu^0 \in \mathcal{P}(\mathbb{R}^d) \iff$ **Cauchy-Lipschitz** regularity on $u^*(t, \cdot)$

Multi-agent systems – *Micro-macro correspondence*

Mean-field controls (back to discrete problems)

(**MFOCP**) on measures $\mu(\cdot) \implies$ sol. $u^*(\cdot, \cdot)$

Plug in $\mu_N(\cdot) = \frac{1}{N} \sum_{i=1}^N \delta_{x_i(\cdot)}$

discrete classical trajectories $(x_1(\cdot), \dots, x_N(\cdot))$

Stirring $(x_1(\cdot), \dots, x_N(\cdot))$ with the mean-field control $u^*(\cdot, \cdot)$ yields

$$\dot{x}_i(t) = \mathbf{v}_N(t, \mathbf{x}_N(t), x_i(t)) + u^*(t, x_i(t))$$

May be ill-defined !

Well-posedness of continuity equations (general idea) [DpL'89 & A'04]

Arbitrary $\mu^0 \in \mathcal{P}(\mathbb{R}^d) \iff$ Cauchy-Lipschitz regularity on $u^*(t, \cdot)$

Multi-agent systems – *Micro-macro correspondence*

Mean-field controls (back to discrete problems)

(**MFOCP**) on measures $\mu(\cdot) \implies$ sol. $u^*(\cdot, \cdot)$

Plug in $\mu_N(\cdot) = \frac{1}{N} \sum_{i=1}^N \delta_{x_i(\cdot)}$

discrete classical trajectories $(x_1(\cdot), \dots, x_N(\cdot))$

Stirring $(x_1(\cdot), \dots, x_N(\cdot))$ with the mean-field control $u^*(\cdot, \cdot)$ yields

$$\dot{x}_i(t) = \mathbf{v}_N(t, \mathbf{x}_N(t), x_i(t)) + u^*(t, x_i(t))$$

May be ill-defined !

Well-posedness of continuity equations (general idea) [DpL'89 & A'04]

Arbitrary $\mu^0 \in \mathcal{P}(\mathbb{R}^d) \iff$ Cauchy-Lipschitz regularity on $u^*(t, \cdot)$

Multi-agent systems – *Micro-macro correspondence*

Mean-field controls (back to discrete problems)

(MFOCP) on measures $\mu(\cdot) \implies$ sol. $u^*(\cdot, \cdot)$

Plug in $\mu_N(\cdot) = \frac{1}{N} \sum_{i=1}^N \delta_{x_i(\cdot)}$

discrete classical trajectories $(x_1(\cdot), \dots, x_N(\cdot))$

Stirring $(x_1(\cdot), \dots, x_N(\cdot))$ with the mean-field control $u^*(\cdot, \cdot)$ yields

$$\dot{x}_i(t) = \mathbf{v}_N(t, \mathbf{x}_N(t), x_i(t)) + u^*(t, x_i(t))$$

May be ill-defined !

Well-posedness of continuity equations (general idea) [DpL'89 & A'04]

Arbitrary $\mu^0 \in \mathcal{P}(\mathbb{R}^d) \iff$ **Cauchy-Lipschitz** regularity on $u^*(t, \cdot)$

Multi-agent systems – *Lipschitz regularity of MFOC*

Theorem 1 (Sufficient Lip conditions for **(MFOCP)**)[B. & Rossi'21]

Suppose that the functionals in **(MFOCP)** are $C_{\text{loc}}^{2,1}$ in (μ, x) and that $\psi : U \rightarrow \mathbb{R}$ is **strongly convex** with $\lambda_\psi > 0$ large enough.

Then, there exist a constant $\mathcal{L}_U > 0$ and an optimal control $u^*(\cdot, \cdot)$ for **(MFOCP)** such that $\text{Lip}(u^*(t, \cdot); \mathbb{R}^d) \leq \mathcal{L}_U$ for a.e. $t \in [0, T]$.

Scheme of proof

1. Discretize **(MFOCP)** with $\mu_N^0 = \frac{1}{N} \sum_{i=1}^N \delta_{x_i^0} \rightharpoonup \mu^0 \rightsquigarrow \text{(OCP}_N\text{)}$
2. Build a sequence of **uniformly Lipschitz** $u_N^*(\cdot, \cdot)$ for **(OCP}_N\text{)}**
3. Show that $u^*(\cdot, \cdot) = \lim_{k \rightarrow +\infty} u_{N_k}^*(\cdot, \cdot)$ is optimal for **(MFOCP)**

\hookrightarrow Let us look into the case $N = 1$, i.e. classical OCPs !

Multi-agent systems – *Lipschitz regularity of MFOC*

Theorem 1 (Sufficient Lip conditions for **(MFOCP)**)[B. & Rossi'21]

Suppose that the functionals in **(MFOCP)** are $C_{\text{loc}}^{2,1}$ in (μ, x) and that $\psi : U \rightarrow \mathbb{R}$ is **strongly convex** with $\lambda_\psi > 0$ large enough.

Then, there exist a constant $\mathcal{L}_U > 0$ and an optimal control $u^*(\cdot, \cdot)$ for **(MFOCP)** such that $\text{Lip}(u^*(t, \cdot); \mathbb{R}^d) \leq \mathcal{L}_U$ for a.e. $t \in [0, T]$.

Scheme of proof

1. Discretize**(MFOCP)** with $\mu_N^0 = \frac{1}{N} \sum_{i=1}^N \delta_{x_i^0} \rightharpoonup \mu^0 \rightsquigarrow \text{(OCP}_N\text{)}$
2. Build a sequence of **uniformly Lipschitz** $u_N^*(\cdot, \cdot)$ for **(OCP}_N\text{)}**
3. Show that $u^*(\cdot, \cdot) = \lim_{k \rightarrow +\infty} u_{N_k}^*(\cdot, \cdot)$ is optimal for **(MFOCP)**

\hookrightarrow Let us look into the case $N = 1$, i.e. classical OCPs !

Multi-agent systems – *Lipschitz regularity of MFOC*

Theorem 1 (Sufficient Lip conditions for **(MFOCP)**)[B. & Rossi'21]

Suppose that the functionals in **(MFOCP)** are $C_{\text{loc}}^{2,1}$ in (μ, x) and that $\psi : U \rightarrow \mathbb{R}$ is **strongly convex** with $\lambda_\psi > 0$ large enough.

Then, there exist a constant $\mathcal{L}_U > 0$ and an optimal control $u^*(\cdot, \cdot)$ for **(MFOCP)** such that $\text{Lip}(u^*(t, \cdot); \mathbb{R}^d) \leq \mathcal{L}_U$ for a.e. $t \in [0, T]$.

Scheme of proof

1. Discretize **(MFOCP)** with $\mu_N^0 = \frac{1}{N} \sum_{i=1}^N \delta_{x_i^0} \rightharpoonup \mu^0 \rightsquigarrow \text{(OCP}_N\text{)}$
2. Build a sequence of **uniformly Lipschitz** $u_N^*(\cdot, \cdot)$ for **(OCP}_N\text{)}**
3. Show that $u^*(\cdot, \cdot) = \lim_{k \rightarrow +\infty} u_{N_k}^*(\cdot, \cdot)$ is optimal for **(MFOCP)**

\hookrightarrow Let us look into the case $N = 1$, i.e. classical OCPs !

Multi-agent systems – Lipschitz regularity of MFOC

Theorem 1 (Sufficient Lip conditions for **(MFOCP)**)[B. & Rossi'21]

Suppose that the functionals in **(MFOCP)** are $C_{\text{loc}}^{2,1}$ in (μ, x) and that $\psi : U \rightarrow \mathbb{R}$ is **strongly convex** with $\lambda_\psi > 0$ large enough.

Then, there exist a constant $\mathcal{L}_U > 0$ and an optimal control $u^*(\cdot, \cdot)$ for **(MFOCP)** such that $\text{Lip}(u^*(t, \cdot); \mathbb{R}^d) \leq \mathcal{L}_U$ for a.e. $t \in [0, T]$.

Scheme of proof

1. Discretize **(MFOCP)** with $\mu_N^0 = \frac{1}{N} \sum_{i=1}^N \delta_{x_i^0} \rightharpoonup \mu^0 \rightsquigarrow \textbf{(OCP}_N\textbf{)}$
2. Build a sequence of **uniformly Lipschitz** $u_N^*(\cdot, \cdot)$ for **(OCP}_N\textbf{)}**
3. Show that $u^*(\cdot, \cdot) = \lim_{k \rightarrow +\infty} u_{N_k}^*(\cdot, \cdot)$ is optimal for **(MFOCP)**

\hookrightarrow Let us look into the case $N = 1$, i.e. classical OCPs !

Multi-agent systems – Lipschitz regularity of MFOC

Theorem 1 (Sufficient Lip conditions for **(MFOCP)**)[B. & Rossi'21]

Suppose that the functionals in **(MFOCP)** are $C_{\text{loc}}^{2,1}$ in (μ, x) and that $\psi : U \rightarrow \mathbb{R}$ is **strongly convex** with $\lambda_\psi > 0$ large enough.

Then, there exist a constant $\mathcal{L}_U > 0$ and an optimal control $u^*(\cdot, \cdot)$ for **(MFOCP)** such that $\text{Lip}(u^*(t, \cdot); \mathbb{R}^d) \leq \mathcal{L}_U$ for a.e. $t \in [0, T]$.

Scheme of proof

1. Discretize **(MFOCP)** with $\mu_N^0 = \frac{1}{N} \sum_{i=1}^N \delta_{x_i^0} \rightharpoonup \mu^0 \rightsquigarrow$ **(OCP_N)**
2. Build a sequence of **uniformly Lipschitz** $u_N^*(\cdot, \cdot)$ for **(OCP_N)**
3. Show that $u^*(\cdot, \cdot) = \lim_{k \rightarrow +\infty} u_{N_k}^*(\cdot, \cdot)$ is optimal for **(MFOCP)**

\hookrightarrow Let us look into the case $N = 1$, i.e. classical OCPs !

Multi-agent systems – Lipschitz regularity of MFOC

Theorem 1 (Sufficient Lip conditions for **(MFOCP)**)[B. & Rossi'21]

Suppose that the functionals in **(MFOCP)** are $C_{\text{loc}}^{2,1}$ in (μ, x) and that $\psi : U \rightarrow \mathbb{R}$ is **strongly convex** with $\lambda_\psi > 0$ large enough.

Then, there exist a constant $\mathcal{L}_U > 0$ and an optimal control $u^*(\cdot, \cdot)$ for **(MFOCP)** such that $\text{Lip}(u^*(t, \cdot); \mathbb{R}^d) \leq \mathcal{L}_U$ for a.e. $t \in [0, T]$.

Scheme of proof

1. Discretize **(MFOCP)** with $\mu_N^0 = \frac{1}{N} \sum_{i=1}^N \delta_{x_i^0} \rightharpoonup \mu^0 \rightsquigarrow \textbf{(OCP}_N\textbf{)}$
2. Build a sequence of **uniformly Lipschitz** $u_N^*(\cdot, \cdot)$ for **(OCP}_N\textbf{)}**
3. Show that $u^*(\cdot, \cdot) = \lim_{k \rightarrow +\infty} u_{N_k}^*(\cdot, \cdot)$ is optimal for **(MFOCP)**

\hookrightarrow Let us look into the case $N = 1$, i.e. classical OCPs !

Multi-agent systems – Lipschitz regularity of MFOC

Theorem 1 (Sufficient Lip conditions for **(MFOCP)**)[B. & Rossi'21]

Suppose that the functionals in **(MFOCP)** are $C_{\text{loc}}^{2,1}$ in (μ, x) and that $\psi : U \rightarrow \mathbb{R}$ is **strongly convex** with $\lambda_\psi > 0$ large enough.

Then, there exist a constant $\mathcal{L}_U > 0$ and an optimal control $u^*(\cdot, \cdot)$ for **(MFOCP)** such that $\text{Lip}(u^*(t, \cdot); \mathbb{R}^d) \leq \mathcal{L}_U$ for a.e. $t \in [0, T]$.

Scheme of proof

1. Discretize **(MFOCP)** with $\mu_N^0 = \frac{1}{N} \sum_{i=1}^N \delta_{x_i^0} \rightharpoonup \mu^0 \rightsquigarrow \textbf{(OCP}_N\textbf{)}$
2. Build a sequence of **uniformly Lipschitz** $u_N^*(\cdot, \cdot)$ for **(OCP}_N\textbf{)}**
3. Show that $u^*(\cdot, \cdot) = \lim_{k \rightarrow +\infty} u_{N_k}^*(\cdot, \cdot)$ is optimal for **(MFOCP)**

↪ Let us look into the case $N = 1$, i.e. classical OCPs !

Outline of the talk

Multi-agent systems and mean-field optimal control

Lipschitz feedbacks in classical OCPs

Lipschitz feedbacks in mean-field OCPs

Example in 1D – Maximisation of the variance

Conclusion and open problems

Classical OCPs – Formulation of the problem

Let $(u^*(\cdot), x^*(\cdot))$ be an optimal pair for

$$(\text{OCP}) \quad \begin{cases} \min_{u(\cdot) \in \mathcal{U}} \left[\int_0^T \left(L(t, x(t)) + \psi(u(t)) \right) dt + \varphi(x(T)) \right] \\ \text{s.t.} \quad \begin{cases} \dot{x}(t) = f(t, x(t)) + u(t), \\ x(0) = x^0. \end{cases} \end{cases}$$

Main question (Lipschitz feedbacks)

When does there exist a map $\bar{u} : [0, T] \times \mathbb{R}^d \rightarrow U$ such that

- (i) $u^*(t) = \bar{u}(t, x^*(t))$ for all times $t \in [0, T]$,
- (ii) the map $\bar{u}(\cdot, \cdot)$ is locally optimal for (OCP),
- (iii) the application $x \in \mathbb{R}^d \mapsto \bar{u}(t, x) \in U$ is Lipschitz.

Hamilton-Jacobi approach & $C^{1,1}$ value function ?

→ Pontryagin approach following [Dontchev et.al 19] !

Classical OCPs – Formulation of the problem

Let $(u^*(\cdot), x^*(\cdot))$ be an optimal pair for

$$(\text{OCP}) \quad \begin{cases} \min_{u(\cdot) \in \mathcal{U}} \left[\int_0^T \left(L(t, x(t)) + \psi(u(t)) \right) dt + \varphi(x(T)) \right] \\ \text{s.t.} \quad \begin{cases} \dot{x}(t) = f(t, x(t)) + u(t), \\ x(0) = x^0. \end{cases} \end{cases}$$

Main question (Lipschitz feedbacks)

When does there exist a map $\bar{u} : [0, T] \times \mathbb{R}^d \rightarrow U$ such that

- (i) $u^*(t) = \bar{u}(t, x^*(t))$ for all times $t \in [0, T]$,
- (ii) the map $\bar{u}(\cdot, \cdot)$ is **locally optimal** for (OCP),
- (iii) the application $x \in \mathbb{R}^d \mapsto \bar{u}(t, x) \in U$ is Lipschitz.

Hamilton-Jacobi approach & $C^{1,1}$ value function ?

→ Pontryagin approach following [Dontchev et.al 19] !

Classical OCPs – Formulation of the problem

Let $(u^*(\cdot), x^*(\cdot))$ be an optimal pair for

$$(\text{OCP}) \quad \begin{cases} \min_{u(\cdot) \in \mathcal{U}} \left[\int_0^T \left(L(t, x(t)) + \psi(u(t)) \right) dt + \varphi(x(T)) \right] \\ \text{s.t.} \quad \begin{cases} \dot{x}(t) = f(t, x(t)) + u(t), \\ x(0) = x^0. \end{cases} \end{cases}$$

Main question (Lipschitz feedbacks)

When does there exist a map $\bar{u} : [0, T] \times \mathbb{R}^d \rightarrow U$ such that

- (i) $u^*(t) = \bar{u}(t, x^*(t))$ for all times $t \in [0, T]$,
- (ii) the map $\bar{u}(\cdot, \cdot)$ is **locally optimal** for (OCP),
- (iii) the application $x \in \mathbb{R}^d \mapsto \bar{u}(t, x) \in U$ is **Lipschitz**.

Hamilton-Jacobi approach & $C^{1,1}$ value function ?

→ Pontryagin approach following [Dontchev et.al 19] !

Classical OCPs – Formulation of the problem

Let $(u^*(\cdot), x^*(\cdot))$ be an optimal pair for

$$(\text{OCP}) \quad \begin{cases} \min_{u(\cdot) \in \mathcal{U}} \left[\int_0^T \left(L(t, x(t)) + \psi(u(t)) \right) dt + \varphi(x(T)) \right] \\ \text{s.t.} \quad \begin{cases} \dot{x}(t) = f(t, x(t)) + u(t), \\ x(0) = x^0. \end{cases} \end{cases}$$

Main question (Lipschitz feedbacks)

When does there exist a map $\bar{u} : [0, T] \times \mathbb{R}^d \rightarrow U$ such that

- (i) $u^*(t) = \bar{u}(t, x^*(t))$ for all times $t \in [0, T]$,
- (ii) the map $\bar{u}(\cdot, \cdot)$ is **locally optimal** for (OCP),
- (iii) the application $x \in \mathbb{R}^d \mapsto \bar{u}(t, x) \in U$ is **Lipschitz**.

Hamilton-Jacobi approach & $C^{1,1}$ value function ?

\hookrightarrow Pontryagin approach following [Dontchev et.al 19] !

Classical OCPs – Formulation of the problem

Let $(u^*(\cdot), x^*(\cdot))$ be an optimal pair for

$$(\text{OCP}) \quad \begin{cases} \min_{u(\cdot) \in \mathcal{U}} \left[\int_0^T \left(L(t, x(t)) + \psi(u(t)) \right) dt + \varphi(x(T)) \right] \\ \text{s.t.} \quad \begin{cases} \dot{x}(t) = f(t, x(t)) + u(t), \\ x(0) = x^0. \end{cases} \end{cases}$$

Main question (Lipschitz feedbacks)

When does there exist a map $\bar{u} : [0, T] \times \mathbb{R}^d \rightarrow U$ such that

- (i) $u^*(t) = \bar{u}(t, x^*(t))$ for all times $t \in [0, T]$,
- (ii) the map $\bar{u}(\cdot, \cdot)$ is **locally optimal** for (OCP),
- (iii) the application $x \in \mathbb{R}^d \mapsto \bar{u}(t, x) \in U$ is **Lipschitz**.

Hamilton-Jacobi approach & $C^{1,1}$ value function ?

↔ Pontryagin approach following [Dontchev et.al 19] !

Classical OCPs – Formulation of the problem

Let $(u^*(\cdot), x^*(\cdot))$ be an optimal pair for

$$(\text{OCP}) \quad \begin{cases} \min_{u(\cdot) \in \mathcal{U}} \left[\int_0^T \left(L(t, x(t)) + \psi(u(t)) \right) dt + \varphi(x(T)) \right] \\ \text{s.t.} \quad \begin{cases} \dot{x}(t) = f(t, x(t)) + u(t), \\ x(0) = x^0. \end{cases} \end{cases}$$

Main question (Lipschitz feedbacks)

When does there exist a map $\bar{u} : [0, T] \times \mathbb{R}^d \rightarrow U$ such that

- (i) $u^*(t) = \bar{u}(t, x^*(t))$ for all times $t \in [0, T]$,
- (ii) the map $\bar{u}(\cdot, \cdot)$ is **locally optimal** for (OCP),
- (iii) the application $x \in \mathbb{R}^d \mapsto \bar{u}(t, x) \in U$ is **Lipschitz**.

Hamilton-Jacobi approach & $C^{1,1}$ **value function** ?

↔ **Pontryagin** approach following [Dontchev et.al 19] !

Classical OCPs – Formulation of the problem

Let $(u^*(\cdot), x^*(\cdot))$ be an optimal pair for

$$(\text{OCP}) \quad \begin{cases} \min_{u(\cdot) \in \mathcal{U}} \left[\int_0^T \left(L(t, x(t)) + \psi(u(t)) \right) dt + \varphi(x(T)) \right] \\ \text{s.t.} \quad \begin{cases} \dot{x}(t) = f(t, x(t)) + u(t), \\ x(0) = x^0. \end{cases} \end{cases}$$

Main question (Lipschitz feedbacks)

When does there exist a map $\bar{u} : [0, T] \times \mathbb{R}^d \rightarrow U$ such that

- (i) $u^*(t) = \bar{u}(t, x^*(t))$ for all times $t \in [0, T]$,
- (ii) the map $\bar{u}(\cdot, \cdot)$ is **locally optimal** for (OCP),
- (iii) the application $x \in \mathbb{R}^d \mapsto \bar{u}(t, x) \in U$ is **Lipschitz**.

Hamilton-Jacobi approach & $C^{1,1}$ **value function** ?

↪ **Pontryagin** approach following [Dontchev et.al 19] !

Classical OCPs – *Necessary optimality conditions*

Idea: Apply 1st-order necessary optimality conditions \rightsquigarrow **PMP**

Theorem (Pontryagin's maximum principle)

Define the **Hamiltonian function** associated to **(OCP)** by

$$H(t, x, p, u) = \langle p, f(t, x) + u \rangle - L(t, x) - \psi(u).$$

Then, there exists a **costate curve** $p^*(\cdot) \in \text{Lip}([0, T], \mathbb{R}^d)$ such that the **Pontryagin triple** $(x^*(\cdot), p^*(\cdot), u^*(\cdot))$ satisfies

$$\begin{cases} \dot{x}^*(t) = \nabla_p H(t, x^*(t), p^*(t), u^*(t)), & x^*(0) = x^0, \\ \dot{p}^*(t) = -\nabla_x H(t, x^*(t), p^*(t), u^*(t)), & p^*(T) = -\nabla \varphi(x^*(T)), \end{cases}$$

and

$$u^*(t) \in \underset{\omega \in U}{\operatorname{argmax}} H(t, x^*(t), p^*(t), \omega),$$

for \mathcal{L}^1 -almost every $t \in [0, T]$.

Classical OCPs – *Necessary optimality conditions*

Idea: Apply 1st-order necessary optimality conditions \rightsquigarrow **PMP**

Theorem (Pontryagin's maximum principle)

Define the **Hamiltonian function** associated to **(OCP)** by

$$H(t, x, p, u) = \langle p, f(t, x) + u \rangle - L(t, x) - \psi(u).$$

Then, there exists a **costate curve** $p^*(\cdot) \in \text{Lip}([0, T], \mathbb{R}^d)$ such that the **Pontryagin triple** $(x^*(\cdot), p^*(\cdot), u^*(\cdot))$ satisfies

$$\begin{cases} \dot{x}^*(t) = \nabla_p H(t, x^*(t), p^*(t), u^*(t)), & x^*(0) = x^0, \\ \dot{p}^*(t) = -\nabla_x H(t, x^*(t), p^*(t), u^*(t)), & p^*(T) = -\nabla \varphi(x^*(T)), \end{cases}$$

and

$$u^*(t) \in \underset{\omega \in U}{\operatorname{argmax}} H(t, x^*(t), p^*(t), \omega),$$

for \mathcal{L}^1 -almost every $t \in [0, T]$.

Classical OCPs – *Necessary optimality conditions*

Idea: Apply 1st-order necessary optimality conditions \rightsquigarrow **PMP**

Theorem (Pontryagin's maximum principle)

Define the **Hamiltonian function** associated to **(OCP)** by

$$H(t, x, p, u) = \langle p, f(t, x) + u \rangle - L(t, x) - \psi(u).$$

Then, there exists a **costate curve** $p^*(\cdot) \in \text{Lip}([0, T], \mathbb{R}^d)$ such that the **Pontryagin triple** $(x^*(\cdot), p^*(\cdot), u^*(\cdot))$ satisfies

$$\begin{cases} \dot{x}^*(t) = \nabla_p H(t, x^*(t), p^*(t), u^*(t)), & x^*(0) = x^0, \\ \dot{p}^*(t) = -\nabla_x H(t, x^*(t), p^*(t), u^*(t)), & p^*(T) = -\nabla \varphi(x^*(T)), \end{cases}$$

and

$$u^*(t) \in \underset{\omega \in U}{\operatorname{argmax}} H(t, x^*(t), p^*(t), \omega),$$

for \mathcal{L}^1 -almost every $t \in [0, T]$.

Classical OCPs – *Necessary optimality conditions*

Idea: Apply 1st-order necessary optimality conditions \rightsquigarrow **PMP**

Theorem (Pontryagin's maximum principle)

Define the **Hamiltonian function** associated to **(OCP)** by

$$H(t, x, p, u) = \langle p, f(t, x) + u \rangle - L(t, x) - \psi(u).$$

Then, there exists a **costate curve** $p^*(\cdot) \in \text{Lip}([0, T], \mathbb{R}^d)$ such that the **Pontryagin triple** $(x^*(\cdot), p^*(\cdot), u^*(\cdot))$ satisfies

$$\begin{cases} \dot{x}^*(t) = \nabla_p H(t, x^*(t), p^*(t), u^*(t)), & x^*(0) = x^0, \\ \dot{p}^*(t) = -\nabla_x H(t, x^*(t), p^*(t), u^*(t)), & p^*(T) = -\nabla \varphi(x^*(T)), \end{cases}$$

and

$$u^*(t) \in \underset{\omega \in U}{\operatorname{argmax}} H(t, x^*(t), p^*(t), \omega),$$

for \mathcal{L}^1 -almost every $t \in [0, T]$.

Locally optimal Lipschitz feedback – *Statement*

Theorem (Pontryagin approach to Lipschitz feedbacks) [DKV'19]

Suppose that the data of **(OCP)** are $C_{\text{loc}}^{2,1}$, that $\psi(\cdot)$ is **strictly convex**, and that there exists $\rho > 0$ such that any **linearised pairs**

$$\dot{y}(t) = D_x f(t, x^*(t))y(t) + w(t), \quad u^*(t) + w(t) \in U$$

satisfies the following **uniform coercivity estimate** holds

$$\begin{aligned} & \langle \nabla_x^2 \varphi(x^*(T))y(T), y(T) \rangle \\ & - \int_0^T \langle \nabla_x^2 H(t, x^*(t), p^*(t), u^*(t))y(t), y(t) \rangle dt \\ & - \int_0^T \langle \nabla_u^2 H(t, x^*(t), p^*(t), u^*(t))w(t), w(t) \rangle dt \geq \rho \int_0^T |w(t)|^2 dt. \end{aligned}$$

Then, there exists a **locally optimal feedback** $\bar{u} \in \text{Lip}(\mathcal{N}, U)$ for **(OCP)**, where $\mathcal{N} \subset [0, T] \times \mathbb{R}^d$ is an ε -neigh. of $\text{Graph}(x^*(\cdot))$.

Locally optimal Lipschitz feedback – *Statement*

Theorem (Pontryagin approach to Lipschitz feedbacks) [DKV'19]

Suppose that the data of **(OCP)** are $C_{\text{loc}}^{2,1}$, that $\psi(\cdot)$ is **strictly convex**, and that there exists $\rho > 0$ such that any **linearised pairs**

$$\dot{y}(t) = D_x f(t, x^*(t))y(t) + w(t), \quad u^*(t) + w(t) \in U$$

satisfies the following **uniform coercivity estimate** holds

$$\begin{aligned} & \langle \nabla_x^2 \varphi(x^*(T))y(T), y(T) \rangle \\ & - \int_0^T \langle \nabla_x^2 H(t, x^*(t), p^*(t), u^*(t))y(t), y(t) \rangle dt \\ & - \int_0^T \langle \nabla_u^2 H(t, x^*(t), p^*(t), u^*(t))w(t), w(t) \rangle dt \geq \rho \int_0^T |w(t)|^2 dt. \end{aligned}$$

Then, there exists a **locally optimal feedback** $\bar{u} \in \text{Lip}(\mathcal{N}, U)$ for **(OCP)**, where $\mathcal{N} \subset [0, T] \times \mathbb{R}^d$ is an ϵ -neigh. of $\text{Graph}(x^*(\cdot))$.

Locally optimal Lipschitz feedback – *Statement*

Theorem (Pontryagin approach to Lipschitz feedbacks) [DKV'19]

Suppose that the data of **(OCP)** are $C_{\text{loc}}^{2,1}$, that $\psi(\cdot)$ is **strictly convex**, and that there exists $\rho > 0$ such that any **linearised pairs**

$$\dot{y}(t) = D_x f(t, x^*(t))y(t) + w(t), \quad u^*(t) + w(t) \in U$$

satisfies the following **uniform coercivity estimate** holds

$$\begin{aligned} & \langle \nabla_x^2 \varphi(x^*(T))y(T), y(T) \rangle \\ & - \int_0^T \langle \nabla_x^2 H(t, x^*(t), p^*(t), u^*(t))y(t), y(t) \rangle dt \\ & - \int_0^T \langle \nabla_u^2 H(t, x^*(t), p^*(t), u^*(t))w(t), w(t) \rangle dt \geq \rho \int_0^T |w(t)|^2 dt. \end{aligned}$$

Then, there exists a **locally optimal feedback** $\bar{u} \in \text{Lip}(\mathcal{N}, U)$ for **(OCP)**, where $\mathcal{N} \subset [0, T] \times \mathbb{R}^d$ is an ϵ -neigh. of $\text{Graph}(x^*(\cdot))$.

Locally optimal Lipschitz feedback – *Statement*

Theorem (Pontryagin approach to Lipschitz feedbacks) [DKV'19]

Suppose that the data of **(OCP)** are $C_{\text{loc}}^{2,1}$, that $\psi(\cdot)$ is **strictly convex**, and that there exists $\rho > 0$ such that any **linearised pairs**


$$\dot{y}(t) = D_x f(t, x^*(t))y(t) + w(t), \quad u^*(t) + w(t) \in U$$

satisfies the following **uniform coercivity estimate** holds

$$\begin{aligned} & \langle \nabla_x^2 \varphi(x^*(T))y(T), y(T) \rangle \\ & - \int_0^T \langle \nabla_x^2 H(t, x^*(t), p^*(t), u^*(t))y(t), y(t) \rangle dt \\ & - \int_0^T \langle \nabla_u^2 H(t, x^*(t), p^*(t), u^*(t))w(t), w(t) \rangle dt \geq \rho \int_0^T |w(t)|^2 dt. \end{aligned}$$


Then, there exists a **locally optimal feedback** $\bar{u} \in \text{Lip}(\mathcal{N}, U)$ for **(OCP)**, where $\mathcal{N} \subset [0, T] \times \mathbb{R}^d$ is an ϵ -neigh. of $\text{Graph}(x^*(\cdot))$.

Locally optimal Lipschitz feedback – *Illustration*




1. Consider the curve $x^*(\cdot)$ and the ϵ -neighbourhood \mathcal{N} ,
2. For $(\tau, \xi) \in [0, T] \times \mathcal{N}_\tau$, let $x^*_{(\tau,\xi)}(\cdot)$ be an optimal curve such that $x^*_{(\tau,\xi)}(\tau) = \xi$,
3. Then, $t \in [\tau, T] \mapsto \bar{u}(t, x^*_{(\tau,\xi)}(t)) \in U$ is **better** than every admissible open-loop control generating $x^*_{(\tau,\xi)}(\cdot)$.

Locally optimal Lipschitz feedback – *Illustration*




1. Consider the curve $x^*(\cdot)$ and the ϵ -neighbourhood \mathcal{N} ,
2. For $(\tau, \xi) \in [0, T] \times \mathcal{N}_\tau$, let $x^*_{(\tau,\xi)}(\cdot)$ be an optimal curve such that $x^*_{(\tau,\xi)}(\tau) = \xi$,
3. Then, $t \in [\tau, T] \mapsto \bar{u}(t, x^*_{(\tau,\xi)}(t)) \in U$ is **better** than every admissible open-loop control generating $x^*_{(\tau,\xi)}(\cdot)$.

Locally optimal Lipschitz feedback – *Illustration*



1. Consider the curve $x^*(\cdot)$ and the ϵ -neighbourhood \mathcal{N} ,
2. For $(\tau, \xi) \in [0, T] \times \mathcal{N}_\tau$, let $x^*_{(\tau,\xi)}(\cdot)$ be an optimal curve such that $x^*_{(\tau,\xi)}(\tau) = \xi$,
3. Then, $t \in [\tau, T] \mapsto \bar{u}(t, x^*_{(\tau,\xi)}(t)) \in U$ is **better** than every admissible open-loop control generating $x^*_{(\tau,\xi)}(\cdot)$.

Locally optimal Lipschitz feedback – Illustration



1. Consider the curve $x^*(\cdot)$ and the ϵ -neighbourhood \mathcal{N} ,
2. For $(\tau, \xi) \in [0, T] \times \mathcal{N}_\tau$, let $x_{(\tau, \xi)}^*(\cdot)$ be an optimal curve such that $x_{(\tau, \xi)}^*(\tau) = \xi$,
3. Then, $t \in [\tau, T] \mapsto \bar{u}(t, x_{(\tau, \xi)}^*(t)) \in U$ is **better** than every admissible open-loop control generating $x_{(\tau, \xi)}^*(\cdot)$.

Locally optimal Lipschitz feedback – Proof scheme (1)

Proof (General idea)

Show that the PMP system is **invertible** in a **Lipschitz way** with respect to perturbations of the form $x(\tau) = \xi \in \mathcal{N}_\tau$.

1. Rewrite the maximisation condition as

$$\nabla_u H(t, x^*(t), p^*(t), u^*(t)) \in N_U(u^*(t)).$$

2. Rewrite the PMP as the **dynamical differential inclusion**

$$0 \in F(x^*(\cdot), p^*(\cdot), u^*(\cdot)) + G(x^*(\cdot), p^*(\cdot), u^*(\cdot))$$

where F encodes the **state-costate dynamics** and G the **maximisation condition**.

↪ Quantitative version of the non-smooth IFT (**Robinson**)

Locally optimal Lipschitz feedback – Proof scheme (1)

Proof (General idea)

Show that the PMP system is **invertible** in a **Lipschitz way** with respect to perturbations of the form $x(\tau) = \xi \in \mathcal{N}_\tau$.

1. Rewrite the maximisation condition as

$$\nabla_u H(t, x^*(t), p^*(t), u^*(t)) \in N_U(u^*(t)).$$

2. Rewrite the PMP as the **dynamical differential inclusion**

$$0 \in F(x^*(\cdot), p^*(\cdot), u^*(\cdot)) + G(x^*(\cdot), p^*(\cdot), u^*(\cdot))$$

where F encodes the **state-costate dynamics** and G the **maximisation condition**.

↪ Quantitative version of the non-smooth IFT (**Robinson**)

Locally optimal Lipschitz feedback – Proof scheme (1)

Proof (General idea)

Show that the PMP system is **invertible** in a **Lipschitz way** with respect to perturbations of the form $x(\tau) = \xi \in \mathcal{N}_\tau$.

1. Rewrite the maximisation condition as

$$\nabla_u H(t, x^*(t), p^*(t), u^*(t)) \in N_U(u^*(t)).$$

2. Rewrite the PMP as the **dynamical differential inclusion**

$$0 \in F(x^*(\cdot), p^*(\cdot), u^*(\cdot)) + G(x^*(\cdot), p^*(\cdot), u^*(\cdot))$$

where F encodes the **state-costate dynamics** and G the **maximisation condition**.

↪ Quantitative version of the non-smooth IFT (**Robinson**)

Locally optimal Lipschitz feedback – Proof scheme (1)

Proof (General idea)

Show that the PMP system is **invertible** in a **Lipschitz way** with respect to perturbations of the form $x(\tau) = \xi \in \mathcal{N}_\tau$.

1. Rewrite the maximisation condition as

$$\nabla_u H(t, x^*(t), p^*(t), u^*(t)) \in N_U(u^*(t)).$$

2. Rewrite the PMP as the **dynamical differential inclusion**

$$0 \in F(x^*(\cdot), p^*(\cdot), u^*(\cdot)) + G(x^*(\cdot), p^*(\cdot), u^*(\cdot))$$

where F encodes the **state-costate dynamics** and G the **maximisation condition**.

↪ **Quantitative** version of the non-smooth IFT (**Robinson**)

Locally optimal Lipschitz feedback – Proof scheme (1)

Proof (General idea)

Show that the PMP system is **invertible** in a **Lipschitz way** with respect to perturbations of the form $x(\tau) = \xi \in \mathcal{N}_\tau$.

1. Rewrite the maximisation condition as

$$\nabla_u H(t, x^*(t), p^*(t), u^*(t)) \in N_U(u^*(t)).$$

2. Rewrite the PMP as the **dynamical differential inclusion**

$$0 \in F(x^*(\cdot), p^*(\cdot), u^*(\cdot)) + G(x^*(\cdot), p^*(\cdot), u^*(\cdot))$$

where F encodes the **state-costate dynamics** and G the **maximisation condition**.

↪ **Quantitative** version of the non-smooth IFT (**Robinson**)

Locally optimal Lipschitz feedback – *Proof scheme* (2)

3. The **partially linearised** dynamical inclusion

$$0 \in DF(x^*(\cdot), p^*(\cdot), u^*(\cdot))(y, q, w) + G(y, q, w)$$

is in fact the PMP of the **LQ problem** on $(y(\cdot), w(\cdot))$

$$\begin{cases} \min_{w(\cdot)} \left[\int_0^T \left(\langle A^*(t)y(t), y(t) \rangle + \langle B^*(t)w(t), w(t) \rangle \right) dt + \langle C^*(T)y(T), y(T) \rangle \right] \\ \text{s.t.} \begin{cases} \dot{y}(t) = D_x f(t, x^*(t))y(t) + w(t), \\ y(0) = 0, \quad u^*(t) + w(t) \in U, \end{cases} \end{cases}$$

with **costate** $q(\cdot)$, and where

$$A^* = -\nabla_x^2 H^*, \quad B^* = -\nabla_u^2 H^* \quad \text{and} \quad C^*(T) = \nabla_x^2 \varphi(x^*(T)).$$

4. Coercivity estimate: quantitative second-order convexity condition for the LQ problem \rightsquigarrow **Lipschitz inverse** [HD'93]

Locally optimal Lipschitz feedback – Proof scheme (2)

3. The **partially linearised** dynamical inclusion

$$0 \in DF(x^*(\cdot), p^*(\cdot), u^*(\cdot))(y, q, w) + G(y, q, w)$$

is in fact the PMP of the **LQ problem** on $(y(\cdot), w(\cdot))$

$$\begin{cases} \min_{w(\cdot)} \left[\int_0^T (\langle A^*(t)y(t), y(t) \rangle + \langle B^*(t)w(t), w(t) \rangle) dt + \langle C^*(T)y(T), y(T) \rangle \right] \\ \text{s.t.} \begin{cases} \dot{y}(t) = D_x f(t, x^*(t))y(t) + w(t), \\ y(0) = 0, \quad u^*(t) + w(t) \in U, \end{cases} \end{cases}$$

with **costate** $q(\cdot)$, and where

$$A^* = -\nabla_x^2 H^*, \quad B^* = -\nabla_u^2 H^* \quad \text{and} \quad C^*(T) = \nabla_x^2 \varphi(x^*(T)).$$

4. **Coercivity estimate**: quantitative **second-order convexity condition** for the LQ problem \rightsquigarrow **Lipschitz inverse!** [HD'93]

Locally optimal Lipschitz feedback – Proof scheme (2)

3. The **partially linearised** dynamical inclusion

$$0 \in DF(x^*(\cdot), p^*(\cdot), u^*(\cdot))(y, q, w) + G(y, q, w)$$

is in fact the PMP of the **LQ problem** on $(y(\cdot), w(\cdot))$

$$\begin{cases} \min_{w(\cdot)} \left[\int_0^T (\langle A^*(t)y(t), y(t) \rangle + \langle B^*(t)w(t), w(t) \rangle) dt + \langle C^*(T)y(T), y(T) \rangle \right] \\ \text{s.t.} \begin{cases} \dot{y}(t) = D_x f(t, x^*(t))y(t) + w(t), \\ y(0) = 0, \quad u^*(t) + w(t) \in U, \end{cases} \end{cases}$$

with **costate** $q(\cdot)$, and where

$$A^* = -\nabla_x^2 H^*, \quad B^* = -\nabla_u^2 H^* \quad \text{and} \quad C^*(T) = \nabla_x^2 \varphi(x^*(T)).$$

4. **Coercivity estimate**: quantitative **second-order convexity condition** for the LQ problem \rightsquigarrow **Lipschitz inverse!** [HD'93]

Locally optimal Lipschitz feedback – Proof scheme (2)

3. The **partially linearised** dynamical inclusion

$$0 \in DF(x^*(\cdot), p^*(\cdot), u^*(\cdot))(y, q, w) + G(y, q, w)$$

is in fact the PMP of the **LQ problem** on $(y(\cdot), w(\cdot))$

$$\begin{cases} \min_{w(\cdot)} \left[\int_0^T (\langle A^*(t)y(t), y(t) \rangle + \langle B^*(t)w(t), w(t) \rangle) dt + \langle C^*(T)y(T), y(T) \rangle \right] \\ \text{s.t.} \begin{cases} \dot{y}(t) = D_x f(t, x^*(t))y(t) + w(t), \\ y(0) = 0, \quad u^*(t) + w(t) \in U, \end{cases} \end{cases}$$

with **costate** $q(\cdot)$, and where

$$A^* = -\nabla_x^2 H^*, \quad B^* = -\nabla_u^2 H^* \quad \text{and} \quad C^*(T) = \nabla_x^2 \varphi(x^*(T)).$$

4. **Coercivity estimate:** quantitative **second-order convexity condition** for the LQ problem \rightsquigarrow **Lipschitz inverse!** [HD'93]

Outline of the talk

Multi-agent systems and mean-field optimal control

Lipschitz feedbacks in classical OCPs

Lipschitz feedbacks in mean-field OCPs

Example in 1D – Maximisation of the variance

Conclusion and open problems

Back to multi-agent systems – *Adaptation to (CO_N)*

Going back to general systems with $N \geq 1$ interacting agents

General multi-agent optimal control problem (**OCP_N**)

$$\begin{cases} \min_{\mathbf{u}(\cdot)} \left[\frac{1}{N} \sum_{i=1}^N \int_0^T \left(\mathbf{L}_N(t, \mathbf{x}_N(t), x_i(t)) + \psi(u_i(t)) \right) dt + \varphi_N(\mathbf{x}_N(T)) \right] \\ \text{s.t.} \begin{cases} \dot{x}_i(t) = \mathbf{v}_N(t, \mathbf{x}_N(t), x_i(t)) + u_i(t), \\ x_i(0) = x_i^0. \end{cases} \end{cases}$$

Idea: See (**OCP_N**) as a classical OCP in $(\mathbb{R}^d)^N$ with trajectory-control pairs $(\mathbf{x}(\cdot), \mathbf{u}(\cdot)) \in \text{Lip}([0, T], (\mathbb{R}^d)^N) \times L^\infty([0, T], U^N)$.

Question: How do we write a coercivity estimate adapted to the mean-field structure, i.e. **uniform in N** ? \rightsquigarrow **Wasserstein calculus!**

Back to multi-agent systems – *Adaptation to (CO_N)*

Going back to general systems with $N \geq 1$ interacting agents

General multi-agent optimal control problem (**OCP_N**)

$$\begin{cases} \min_{\mathbf{u}(\cdot)} \left[\frac{1}{N} \sum_{i=1}^N \int_0^T \left(\mathbf{L}_N(t, \mathbf{x}_N(t), x_i(t)) + \psi(u_i(t)) \right) dt + \varphi_N(\mathbf{x}_N(T)) \right] \\ \text{s.t.} \begin{cases} \dot{x}_i(t) = \mathbf{v}_N(t, \mathbf{x}_N(t), x_i(t)) + u_i(t), \\ x_i(0) = x_i^0. \end{cases} \end{cases}$$

Idea: See (**OCP_N**) as a classical OCP in $(\mathbb{R}^d)^N$ with trajectory-control pairs $(\mathbf{x}(\cdot), \mathbf{u}(\cdot)) \in \text{Lip}([0, T], (\mathbb{R}^d)^N) \times L^\infty([0, T], U^N)$.

Question: How do we write a coercivity estimate adapted to the mean-field structure, i.e. **uniform in N** ? \rightsquigarrow **Wasserstein calculus!**

Back to multi-agent systems – *Adaptation to (CO_N)*

Going back to general systems with $N \geq 1$ interacting agents

General multi-agent optimal control problem (**OCP_N**)

$$\begin{cases} \min_{\mathbf{u}(\cdot)} \left[\frac{1}{N} \sum_{i=1}^N \int_0^T \left(\mathbf{L}_N(t, \mathbf{x}_N(t), \mathbf{x}_i(t)) + \psi(\mathbf{u}_i(t)) \right) dt + \varphi_N(\mathbf{x}_N(T)) \right] \\ \text{s.t.} \begin{cases} \dot{\mathbf{x}}_i(t) = \mathbf{v}_N(t, \mathbf{x}_N(t), \mathbf{x}_i(t)) + \mathbf{u}_i(t), \\ \mathbf{x}_i(0) = \mathbf{x}_i^0. \end{cases} \end{cases}$$

Idea: See (**OCP_N**) as a classical OCP in $(\mathbb{R}^d)^N$ with trajectory-control pairs $(\mathbf{x}(\cdot), \mathbf{u}(\cdot)) \in \text{Lip}([0, T], (\mathbb{R}^d)^N) \times L^\infty([0, T], U^N)$.

Question: How do we write a coercivity estimate adapted to the mean-field structure, i.e. **uniform in N** ? \rightsquigarrow **Wasserstein calculus!**

Back to multi-agent systems – *Adaptation to (CO_N)*

Going back to general systems with $N \geq 1$ interacting agents

General multi-agent optimal control problem (**OCP_N**)

$$\begin{cases} \min_{\mathbf{u}(\cdot)} \left[\frac{1}{N} \sum_{i=1}^N \int_0^T \left(\mathbf{L}_N(t, \mathbf{x}_N(t), \mathbf{x}_i(t)) + \psi(\mathbf{u}_i(t)) \right) dt + \varphi_N(\mathbf{x}_N(T)) \right] \\ \text{s.t.} \begin{cases} \dot{\mathbf{x}}_i(t) = \mathbf{v}_N(t, \mathbf{x}_N(t), \mathbf{x}_i(t)) + \mathbf{u}_i(t), \\ \mathbf{x}_i(0) = \mathbf{x}_i^0. \end{cases} \end{cases}$$

Idea: See (**OCP_N**) as a classical OCP in $(\mathbb{R}^d)^N$ with trajectory-control pairs $(\mathbf{x}(\cdot), \mathbf{u}(\cdot)) \in \text{Lip}([0, T], (\mathbb{R}^d)^N) \times L^\infty([0, T], U^N)$.

Question: How do we write a coercivity estimate adapted to the mean-field structure, i.e. **uniform in N** ? \rightsquigarrow **Wasserstein calculus!**


Optimal transport toolbox – *The modern foundations*

Optimal transport problem (Kantorovich's formulation)

Given $\mu, \nu \in \mathcal{P}(\mathbb{R}^d)$ and a cost function $c : \mathbb{R}^{2d} \rightarrow (-\infty, +\infty]$, find a probability measure $\gamma^* \in \mathcal{P}(\mathbb{R}^{2d})$ s.t.

- (i) $\gamma^* \in \Gamma(\mu, \nu)$, i.e. $\pi_{\#}^1 \gamma^* = \mu$ and $\pi_{\#}^2 \gamma^* = \nu$
- (ii) γ^* solves the optimisation problem

$$\int_{\mathbb{R}^{2d}} c(x, y) d\gamma^*(x, y) = \min_{\gamma \in \Gamma(\mu, \nu)} \int_{\mathbb{R}^{2d}} c(x, y) d\gamma(x, y)$$




Optimal transport toolbox – *The modern foundations*

Optimal transport problem (Kantorovich's formulation)

Given $\mu, \nu \in \mathcal{P}(\mathbb{R}^d)$ and a cost function $c : \mathbb{R}^{2d} \rightarrow (-\infty, +\infty]$, find a probability measure $\gamma^* \in \mathcal{P}(\mathbb{R}^{2d})$ s.t.

- (i) $\gamma^* \in \Gamma(\mu, \nu)$, i.e. $\pi_{\#}^1 \gamma^* = \mu$ and $\pi_{\#}^2 \gamma^* = \nu$
- (ii) γ^* solves the optimisation problem

$$\int_{\mathbb{R}^{2d}} c(x, y) d\gamma^*(x, y) = \min_{\gamma \in \Gamma(\mu, \nu)} \int_{\mathbb{R}^{2d}} c(x, y) d\gamma(x, y)$$




Optimal transport toolbox – *The modern foundations*

Optimal transport problem (Kantorovich's formulation)

Given $\mu, \nu \in \mathcal{P}(\mathbb{R}^d)$ and a cost function $c : \mathbb{R}^{2d} \rightarrow (-\infty, +\infty]$, find a probability measure $\gamma^* \in \mathcal{P}(\mathbb{R}^{2d})$ s.t.

- (i) $\gamma^* \in \Gamma(\mu, \nu)$, i.e. $\pi_{\#}^1 \gamma^* = \mu$ and $\pi_{\#}^2 \gamma^* = \nu$
- (ii) γ^* solves the optimisation problem

$$\int_{\mathbb{R}^{2d}} c(x, y) d\gamma^*(x, y) = \min_{\gamma \in \Gamma(\mu, \nu)} \int_{\mathbb{R}^{2d}} c(x, y) d\gamma(x, y)$$




Optimal transport toolbox – *The modern foundations*

Optimal transport problem (Kantorovich's formulation)

Given $\mu, \nu \in \mathcal{P}(\mathbb{R}^d)$ and a cost function $c : \mathbb{R}^{2d} \rightarrow (-\infty, +\infty]$, find a probability measure $\gamma^* \in \mathcal{P}(\mathbb{R}^{2d})$ s.t.

- (i) $\gamma^* \in \Gamma(\mu, \nu)$, i.e. $\pi_{\#}^1 \gamma^* = \mu$ and $\pi_{\#}^2 \gamma^* = \nu$
- (ii) γ^* solves the optimisation problem

$$\int_{\mathbb{R}^{2d}} c(x, y) d\gamma^*(x, y) = \min_{\gamma \in \Gamma(\mu, \nu)} \int_{\mathbb{R}^{2d}} c(x, y) d\gamma(x, y)$$



Optimal transport toolbox – *Wasserstein spaces*

Idea: choose $c(x, y) := |x - y|^2 \rightsquigarrow$ **distance** between measures

Definition (Wasserstein distance)

The quantity

$$W_2(\mu, \nu) = \min_{\gamma \in \Gamma(\mu, \nu)} \left(\int_{\mathbb{R}^d} |x - y|^2 d\gamma(x, y) \right)^{1/2}$$

defines a **distance** over $\mathcal{P}_2(\mathbb{R}^d)$.

Theorem (Structure of Wasserstein spaces)

The metric space $(\mathcal{P}_2(\mathbb{R}^d), W_2)$

- (i) gives an **intrinsic meaning** to continuity equations [AGS'08]
- (ii) has a **weak Riemannian** structure [McCann'97, Otto'01]
- (iii) supports a **Levi-Civita connection** [AmbrosioGigli'08, Lott'07]

Optimal transport toolbox – *Wasserstein spaces*

Idea: choose $c(x, y) := |x - y|^2 \rightsquigarrow$ **distance** between measures

Definition (Wasserstein distance)

The quantity

$$W_2(\mu, \nu) = \min_{\gamma \in \Gamma(\mu, \nu)} \left(\int_{\mathbb{R}^d} |x - y|^2 d\gamma(x, y) \right)^{1/2}$$

defines a **distance** over $\mathcal{P}_2(\mathbb{R}^d)$.

Theorem (Structure of Wasserstein spaces)

The metric space $(\mathcal{P}_2(\mathbb{R}^d), W_2)$

- (i) gives an **intrinsic meaning** to continuity equations [AGS'08]
- (ii) has a **weak Riemannian** structure [McCann'97, Otto'01]
- (iii) supports a **Levi-Civita connection** [AmbrosioGigli'08, Lott'07]

Optimal transport toolbox – *Wasserstein spaces*

Idea: choose $c(x, y) := |x - y|^2 \rightsquigarrow$ **distance** between measures

Definition (Wasserstein distance)

The quantity

$$W_2(\mu, \nu) = \min_{\gamma \in \Gamma(\mu, \nu)} \left(\int_{\mathbb{R}^d} |x - y|^2 d\gamma(x, y) \right)^{1/2}$$

defines a **distance** over $\mathcal{P}_2(\mathbb{R}^d)$.

Theorem (Structure of Wasserstein spaces)

The metric space $(\mathcal{P}_2(\mathbb{R}^d), W_2)$

- (i) gives an **intrinsic meaning** to continuity equations [AGS'08]
- (ii) has a **weak Riemannian** structure [McCann'97, Otto'01]
- (iii) supports a **Levi-Civita connection** [AmbrosioGigli'08, Lott'07]

Optimal transport toolbox – *Wasserstein spaces*

Idea: choose $c(x, y) := |x - y|^2 \rightsquigarrow$ **distance** between measures

Definition (Wasserstein distance)

The quantity

$$W_2(\mu, \nu) = \min_{\gamma \in \Gamma(\mu, \nu)} \left(\int_{\mathbb{R}^d} |x - y|^2 d\gamma(x, y) \right)^{1/2}$$

defines a **distance** over $\mathcal{P}_2(\mathbb{R}^d)$.

Theorem (Structure of Wasserstein spaces)

The metric space $(\mathcal{P}_2(\mathbb{R}^d), W_2)$

- (i) gives an **intrinsic meaning** to continuity equations [AGS'08]
- (ii) has a **weak Riemannian** structure [McCann'97, Otto'01]
- (iii) supports a **Levi-Civita connection** [AmbrosioGigli'08, Lott'07]

Optimal transport toolbox – Wasserstein spaces

Idea: choose $c(x, y) := |x - y|^2 \rightsquigarrow$ **distance** between measures

Definition (Wasserstein distance)

The quantity

$$W_2(\mu, \nu) = \min_{\gamma \in \Gamma(\mu, \nu)} \left(\int_{\mathbb{R}^d} |x - y|^2 d\gamma(x, y) \right)^{1/2}$$

defines a **distance** over $\mathcal{P}_2(\mathbb{R}^d)$.

Theorem (Structure of Wasserstein spaces)

The metric space $(\mathcal{P}_2(\mathbb{R}^d), W_2)$

- (i) gives an **intrinsic meaning** to continuity equations [AGS'08]
- (ii) has a **weak Riemannian** structure [McCann'97, Otto'01]
- (iii) supports a **Levi-Civita connection** [AmbrosioGigli'08, Lott'07]

Optimal transport toolbox – *Wasserstein spaces*

Idea: choose $c(x, y) := |x - y|^2 \rightsquigarrow$ **distance** between measures

Definition (Wasserstein distance)

The quantity

$$W_2(\mu, \nu) = \min_{\gamma \in \Gamma(\mu, \nu)} \left(\int_{\mathbb{R}^d} |x - y|^2 d\gamma(x, y) \right)^{1/2}$$

defines a **distance** over $\mathcal{P}_2(\mathbb{R}^d)$.

Theorem (Structure of Wasserstein spaces)

The metric space $(\mathcal{P}_2(\mathbb{R}^d), W_2)$

- (i) gives an **intrinsic meaning** to continuity equations [AGS'08]
- (ii) has a **weak Riemannian** structure [McCann'97, Otto'01]
- (iii) supports a **Levi-Civita connection** [AmbrosioGigli'08, Lott'07]

Optimal transport toolbox – *Mean-field derivatives*

Wasserstein calculus (Heuristic definitions)

If $\varphi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ is “nice enough” at μ , there exists

- (i) a natural **gradient** map $\nabla_\mu \varphi(\mu) \in \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \subset L^2(\mathbb{R}^d, \mathbb{R}^d; \mu)$
- (ii) a **Hessian** bilinear form $\text{Hess } \varphi[\mu] : \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \times \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$

Special case (Empirical distributions)

If $\mu_N := \mu[\mathbf{x}] = \frac{1}{N} \sum_{i=1}^N \delta_{\mathbf{x}_i}$, then

$$\mathcal{T}_{\mu_N} \mathcal{P}_2(\mathbb{R}^d) \sim ((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N),$$

where $\langle \mathbf{x}, \mathbf{y} \rangle_N = \frac{1}{N} \sum_{i=1}^N \langle \mathbf{x}_i, \mathbf{y}_i \rangle$. Given $\varphi_N : \mathbf{x} \mapsto \varphi(\mu[\mathbf{x}])$, define

$$\mathbf{Grad}_x \varphi_N := (\nabla_\mu \varphi(\mu[\mathbf{x}])(\mathbf{x}_i))_{1 \leq i \leq N}, \quad \mathbf{Hess}_x \varphi_N[\mathbf{x}] := \text{Hess } \varphi[\mu_N].$$

Idea: Formulate **coercivity estimates** for (OCP_N) using **Wasserstein derivatives** in the natural Hilbert space $((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N)$

Optimal transport toolbox – *Mean-field derivatives*

Wasserstein calculus (Heuristic definitions)

If $\varphi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ is “nice enough” at μ , there exists

- (i) a natural **gradient** map $\nabla_\mu \varphi(\mu) \in \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \subset L^2(\mathbb{R}^d, \mathbb{R}^d; \mu)$
- (ii) a **Hessian** bilinear form $\text{Hess } \varphi[\mu] : \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \times \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$

Special case (Empirical distributions)

If $\mu_N := \mu[\mathbf{x}] = \frac{1}{N} \sum_{i=1}^N \delta_{x_i}$, then

$$\mathcal{T}_{\mu_N} \mathcal{P}_2(\mathbb{R}^d) \sim ((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N),$$

where $\langle \mathbf{x}, \mathbf{y} \rangle_N = \frac{1}{N} \sum_{i=1}^N \langle x_i, y_i \rangle$. Given $\varphi_N : \mathbf{x} \mapsto \varphi(\mu[\mathbf{x}])$, define

$$\text{Grad}_{\mathbf{x}} \varphi_N := (\nabla_\mu \varphi(\mu[\mathbf{x}])(x_i))_{1 \leq i \leq N}, \quad \text{Hess}_{\mathbf{x}} \varphi_N[\mathbf{x}] := \text{Hess } \varphi[\mu_N].$$

Idea: Formulate **coercivity estimates** for (OCP_N) using **Wasserstein derivatives** in the natural Hilbert space $((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N)$

Optimal transport toolbox – *Mean-field derivatives*

Wasserstein calculus (Heuristic definitions)

If $\varphi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ is “nice enough” at μ , there exists

- (i) a natural **gradient** map $\nabla_\mu \varphi(\mu) \in \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \subset L^2(\mathbb{R}^d, \mathbb{R}^d; \mu)$
- (ii) a **Hessian** bilinear form $\text{Hess } \varphi[\mu] : \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \times \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$

Special case (Empirical distributions)

If $\mu_N := \mu[\mathbf{x}] = \frac{1}{N} \sum_{i=1}^N \delta_{\mathbf{x}_i}$, then

$$\mathcal{T}_{\mu_N} \mathcal{P}_2(\mathbb{R}^d) \sim ((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N),$$

where $\langle \mathbf{x}, \mathbf{y} \rangle_N = \frac{1}{N} \sum_{i=1}^N \langle \mathbf{x}_i, \mathbf{y}_i \rangle$. Given $\varphi_N : \mathbf{x} \mapsto \varphi(\mu[\mathbf{x}])$, define

$$\text{Grad}_{\mathbf{x}} \varphi_N := (\nabla_\mu \varphi(\mu[\mathbf{x}])(\mathbf{x}_i))_{1 \leq i \leq N}, \quad \text{Hess}_{\mathbf{x}} \varphi_N[\mathbf{x}] := \text{Hess } \varphi[\mu_N].$$

Idea: Formulate **coercivity estimates** for (OCP_N) using **Wasserstein derivatives** in the natural Hilbert space $((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N)$

Optimal transport toolbox – *Mean-field derivatives*

Wasserstein calculus (Heuristic definitions)

If $\varphi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ is “nice enough” at μ , there exists

- (i) a natural **gradient** map $\nabla_\mu \varphi(\mu) \in \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \subset L^2(\mathbb{R}^d, \mathbb{R}^d; \mu)$
- (ii) a **Hessian** bilinear form $\text{Hess } \varphi[\mu] : \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \times \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$

Special case (Empirical distributions)

If $\mu_N := \mu[\mathbf{x}] = \frac{1}{N} \sum_{i=1}^N \delta_{\mathbf{x}_i}$, then

$$\mathcal{T}_{\mu_N} \mathcal{P}_2(\mathbb{R}^d) \sim ((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N),$$

where $\langle \mathbf{x}, \mathbf{y} \rangle_N = \frac{1}{N} \sum_{i=1}^N \langle \mathbf{x}_i, \mathbf{y}_i \rangle$. Given $\varphi_N : \mathbf{x} \mapsto \varphi(\mu[\mathbf{x}])$, define

$$\text{Grad}_{\mathbf{x}} \varphi_N := (\nabla_\mu \varphi(\mu[\mathbf{x}])(\mathbf{x}_i))_{1 \leq i \leq N}, \quad \text{Hess}_{\mathbf{x}} \varphi_N[\mathbf{x}] := \text{Hess } \varphi[\mu_N].$$

Idea: Formulate **coercivity estimates** for (OCP_N) using **Wasserstein derivatives** in the natural Hilbert space $((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N)$

Optimal transport toolbox – *Mean-field derivatives*

Wasserstein calculus (Heuristic definitions)

If $\varphi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ is “nice enough” at μ , there exists

- (i) a natural **gradient** map $\nabla_\mu \varphi(\mu) \in \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \subset L^2(\mathbb{R}^d, \mathbb{R}^d; \mu)$
- (ii) a **Hessian** bilinear form $\text{Hess } \varphi[\mu] : \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \times \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$

Special case (Empirical distributions)

If $\mu_N := \mu[\mathbf{x}] = \frac{1}{N} \sum_{i=1}^N \delta_{\mathbf{x}_i}$, then

$$\mathcal{T}_{\mu_N} \mathcal{P}_2(\mathbb{R}^d) \sim ((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N),$$

where $\langle \mathbf{x}, \mathbf{y} \rangle_N = \frac{1}{N} \sum_{i=1}^N \langle \mathbf{x}_i, \mathbf{y}_i \rangle$. Given $\varphi_N : \mathbf{x} \mapsto \varphi(\mu[\mathbf{x}])$, define

$$\text{Grad}_{\mathbf{x}} \varphi_N := (\nabla_\mu \varphi(\mu[\mathbf{x}])(\mathbf{x}_i))_{1 \leq i \leq N}, \quad \text{Hess}_{\mathbf{x}} \varphi_N[\mathbf{x}] := \text{Hess } \varphi[\mu_N].$$

Idea: Formulate **coercivity estimates** for (OCP_N) using **Wasserstein derivatives** in the natural Hilbert space $((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N)$

Optimal transport toolbox – *Mean-field derivatives*

Wasserstein calculus (Heuristic definitions)

If $\varphi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ is “nice enough” at μ , there exists

- (i) a natural **gradient** map $\nabla_\mu \varphi(\mu) \in \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \subset L^2(\mathbb{R}^d, \mathbb{R}^d; \mu)$
- (ii) a **Hessian** bilinear form $\text{Hess } \varphi[\mu] : \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \times \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$

Special case (Empirical distributions)

If $\mu_N := \mu[\mathbf{x}] = \frac{1}{N} \sum_{i=1}^N \delta_{\mathbf{x}_i}$, then

$$\mathcal{T}_{\mu_N} \mathcal{P}_2(\mathbb{R}^d) \sim ((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N),$$

where $\langle \mathbf{x}, \mathbf{y} \rangle_N = \frac{1}{N} \sum_{i=1}^N \langle \mathbf{x}_i, \mathbf{y}_i \rangle$. Given $\varphi_N : \mathbf{x} \mapsto \varphi(\mu[\mathbf{x}])$, define

$$\text{Grad}_{\mathbf{x}} \varphi_N := (\nabla_\mu \varphi(\mu[\mathbf{x}])(\mathbf{x}_i))_{1 \leq i \leq N}, \quad \text{Hess}_{\mathbf{x}} \varphi_N[\mathbf{x}] := \text{Hess } \varphi[\mu_N].$$

Idea: Formulate **coercivity estimates** for (OCP_N) using **Wasserstein derivatives** in the natural Hilbert space $((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N)$

Optimal transport toolbox – *Mean-field derivatives*

Wasserstein calculus (Heuristic definitions)

If $\varphi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ is “nice enough” at μ , there exists

- (i) a natural **gradient** map $\nabla_\mu \varphi(\mu) \in \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \subset L^2(\mathbb{R}^d, \mathbb{R}^d; \mu)$
- (ii) a **Hessian** bilinear form $\text{Hess } \varphi[\mu] : \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \times \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$

Special case (Empirical distributions)

If $\mu_N := \mu[\mathbf{x}] = \frac{1}{N} \sum_{i=1}^N \delta_{\mathbf{x}_i}$, then

$$\mathcal{T}_{\mu_N} \mathcal{P}_2(\mathbb{R}^d) \sim ((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N),$$

where $\langle \mathbf{x}, \mathbf{y} \rangle_N = \frac{1}{N} \sum_{i=1}^N \langle \mathbf{x}_i, \mathbf{y}_i \rangle$. Given $\varphi_N : \mathbf{x} \mapsto \varphi(\mu[\mathbf{x}])$, define

$$\mathbf{Grad}_\mathbf{x} \varphi_N := (\nabla_\mu \varphi(\mu[\mathbf{x}])(\mathbf{x}_i))_{1 \leq i \leq N}, \quad \mathbf{Hess}_\mathbf{x} \varphi_N[\mathbf{x}] := \text{Hess } \varphi[\mu_N].$$

Idea: Formulate **coercivity estimates** for (OCP_N) using **Wasserstein derivatives** in the natural Hilbert space $((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N)$

Optimal transport toolbox – *Mean-field derivatives*

Wasserstein calculus (Heuristic definitions)

If $\varphi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$ is “nice enough” at μ , there exists

- (i) a natural **gradient** map $\nabla_\mu \varphi(\mu) \in \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \subset L^2(\mathbb{R}^d, \mathbb{R}^d; \mu)$
- (ii) a **Hessian** bilinear form $\text{Hess } \varphi[\mu] : \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \times \mathcal{T}_\mu \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R}$

Special case (Empirical distributions)

If $\mu_N := \mu[\mathbf{x}] = \frac{1}{N} \sum_{i=1}^N \delta_{\mathbf{x}_i}$, then

$$\mathcal{T}_{\mu_N} \mathcal{P}_2(\mathbb{R}^d) \sim ((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N),$$

where $\langle \mathbf{x}, \mathbf{y} \rangle_N = \frac{1}{N} \sum_{i=1}^N \langle \mathbf{x}_i, \mathbf{y}_i \rangle$. Given $\varphi_N : \mathbf{x} \mapsto \varphi(\mu[\mathbf{x}])$, define

$$\mathbf{Grad}_x \varphi_N := (\nabla_\mu \varphi(\mu[\mathbf{x}])(\mathbf{x}_i))_{1 \leq i \leq N}, \quad \mathbf{Hess}_x \varphi_N[\mathbf{x}] := \text{Hess } \varphi[\mu_N].$$

Idea: Formulate **coercivity estimates** for (OCP_N) using **Wasserstein derivatives** in the natural Hilbert space $((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N)$

Lipschitz regularity in (\mathbf{OCP}_N) – *The coercivity condition*

Define the **mean-field Hamiltonian** associated to (\mathbf{OCP}_N)

$$\mathbb{H}_N(t, \mathbf{x}, \mathbf{r}, \mathbf{u}) = \frac{1}{N} \sum_{i=1}^N \left(\langle \mathbf{r}_i, \mathbf{v}_N(t, \mathbf{x}, \mathbf{x}_i) + \mathbf{u}_i \rangle - \mathbf{L}_N(t, \mathbf{x}, \mathbf{x}_i) - \psi(\mathbf{u}_i) \right)$$

and apply the **PMP** \rightsquigarrow **Pontryagin triple** $(\mathbf{x}_N^*(\cdot), \mathbf{r}_N^*(\cdot), \mathbf{u}_N^*(\cdot))$

Mean-field coercivity estimate (MFCE_N)

(MFCE_N) holds along $(\mathbf{x}_N^*(\cdot), \mathbf{r}_N^*(\cdot), \mathbf{u}_N^*(\cdot))$ iff there exists $\rho > 0$ s.t.

$$\begin{aligned} & \text{Hess}_{\mathbf{x}} \varphi_N[\mathbf{x}_N^*(T)](\mathbf{y}(T), \mathbf{y}(T)) \\ & - \int_0^T \text{Hess}_{\mathbf{x}} \mathbb{H}_N[t, \mathbf{x}_N^*(t), \mathbf{r}_N^*(t), \mathbf{u}_N^*(t)](\mathbf{y}(t), \mathbf{y}(t)) dt \\ & - \int_0^T \text{Hess}_{\mathbf{u}} \mathbb{H}_N[t, \mathbf{x}_N^*(t), \mathbf{r}_N^*(t), \mathbf{u}_N^*(t)](\mathbf{w}(t), \mathbf{w}(t)) dt \geq \rho \int_0^T |\mathbf{w}(t)|_N^2 dt, \end{aligned}$$

for any pair $(\mathbf{y}(\cdot), \mathbf{w}(\cdot))$ sol. the **mean-field linearised dynamics**.

Lipschitz regularity in (OCP_N) – *The coercivity condition*

Define the **mean-field Hamiltonian** associated to (OCP_N)

$$\mathbb{H}_N(t, \mathbf{x}, \mathbf{r}, \mathbf{u}) = \frac{1}{N} \sum_{i=1}^N \left(\langle \mathbf{r}_i, \mathbf{v}_N(t, \mathbf{x}, \mathbf{x}_i) + \mathbf{u}_i \rangle - \mathbf{L}_N(t, \mathbf{x}, \mathbf{x}_i) - \psi(\mathbf{u}_i) \right)$$

and apply the **PMP** \rightsquigarrow **Pontryagin triple** $(\mathbf{x}_N^*(\cdot), \mathbf{r}_N^*(\cdot), \mathbf{u}_N^*(\cdot))$

Mean-field coercivity estimate (MFCE_N)

(MFCE_N) holds along $(\mathbf{x}_N^*(\cdot), \mathbf{r}_N^*(\cdot), \mathbf{u}_N^*(\cdot))$ iff there exists $\rho > 0$ s.t.

$$\begin{aligned} & \text{Hess}_{\mathbf{x}} \varphi_N[\mathbf{x}_N^*(T)](\mathbf{y}(T), \mathbf{y}(T)) \\ & - \int_0^T \text{Hess}_{\mathbf{x}} \mathbb{H}_N[t, \mathbf{x}_N^*(t), \mathbf{r}_N^*(t), \mathbf{u}_N^*(t)](\mathbf{y}(t), \mathbf{y}(t)) dt \\ & - \int_0^T \text{Hess}_{\mathbf{u}} \mathbb{H}_N[t, \mathbf{x}_N^*(t), \mathbf{r}_N^*(t), \mathbf{u}_N^*(t)](\mathbf{w}(t), \mathbf{w}(t)) dt \geq \rho \int_0^T \|\mathbf{w}(t)\|_N^2 dt, \end{aligned}$$

for any pair $(\mathbf{y}(\cdot), \mathbf{w}(\cdot))$ sol. the **mean-field linearised dynamics**.

Lipschitz regularity in (OCP_N) – *The coercivity condition*

Define the **mean-field Hamiltonian** associated to (OCP_N)

$$\mathbb{H}_N(t, \mathbf{x}, \mathbf{r}, \mathbf{u}) = \frac{1}{N} \sum_{i=1}^N \left(\langle \mathbf{r}_i, \mathbf{v}_N(t, \mathbf{x}, \mathbf{x}_i) + \mathbf{u}_i \rangle - \mathbf{L}_N(t, \mathbf{x}, \mathbf{x}_i) - \psi(\mathbf{u}_i) \right)$$

and apply the **PMP** \rightsquigarrow **Pontryagin triple** $(\mathbf{x}_N^*(\cdot), \mathbf{r}_N^*(\cdot), \mathbf{u}_N^*(\cdot))$

Mean-field coercivity estimate (MFCE_N)

(MFCE_N) holds along $(\mathbf{x}_N^*(\cdot), \mathbf{r}_N^*(\cdot), \mathbf{u}_N^*(\cdot))$ iff there exists $\rho > 0$ s.t.

$$\begin{aligned} & \text{Hess}_{\mathbf{x}} \varphi_N[\mathbf{x}_N^*(T)](\mathbf{y}(T), \mathbf{y}(T)) \\ & - \int_0^T \text{Hess}_{\mathbf{x}} \mathbb{H}_N[t, \mathbf{x}_N^*(t), \mathbf{r}_N^*(t), \mathbf{u}_N^*(t)](\mathbf{y}(t), \mathbf{y}(t)) dt \\ & - \int_0^T \text{Hess}_{\mathbf{u}} \mathbb{H}_N[t, \mathbf{x}_N^*(t), \mathbf{r}_N^*(t), \mathbf{u}_N^*(t)](\mathbf{w}(t), \mathbf{w}(t)) dt \geq \rho \int_0^T \|\mathbf{w}(t)\|_N^2 dt, \end{aligned}$$

for any pair $(\mathbf{y}(\cdot), \mathbf{w}(\cdot))$ sol. the **mean-field linearised dynamics**.

Lipschitz regularity in (OCP_N) – Convergence of feedbacks

Theorem 2 (Convergence of Lipschitz feedbacks)[B. & Rossi'21]

Suppose that for all $N \geq 1$, there exists an **optimal Pontryagin triple** $(\mathbf{x}_N^*(\cdot), \mathbf{r}_N^*(\cdot), \mathbf{u}_N^*(\cdot))$ for (OCP_N) which satisfies (MFCE_N) for a given constant $\rho > 0$ **independent of** $N \geq 1$.

Then, there exists $\mathcal{L}_U > 0$ and $\mathbf{u}_N^*(\cdot, \cdot) \in L^\infty([0, T], \text{Lip}(\mathbb{R}^d, U))$ s.t.

- (i) $\mathbf{u}_N^*(t, \mathbf{x}_i^*(t)) = \mathbf{u}_i^*(t)$ for \mathcal{L}^1 -almost every $t \in [0, T]$.
- (ii) The maps $\mathbf{x} \in \mathbb{R}^d \mapsto \mathbf{u}_N^*(t, \mathbf{x}) \in U$ are \mathcal{L}_U -Lipschitz.
- (iii) For $p \in (d, +\infty)$, the cluster points of $(\mathbf{u}_N^*(\cdot, \cdot))$ in the weak-topology of $L^2([0, T], W_{\text{loc}}^{1,p}(\mathbb{R}^d, U))$ are **optimal** for **(MFOCP)**.

1. By adapting [DKV'19] in $((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N)$ there exists a **locally optimal feedback** mapping $\bar{\mathbf{u}}_N : \mathcal{N} \rightarrow U^N$ s.t.

$$|\bar{\mathbf{u}}(t, \mathbf{y}) - \bar{\mathbf{u}}(t, \mathbf{x})|_N \leq \mathcal{L}_U |\mathbf{y} - \mathbf{x}|_N \quad \text{and} \quad \mathbf{u}_N^*(t) = \bar{\mathbf{u}}(t, \mathbf{x}_N^*(t)).$$

Lipschitz regularity in (OCP_N) – Convergence of feedbacks

Theorem 2 (Convergence of Lipschitz feedbacks)[B. & Rossi'21]

Suppose that for all $N \geq 1$, there exists an **optimal Pontryagin triple** $(\mathbf{x}_N^*(\cdot), \mathbf{r}_N^*(\cdot), \mathbf{u}_N^*(\cdot))$ for (OCP_N) which satisfies (MFCE_N) for a given constant $\rho > 0$ **independent of** $N \geq 1$.

Then, there exists $\mathcal{L}_U > 0$ and $\mathbf{u}_N^*(\cdot, \cdot) \in L^\infty([0, T], \text{Lip}(\mathbb{R}^d, U))$ s.t.

- (i) $\mathbf{u}_N^*(t, \mathbf{x}_i^*(t)) = \mathbf{u}_i^*(t)$ for \mathcal{L}^1 -almost every $t \in [0, T]$.
- (ii) The maps $\mathbf{x} \in \mathbb{R}^d \mapsto \mathbf{u}_N^*(t, \mathbf{x}) \in U$ are \mathcal{L}_U -Lipschitz.
- (iii) For $p \in (d, +\infty)$, the cluster points of $(\mathbf{u}_N^*(\cdot, \cdot))$ in the weak-topology of $L^2([0, T], W_{\text{loc}}^{1,p}(\mathbb{R}^d, U))$ are **optimal** for **(MFOCP)**.

1. By adapting [DKV'19] in $((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N)$ there exists a **locally optimal feedback** mapping $\bar{\mathbf{u}}_N : \mathcal{N} \rightarrow U^N$ s.t.

$$|\bar{\mathbf{u}}(t, \mathbf{y}) - \bar{\mathbf{u}}(t, \mathbf{x})|_N \leq \mathcal{L}_U |\mathbf{y} - \mathbf{x}|_N \quad \text{and} \quad \mathbf{u}_N^*(t) = \bar{\mathbf{u}}(t, \mathbf{x}_N^*(t)).$$

Lipschitz regularity in (OCP_N) – Convergence of feedbacks

Theorem 2 (Convergence of Lipschitz feedbacks)[B. & Rossi'21]

Suppose that for all $N \geq 1$, there exists an **optimal Pontryagin triple** $(\mathbf{x}_N^*(\cdot), \mathbf{r}_N^*(\cdot), \mathbf{u}_N^*(\cdot))$ for (OCP_N) which satisfies (MFCE_N) for a given constant $\rho > 0$ **independent of** $N \geq 1$.

Then, there exists $\mathcal{L}_U > 0$ and $\mathbf{u}_N^*(\cdot, \cdot) \in L^\infty([0, T], \text{Lip}(\mathbb{R}^d, U))$ s.t.

- (i) $\mathbf{u}_N^*(t, \mathbf{x}_i^*(t)) = \mathbf{u}_i^*(t)$ for \mathcal{L}^1 -almost every $t \in [0, T]$.
- (ii) The maps $\mathbf{x} \in \mathbb{R}^d \mapsto \mathbf{u}_N^*(t, \mathbf{x}) \in U$ are \mathcal{L}_U -Lipschitz.
- (iii) For $p \in (d, +\infty)$, the cluster points of $(\mathbf{u}_N^*(\cdot, \cdot))$ in the weak-topology of $L^2([0, T], W_{\text{loc}}^{1,p}(\mathbb{R}^d, U))$ are **optimal** for **(MFOCP)**.

1. By adapting [DKV'19] in $((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N)$ there exists a **locally optimal feedback** mapping $\bar{\mathbf{u}}_N : \mathcal{N} \rightarrow U^N$ s.t.

$$|\bar{\mathbf{u}}(t, \mathbf{y}) - \bar{\mathbf{u}}(t, \mathbf{x})|_N \leq \mathcal{L}_U |\mathbf{y} - \mathbf{x}|_N \quad \text{and} \quad \mathbf{u}_N^*(t) = \bar{\mathbf{u}}(t, \mathbf{x}_N^*(t)).$$

Lipschitz regularity in (OCP_N) – Convergence of feedbacks

Theorem 2 (Convergence of Lipschitz feedbacks)[B. & Rossi'21]

Suppose that for all $N \geq 1$, there exists an **optimal Pontryagin triple** $(\mathbf{x}_N^*(\cdot), \mathbf{r}_N^*(\cdot), \mathbf{u}_N^*(\cdot))$ for (OCP_N) which satisfies (MFCE_N) for a given constant $\rho > 0$ **independent of** $N \geq 1$.

Then, there exists $\mathcal{L}_U > 0$ and $\mathbf{u}_N^*(\cdot, \cdot) \in L^\infty([0, T], \text{Lip}(\mathbb{R}^d, U))$ s.t.

- (i) $\mathbf{u}_N^*(t, \mathbf{x}_i^*(t)) = \mathbf{u}_i^*(t)$ for \mathcal{L}^1 -almost every $t \in [0, T]$.
- (ii) The maps $\mathbf{x} \in \mathbb{R}^d \mapsto \mathbf{u}_N^*(t, \mathbf{x}) \in U$ are \mathcal{L}_U -Lipschitz.
- (iii) For $p \in (d, +\infty)$, the cluster points of $(\mathbf{u}_N^*(\cdot, \cdot))$ in the weak-topology of $L^2([0, T], W_{\text{loc}}^{1,p}(\mathbb{R}^d, U))$ are **optimal** for **(MFOCP)**.

1. By adapting [DKV'19] in $((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N)$ there exists a **locally optimal feedback** mapping $\bar{\mathbf{u}}_N : \mathcal{N} \rightarrow U^N$ s.t.

$$|\bar{\mathbf{u}}(t, \mathbf{y}) - \bar{\mathbf{u}}(t, \mathbf{x})|_N \leq \mathcal{L}_U |\mathbf{y} - \mathbf{x}|_N \quad \text{and} \quad \mathbf{u}_N^*(t) = \bar{\mathbf{u}}(t, \mathbf{x}_N^*(t)).$$

Lipschitz regularity in (OCP_N) – Convergence of feedbacks

Theorem 2 (Convergence of Lipschitz feedbacks)[B. & Rossi'21]

Suppose that for all $N \geq 1$, there exists an **optimal Pontryagin triple** $(\mathbf{x}_N^*(\cdot), \mathbf{r}_N^*(\cdot), \mathbf{u}_N^*(\cdot))$ for (OCP_N) which satisfies (MFCE_N) for a given constant $\rho > 0$ **independent of** $N \geq 1$.

Then, there exists $\mathcal{L}_U > 0$ and $\mathbf{u}_N^*(\cdot, \cdot) \in L^\infty([0, T], \text{Lip}(\mathbb{R}^d, U))$ s.t.

- (i) $\mathbf{u}_N^*(t, \mathbf{x}_i^*(t)) = \mathbf{u}_i^*(t)$ for \mathcal{L}^1 -almost every $t \in [0, T]$.
- (ii) The maps $\mathbf{x} \in \mathbb{R}^d \mapsto \mathbf{u}_N^*(t, \mathbf{x}) \in U$ are \mathcal{L}_U -**Lipschitz**.
- (iii) For $p \in (d, +\infty)$, the cluster points of $(\mathbf{u}_N^*(\cdot, \cdot))$ in the weak-topology of $L^2([0, T], W_{\text{loc}}^{1,p}(\mathbb{R}^d, U))$ are **optimal** for **(MFOCP)**.

1. By adapting [DKV'19] in $((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N)$ there exists a **locally optimal feedback** mapping $\bar{\mathbf{u}}_N : \mathcal{N} \rightarrow U^N$ s.t.

$$|\bar{\mathbf{u}}(t, \mathbf{y}) - \bar{\mathbf{u}}(t, \mathbf{x})|_N \leq \mathcal{L}_U |\mathbf{y} - \mathbf{x}|_N \quad \text{and} \quad \mathbf{u}_N^*(t) = \bar{\mathbf{u}}(t, \mathbf{x}_N^*(t)).$$

Lipschitz regularity in (OCP_N) – Convergence of feedbacks

Theorem 2 (Convergence of Lipschitz feedbacks)[B. & Rossi'21]

Suppose that for all $N \geq 1$, there exists an **optimal Pontryagin triple** $(\mathbf{x}_N^*(\cdot), \mathbf{r}_N^*(\cdot), \mathbf{u}_N^*(\cdot))$ for (OCP_N) which satisfies (MFCE_N) for a given constant $\rho > 0$ **independent of** $N \geq 1$.

Then, there exists $\mathcal{L}_U > 0$ and $\mathbf{u}_N^*(\cdot, \cdot) \in L^\infty([0, T], \text{Lip}(\mathbb{R}^d, U))$ s.t.

- (i) $\mathbf{u}_N^*(t, \mathbf{x}_i^*(t)) = \mathbf{u}_i^*(t)$ for \mathcal{L}^1 -almost every $t \in [0, T]$.
- (ii) The maps $\mathbf{x} \in \mathbb{R}^d \mapsto \mathbf{u}_N^*(t, \mathbf{x}) \in U$ are \mathcal{L}_U -**Lipschitz**.
- (iii) For $p \in (d, +\infty)$, the cluster points of $(\mathbf{u}_N^*(\cdot, \cdot))$ in the weak-topology of $L^2([0, T], W_{\text{loc}}^{1,p}(\mathbb{R}^d, U))$ are **optimal** for **(MFOCP)**.

1. By adapting [DKV'19] in $((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N)$ there exists a **locally optimal feedback** mapping $\bar{\mathbf{u}}_N : \mathcal{N} \rightarrow U^N$ s.t.

$$|\bar{\mathbf{u}}(t, \mathbf{y}) - \bar{\mathbf{u}}(t, \mathbf{x})|_N \leq \mathcal{L}_U |\mathbf{y} - \mathbf{x}|_N \quad \text{and} \quad \mathbf{u}_N^*(t) = \bar{\mathbf{u}}(t, \mathbf{x}_N^*(t)).$$

Lipschitz regularity in (OCP_N) – Convergence of feedbacks

Theorem 2 (Convergence of Lipschitz feedbacks)[B. & Rossi'21]

Suppose that for all $N \geq 1$, there exists an **optimal Pontryagin triple** $(\mathbf{x}_N^*(\cdot), \mathbf{r}_N^*(\cdot), \mathbf{u}_N^*(\cdot))$ for (OCP_N) which satisfies (MFCE_N) for a given constant $\rho > 0$ **independent of** $N \geq 1$.

Then, there exists $\mathcal{L}_U > 0$ and $\mathbf{u}_N^*(\cdot, \cdot) \in L^\infty([0, T], \text{Lip}(\mathbb{R}^d, U))$ s.t.

- (i) $\mathbf{u}_N^*(t, \mathbf{x}_i^*(t)) = \mathbf{u}_i^*(t)$ for \mathcal{L}^1 -almost every $t \in [0, T]$.
- (ii) The maps $\mathbf{x} \in \mathbb{R}^d \mapsto \mathbf{u}_N^*(t, \mathbf{x}) \in U$ are \mathcal{L}_U -**Lipschitz**.
- (iii) For $p \in (d, +\infty)$, the cluster points of $(\mathbf{u}_N^*(\cdot, \cdot))$ in the weak-topology of $L^2([0, T], W_{\text{loc}}^{1,p}(\mathbb{R}^d, U))$ are **optimal** for **(MFOCP)**.

1. By adapting [DKV'19] in $((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N)$ there exists a **locally optimal feedback** mapping $\bar{\mathbf{u}}_N : \mathcal{N} \rightarrow U^N$ s.t.

$$|\bar{\mathbf{u}}(t, \mathbf{y}) - \bar{\mathbf{u}}(t, \mathbf{x})|_N \leq \mathcal{L}_U |\mathbf{y} - \mathbf{x}|_N \quad \text{and} \quad \mathbf{u}_N^*(t) = \bar{\mathbf{u}}(t, \mathbf{x}_N^*(t)).$$

Lipschitz regularity in (OCP_N) – Convergence of feedbacks

Theorem 2 (Convergence of Lipschitz feedbacks)[B. & Rossi'21]

Suppose that for all $N \geq 1$, there exists an **optimal Pontryagin triple** $(\mathbf{x}_N^*(\cdot), \mathbf{r}_N^*(\cdot), \mathbf{u}_N^*(\cdot))$ for (OCP_N) which satisfies (MFCE_N) for a given constant $\rho > 0$ **independent of** $N \geq 1$.

Then, there exists $\mathcal{L}_U > 0$ and $\mathbf{u}_N^*(\cdot, \cdot) \in L^\infty([0, T], \text{Lip}(\mathbb{R}^d, U))$ s.t.

- (i) $\mathbf{u}_N^*(t, \mathbf{x}_i^*(t)) = \mathbf{u}_i^*(t)$ for \mathcal{L}^1 -almost every $t \in [0, T]$.
- (ii) The maps $\mathbf{x} \in \mathbb{R}^d \mapsto \mathbf{u}_N^*(t, \mathbf{x}) \in U$ are \mathcal{L}_U -**Lipschitz**.
- (iii) For $p \in (d, +\infty)$, the cluster points of $(\mathbf{u}_N^*(\cdot, \cdot))$ in the weak-topology of $L^2([0, T], W_{\text{loc}}^{1,p}(\mathbb{R}^d, U))$ are **optimal** for **(MFOCP)**.

1. By adapting [DKV'19] in $((\mathbb{R}^d)^N, \langle \cdot, \cdot \rangle_N)$ there exists a **locally optimal feedback** mapping $\bar{\mathbf{u}}_N : \mathcal{N} \rightarrow U^N$ s.t.

$$|\bar{\mathbf{u}}(t, \mathbf{y}) - \bar{\mathbf{u}}(t, \mathbf{x})|_N \leq \mathcal{L}_U |\mathbf{y} - \mathbf{x}|_N \quad \text{and} \quad \mathbf{u}_N^*(t) = \bar{\mathbf{u}}(t, \mathbf{x}_N^*(t)).$$

Lipschitz regularity in (OCP_N) – Proof of Theorem 2

2. The **agent-based** feedbacks $\bar{u}_i : [0, T] \times \mathcal{N}_i \rightarrow \mathbb{R}^d$ defined by

$$\bar{u}_i(t, x) := \bar{u}_i\left(t, x_1^*(t), \dots, x_{i-1}^*(t), x, x_{i+1}^*(t), \dots, x_N^*(t)\right),$$

are **Lipschitz** on \mathcal{N}_i , i.e. $|\bar{u}_i(t, y) - \bar{u}_i(t, x)| \leq \mathcal{L}_U |x - y|$.

\hookrightarrow Collection of **feedback-neighbourhood** pairs $\{\bar{u}_i(\cdot, \cdot), \mathcal{N}_i\}_{i=1}^N$.

3. Build a **vector field** $u_N^* : [0, T] \times \mathbb{R}^d \rightarrow U$ by **patching together** the collection $\{\bar{u}_i(\cdot, \cdot), \mathcal{N}_i\}_{i=1}^N \oplus$ **Lip extension**.

\hookrightarrow Sequence $(u_N^*(\cdot, \cdot))$ of **loc. optimal feedbacks** with a **uniform Lipschitz regularity** in the space variable.

4. It holds $u_{N_k}^*(\cdot, \cdot) \rightharpoonup u^*(\cdot, \cdot)$ where $\text{Lip}(u^*(t, \cdot), \mathbb{R}^d) \leq \mathcal{L}_U$ and $u^*(\cdot, \cdot)$ is **optimal** for **(MFOCP)** $\rightsquigarrow \Gamma$ -conv. [FS'14, FLOS'19] \square

Lipschitz regularity in (OCP_N) – Proof of Theorem 2

2. The **agent-based** feedbacks $\bar{u}_i : [0, T] \times \mathcal{N}_i \rightarrow \mathbb{R}^d$ defined by

$$\bar{u}_i(t, \mathbf{x}) := \bar{\mathbf{u}}_i\left(t, \mathbf{x}_1^*(t), \dots, \mathbf{x}_{i-1}^*(t), \mathbf{x}, \mathbf{x}_{i+1}^*(t), \dots, \mathbf{x}_N^*(t)\right),$$

are **Lipschitz** on \mathcal{N}_i , i.e. $|\bar{u}_i(t, \mathbf{y}) - \bar{u}_i(t, \mathbf{x})| \leq \mathcal{L}_U |\mathbf{x} - \mathbf{y}|$.

↪ Collection of **feedback-neighbourhood** pairs $\{\bar{u}_i(\cdot, \cdot), \mathcal{N}_i\}_{i=1}^N$.

3. Build a **vector field** $u_N^* : [0, T] \times \mathbb{R}^d \rightarrow U$ by **patching together** the collection $\{\bar{u}_i(\cdot, \cdot), \mathcal{N}_i\}_{i=1}^N \oplus$ **Lip extension**.

↪ Sequence $(u_N^*(\cdot, \cdot))$ of **loc. optimal feedbacks** with a **uniform Lipschitz regularity** in the space variable.

4. It holds $u_{N_k}^*(\cdot, \cdot) \rightharpoonup u^*(\cdot, \cdot)$ where $\text{Lip}(u^*(t, \cdot), \mathbb{R}^d) \leq \mathcal{L}_U$ and $u^*(\cdot, \cdot)$ is **optimal** for **(MFOCP)** $\rightsquigarrow \Gamma$ -conv. [FS'14, FLOS'19] \square

Lipschitz regularity in (OCP_N) – Proof of Theorem 2

2. The **agent-based** feedbacks $\bar{u}_i : [0, T] \times \mathcal{N}_i \rightarrow \mathbb{R}^d$ defined by

$$\bar{u}_i(t, \mathbf{x}) := \bar{\mathbf{u}}_i\left(t, \mathbf{x}_1^*(t), \dots, \mathbf{x}_{i-1}^*(t), \mathbf{x}, \mathbf{x}_{i+1}^*(t), \dots, \mathbf{x}_N^*(t)\right),$$

are **Lipschitz** on \mathcal{N}_i , i.e. $|\bar{u}_i(t, \mathbf{y}) - \bar{u}_i(t, \mathbf{x})| \leq \mathcal{L}_U |\mathbf{x} - \mathbf{y}|$.

↪ Collection of **feedback-neighbourhood** pairs $\{\bar{u}_i(\cdot, \cdot), \mathcal{N}_i\}_{i=1}^N$.

3. Build a **vector field** $u_N^* : [0, T] \times \mathbb{R}^d \rightarrow U$ by **patching together** the collection $\{\bar{u}_i(\cdot, \cdot), \mathcal{N}_i\}_{i=1}^N \oplus \text{Lip extension}$.

↪ Sequence $(u_N^*(\cdot, \cdot))$ of **loc. optimal** feedbacks with a **uniform Lipschitz regularity** in the space variable.

4. It holds $u_{N_k}^*(\cdot, \cdot) \rightharpoonup u^*(\cdot, \cdot)$ where $\text{Lip}(u^*(t, \cdot), \mathbb{R}^d) \leq \mathcal{L}_U$ and $u^*(\cdot, \cdot)$ is **optimal** for **(MFOCP)** $\rightsquigarrow \Gamma$ -conv. [FS'14, FLOS'19] \square

Lipschitz regularity in (OCP_N) – Proof of Theorem 2

2. The **agent-based** feedbacks $\bar{u}_i : [0, T] \times \mathcal{N}_i \rightarrow \mathbb{R}^d$ defined by

$$\bar{u}_i(t, x) := \bar{u}_i\left(t, x_1^*(t), \dots, x_{i-1}^*(t), x, x_{i+1}^*(t), \dots, x_N^*(t)\right),$$

are **Lipschitz** on \mathcal{N}_i , i.e. $|\bar{u}_i(t, y) - \bar{u}_i(t, x)| \leq \mathcal{L}_U |x - y|$.

↪ Collection of **feedback-neighbourhood** pairs $\{\bar{u}_i(\cdot, \cdot), \mathcal{N}_i\}_{i=1}^N$.

3. Build a **vector field** $u_N^* : [0, T] \times \mathbb{R}^d \rightarrow U$ by **patching together** the collection $\{\bar{u}_i(\cdot, \cdot), \mathcal{N}_i\}_{i=1}^N \oplus \text{Lip extension}$.

↪ Sequence $(u_N^*(\cdot, \cdot))$ of **loc. optimal feedbacks** with a **uniform Lipschitz regularity** in the space variable.

4. It holds $u_{N_k}^*(\cdot, \cdot) \rightharpoonup u^*(\cdot, \cdot)$ where $\text{Lip}(u^*(t, \cdot), \mathbb{R}^d) \leq \mathcal{L}_U$ and $u^*(\cdot, \cdot)$ is **optimal** for $(\text{MFOCP}) \rightsquigarrow \Gamma\text{-conv}$. [FS'14, FLOS'19] \square

Lipschitz regularity in (OCP_N) – Proof of Theorem 2

2. The **agent-based** feedbacks $\bar{u}_i : [0, T] \times \mathcal{N}_i \rightarrow \mathbb{R}^d$ defined by

$$\bar{u}_i(t, x) := \bar{u}_i\left(t, x_1^*(t), \dots, x_{i-1}^*(t), x, x_{i+1}^*(t), \dots, x_N^*(t)\right),$$

are **Lipschitz** on \mathcal{N}_i , i.e. $|\bar{u}_i(t, y) - \bar{u}_i(t, x)| \leq \mathcal{L}_U |x - y|$.

↪ Collection of **feedback-neighbourhood** pairs $\{\bar{u}_i(\cdot, \cdot), \mathcal{N}_i\}_{i=1}^N$.

3. Build a **vector field** $u_N^* : [0, T] \times \mathbb{R}^d \rightarrow U$ by **patching together** the collection $\{\bar{u}_i(\cdot, \cdot), \mathcal{N}_i\}_{i=1}^N \oplus$ **Lip extension**.

↪ Sequence $(u_N^*(\cdot, \cdot))$ of **loc. optimal feedbacks** with a **uniform Lipschitz regularity** in the space variable.

4. It holds $u_{N_k}^*(\cdot, \cdot) \rightharpoonup u^*(\cdot, \cdot)$ where $\text{Lip}(u^*(t, \cdot), \mathbb{R}^d) \leq \mathcal{L}_U$ and $u^*(\cdot, \cdot)$ is **optimal** for **(MFOCP)** $\rightsquigarrow \Gamma$ -conv. [FS'14, FLOS'19] \square

Lipschitz regularity in (OCP_N) – Proof of Theorem 2

2. The **agent-based** feedbacks $\bar{u}_i : [0, T] \times \mathcal{N}_i \rightarrow \mathbb{R}^d$ defined by

$$\bar{u}_i(t, x) := \bar{u}_i\left(t, x_1^*(t), \dots, x_{i-1}^*(t), x, x_{i+1}^*(t), \dots, x_N^*(t)\right),$$

are **Lipschitz** on \mathcal{N}_i , i.e. $|\bar{u}_i(t, y) - \bar{u}_i(t, x)| \leq \mathcal{L}_U |x - y|$.

\hookrightarrow Collection of **feedback-neighbourhood** pairs $\{\bar{u}_i(\cdot, \cdot), \mathcal{N}_i\}_{i=1}^N$.

3. Build a **vector field** $u_N^* : [0, T] \times \mathbb{R}^d \rightarrow U$ by **patching together** the collection $\{\bar{u}_i(\cdot, \cdot), \mathcal{N}_i\}_{i=1}^N \oplus$ **Lip extension**.

\hookrightarrow Sequence $(u_N^*(\cdot, \cdot))$ of **loc. optimal feedbacks** with a **uniform Lipschitz regularity** in the space variable.

4. It holds $u_{N_k}^*(\cdot, \cdot) \rightharpoonup u^*(\cdot, \cdot)$ where $\text{Lip}(u^*(t, \cdot), \mathbb{R}^d) \leq \mathcal{L}_U$ and $u^*(\cdot, \cdot)$ is **optimal** for $(\text{MFOCP}) \rightsquigarrow \Gamma\text{-conv.}$ [FS'14, FLOS'19] \square

Lipschitz regularity in (OCP_N) – Proof of Theorem 2

2. The **agent-based** feedbacks $\bar{u}_i : [0, T] \times \mathcal{N}_i \rightarrow \mathbb{R}^d$ defined by

$$\bar{u}_i(t, x) := \bar{u}_i\left(t, x_1^*(t), \dots, x_{i-1}^*(t), x, x_{i+1}^*(t), \dots, x_N^*(t)\right),$$

are **Lipschitz** on \mathcal{N}_i , i.e. $|\bar{u}_i(t, y) - \bar{u}_i(t, x)| \leq \mathcal{L}_U |x - y|$.

\hookrightarrow Collection of **feedback-neighbourhood** pairs $\{\bar{u}_i(\cdot, \cdot), \mathcal{N}_i\}_{i=1}^N$.

3. Build a **vector field** $u_N^* : [0, T] \times \mathbb{R}^d \rightarrow U$ by **patching together** the collection $\{\bar{u}_i(\cdot, \cdot), \mathcal{N}_i\}_{i=1}^N \oplus$ **Lip extension**.

\hookrightarrow Sequence $(u_N^*(\cdot, \cdot))$ of **loc. optimal feedbacks** with a **uniform Lipschitz regularity** in the space variable.

4. It holds $u_{N_k}^*(\cdot, \cdot) \rightharpoonup u^*(\cdot, \cdot)$ where $\text{Lip}(u^*(t, \cdot), \mathbb{R}^d) \leq \mathcal{L}_U$ and $u^*(\cdot, \cdot)$ is **optimal** for **(MFOCP)** $\rightsquigarrow \Gamma$ -conv. [FS'14, FLOS'19] \square

Lipschitz regularity in (MFOCP) – Wrapping up everything

Question: Precise link with Theorem 1 dealing with (MFOCP) ?

Proposition (Sufficient condition for mean-field coercivity)

Suppose that the dynamics and cost functionals in (MFOCP) are $C_{\text{loc}}^{2,1}$ in the **measure** and **space variables**.

Then for all $N \geq 1$, the coercivity condition (MFCE_N) holds along **every** optimal Pontryagin triples for (OCP_N) **if and only if** $\lambda_\psi > \lambda_{\text{MF}}$ for some $\lambda_{\text{MF}} \geq 0$. In this case, one has $\rho = \lambda_\psi - \lambda_{\text{MF}} > 0$.

Remarks (Interpretation and link with MFGs)

$\lambda_{\text{MF}} \sim$ global **concavity constant** for (MFOCP) over $\mathcal{P}_2(\mathbb{R}^d)$

◇ If (MFOCP) is **sufficiently convex** \rightsquigarrow Lip sol. for any $\lambda_\psi > 0$

◇ Otherwise, one needs $\lambda_\psi > 0$ **large** w.r.t. data of (MFOCP)

\rightsquigarrow Well-posedness of master equation [GS'15,M'19,CCP'20,GM'21]

Lipschitz regularity in (MFOCP) – Wrapping up everything

Question: Precise link with Theorem 1 dealing with (MFOCP) ?

Proposition (Sufficient condition for mean-field coercivity)

Suppose that the dynamics and cost functionals in (MFOCP) are $C_{\text{loc}}^{2,1}$ in the **measure** and **space variables**.

Then for all $N \geq 1$, the coercivity condition (MFCE_N) holds along **every** optimal Pontryagin triples for (OCP_N) **if and only if** $\lambda_\psi > \lambda_{\text{MF}}$ for some $\lambda_{\text{MF}} \geq 0$. In this case, one has $\rho = \lambda_\psi - \lambda_{\text{MF}} > 0$.

Remarks (Interpretation and link with MFGs)

$\lambda_{\text{MF}} \sim$ global **concavity constant** for (MFOCP) over $\mathcal{P}_2(\mathbb{R}^d)$

◇ If (MFOCP) is **sufficiently convex** \rightsquigarrow Lip sol. for any $\lambda_\psi > 0$

◇ Otherwise, one needs $\lambda_\psi > 0$ **large** w.r.t. data of (MFOCP)

\rightsquigarrow Well-posedness of master equation [GS'15, M'19, CCP'20, GM'21]

Lipschitz regularity in (MFOCP) – Wrapping up everything

Question: Precise link with Theorem 1 dealing with (MFOCP) ?

Proposition (Sufficient condition for mean-field coercivity)

Suppose that the dynamics and cost functionals in (MFOCP) are $C_{\text{loc}}^{2,1}$ in the **measure** and **space variables**.

Then for all $N \geq 1$, the coercivity condition (MFCE_N) holds along **every** optimal Pontryagin triples for (OCP_N) **if and only if** $\lambda_\psi > \lambda_{\text{MF}}$ for some $\lambda_{\text{MF}} \geq 0$. In this case, one has $\rho = \lambda_\psi - \lambda_{\text{MF}} > 0$.

Remarks (Interpretation and link with MFGs)

$\lambda_{\text{MF}} \sim$ global **concavity constant** for (MFOCP) over $\mathcal{P}_2(\mathbb{R}^d)$

◇ If (MFOCP) is **sufficiently convex** \rightsquigarrow Lip sol. for any $\lambda_\psi > 0$

◇ Otherwise, one needs $\lambda_\psi > 0$ **large** w.r.t. data of (MFOCP)

\rightsquigarrow Well-posedness of master equation [GS'15, M'19, CCP'20, GM'21]

Lipschitz regularity in (MFOCP) – Wrapping up everything

Question: Precise link with Theorem 1 dealing with (MFOCP) ?

Proposition (Sufficient condition for mean-field coercivity)

Suppose that the dynamics and cost functionals in (MFOCP) are $C_{\text{loc}}^{2,1}$ in the **measure** and **space variables**.

Then for all $N \geq 1$, the coercivity condition (MFCE_N) holds along **every** optimal Pontryagin triples for (OCP_N) **if and only if** $\lambda_\psi > \lambda_{\text{MF}}$ for some $\lambda_{\text{MF}} \geq 0$. In this case, one has $\rho = \lambda_\psi - \lambda_{\text{MF}} > 0$.

Remarks (Interpretation and link with MFGs)

$\lambda_{\text{MF}} \sim$ global **concavity constant** for (MFOCP) over $\mathcal{P}_2(\mathbb{R}^d)$

◇ If (MFOCP) is **sufficiently convex** \rightsquigarrow Lip sol. for any $\lambda_\psi > 0$

◇ Otherwise, one needs $\lambda_\psi > 0$ **large** w.r.t. data of (MFOCP)

\rightsquigarrow Well-posedness of master equation [GS'15, M'19, CCP'20, GM'21]

Lipschitz regularity in (MFOCP) – Wrapping up everything

Question: Precise link with Theorem 1 dealing with (MFOCP) ?

Proposition (Sufficient condition for mean-field coercivity)

Suppose that the dynamics and cost functionals in (MFOCP) are $C_{\text{loc}}^{2,1}$ in the **measure** and **space variables**.

Then for all $N \geq 1$, the coercivity condition (MFCE_N) holds along **every** optimal Pontryagin triples for (OCP_N) **if and only if** $\lambda_\psi > \lambda_{\text{MF}}$ for some $\lambda_{\text{MF}} \geq 0$. In this case, one has $\rho = \lambda_\psi - \lambda_{\text{MF}} > 0$.

Remarks (Interpretation and link with MFGs)

$\lambda_{\text{MF}} \sim$ global **concavity constant** for (MFOCP) over $\mathcal{P}_2(\mathbb{R}^d)$

◇ If (MFOCP) is **sufficiently convex** \rightsquigarrow Lip sol. for any $\lambda_\psi > 0$

◇ Otherwise, one needs $\lambda_\psi > 0$ **large** w.r.t. data of (MFOCP)

\rightsquigarrow Well-posedness of master equation [GS'15, M'19, CCP'20, GM'21]

Lipschitz regularity in (MFOCP) – Wrapping up everything

Question: Precise link with Theorem 1 dealing with (MFOCP) ?

Proposition (Sufficient condition for mean-field coercivity)

Suppose that the dynamics and cost functionals in (MFOCP) are $C_{loc}^{2,1}$ in the **measure** and **space variables**.

Then for all $N \geq 1$, the coercivity condition (MFCE_N) holds along **every** optimal Pontryagin triples for (OCP_N) **if and only if** $\lambda_\psi > \lambda_{MF}$ for some $\lambda_{MF} \geq 0$. In this case, one has $\rho = \lambda_\psi - \lambda_{MF} > 0$.

Remarks (Interpretation and link with MFGs)

$\lambda_{MF} \sim$ global **concavity constant** for (MFOCP) over $\mathcal{P}_2(\mathbb{R}^d)$

◇ If (MFOCP) is **sufficiently convex** \rightsquigarrow Lip sol. for any $\lambda_\psi > 0$

◇ Otherwise, one needs $\lambda_\psi > 0$ **large** w.r.t. data of (MFOCP)

\rightsquigarrow Well-posedness of master equation [GS'15, M'19, CCP'20, GM'21]

Lipschitz regularity in (MFOCP) – Wrapping up everything

Question: Precise link with Theorem 1 dealing with (MFOCP) ?

Proposition (Sufficient condition for mean-field coercivity)

Suppose that the dynamics and cost functionals in (MFOCP) are $C_{\text{loc}}^{2,1}$ in the **measure** and **space variables**.

Then for all $N \geq 1$, the coercivity condition (MFCE_N) holds along **every** optimal Pontryagin triples for (OCP_N) **if and only if** $\lambda_\psi > \lambda_{\text{MF}}$ for some $\lambda_{\text{MF}} \geq 0$. In this case, one has $\rho = \lambda_\psi - \lambda_{\text{MF}} > 0$.

Remarks (Interpretation and link with MFGs)

$\lambda_{\text{MF}} \sim$ global **concavity constant** for (MFOCP) over $\mathcal{P}_2(\mathbb{R}^d)$

◇ If (MFOCP) is **sufficiently convex** \rightsquigarrow Lip sol. for any $\lambda_\psi > 0$

◇ Otherwise, one needs $\lambda_\psi > 0$ **large** w.r.t. data of (MFOCP)

\rightsquigarrow Well-posedness of master equation [GS'15, M'19, CCP'20, GM'21]

Lipschitz regularity in (MFOCP) – Wrapping up everything

Question: Precise link with Theorem 1 dealing with (MFOCP) ?

Proposition (Sufficient condition for mean-field coercivity)

Suppose that the dynamics and cost functionals in (MFOCP) are $C_{\text{loc}}^{2,1}$ in the **measure** and **space variables**.

Then for all $N \geq 1$, the coercivity condition (MFCE_N) holds along **every** optimal Pontryagin triples for (OCP_N) **if and only if** $\lambda_\psi > \lambda_{\text{MF}}$ for some $\lambda_{\text{MF}} \geq 0$. In this case, one has $\rho = \lambda_\psi - \lambda_{\text{MF}} > 0$.

Remarks (Interpretation and link with MFGs)

$\lambda_{\text{MF}} \sim$ global **concavity constant** for (MFOCP) over $\mathcal{P}_2(\mathbb{R}^d)$

- ◇ If (MFOCP) is **sufficiently convex** \rightsquigarrow Lip sol. for any $\lambda_\psi > 0$
- ◇ Otherwise, one needs $\lambda_\psi > 0$ **large** w.r.t. data of (MFOCP)

\hookrightarrow Well-posedness of **master equation** [GS'15, M'19, CCP'20, GM'21]

Lipschitz regularity in (MFOCP) – Wrapping up everything

Question: Precise link with Theorem 1 dealing with (MFOCP) ?

Proposition (Sufficient condition for mean-field coercivity)

Suppose that the dynamics and cost functionals in (MFOCP) are $C_{\text{loc}}^{2,1}$ in the **measure** and **space variables**.

Then for all $N \geq 1$, the coercivity condition (MFCE_N) holds along **every** optimal Pontryagin triples for (OCP_N) **if and only if** $\lambda_\psi > \lambda_{\text{MF}}$ for some $\lambda_{\text{MF}} \geq 0$. In this case, one has $\rho = \lambda_\psi - \lambda_{\text{MF}} > 0$.

Remarks (Interpretation and link with MFGs)

$\lambda_{\text{MF}} \sim$ global **concavity constant** for (MFOCP) over $\mathcal{P}_2(\mathbb{R}^d)$

- ◇ If (MFOCP) is **sufficiently convex** \rightsquigarrow Lip sol. for any $\lambda_\psi > 0$
- ◇ Otherwise, one needs $\lambda_\psi > 0$ **large** w.r.t. data of (MFOCP)

\hookrightarrow Well-posedness of **master equation** [GS'15,M'19,CCP'20,GM'21]

Outline of the talk

Multi-agent systems and mean-field optimal control

Lipschitz feedbacks in classical OCPs

Lipschitz feedbacks in mean-field OCPs

Example in 1D – Maximisation of the variance

Conclusion and open problems

Variance maximisation – Formulation of the problem

Consider the following **(MFOCP)** and the corresponding **(OCP_N)**

Infinite dimensional problem **(MFOCP)**

$$\begin{cases} \min_{u(\cdot, \cdot)} \left[\frac{\lambda}{2} \int_0^T \int_{\mathbb{R}} |u(t, x)|^2 d\mu(t)(x) dt - \frac{1}{2} \int_{\mathbb{R}} |x - \bar{\mu}(T)|^2 d\mu(T) \right] \\ \text{s.t.} \begin{cases} \partial_t \mu(t) + \operatorname{div}_x(u(t, \cdot) \mu(t)) = 0, & \|u^*(t, \cdot)\|_{L^\infty(\mu(t))} \leq 1, \\ \mu(0) = \frac{1}{2} \mathbb{1}_{[-1,1]} \cdot \mathcal{L}^1. \end{cases} \end{cases}$$

Approximating sequence of finite dimensional problems **(OCP_N)**

$$\begin{cases} \min_{u(\cdot)} \left[\frac{\lambda}{2N} \sum_{i=1}^N \int_0^T |u_i(t)|^2 dt - \frac{1}{2N} \sum_{i=1}^N |x_i(T) - \bar{x}(T)|^2 \right] \\ \text{s.t.} \begin{cases} \dot{x}_i(t) = u_i(t), & |u_i(t)| \leq 1 \\ x_i(0) = x_i^0. \end{cases} \end{cases}$$

Variance maximisation – *The role of coercivity*

Idea: there are roughly two ways of increasing the variance

1. **Spreading** the initial indicator function (**smooth** action)
2. **Breaking** the initial indicator function (**non-smooth** action)

↔ the **coercivity estimate determines** each scenario!

Proposition (Variance maximisation)[B. & Rossi'21]

The following statements are equivalent

- (i) (MFCE_N) holds with constant $\rho > 0$ along every triple.
- (ii) It holds $T < \lambda$ and $\rho = \lambda - T > 0$, i.e. $\lambda_{\text{MF}} = T$.
- (iii) The finite-dimensional controls $\{u_i^*(\cdot)\}_{i=1}^N$ satisfy

$$|u_i^*(t) - u_j^*(t)| \leq \frac{1}{\rho} |x_i^*(t) - x_j^*(t)|.$$

Variance maximisation – *The role of coercivity*

Idea: there are roughly two ways of increasing the variance

1. **Spreading** the initial indicator function (**smooth** action)
2. **Breaking** the initial indicator function (**non-smooth** action)

↔ the **coercivity estimate determines** each scenario!

Proposition (Variance maximisation)[B. & Rossi'21]

The following statements are equivalent

- (i) (MFCE_N) holds with constant $\rho > 0$ along every triple.
- (ii) It holds $T < \lambda$ and $\rho = \lambda - T > 0$, i.e. $\lambda_{\text{MF}} = T$.
- (iii) The finite-dimensional controls $\{u_i^*(\cdot)\}_{i=1}^N$ satisfy

$$|u_i^*(t) - u_j^*(t)| \leq \frac{1}{\rho} |x_i^*(t) - x_j^*(t)|.$$

Variance maximisation – *The role of coercivity*

Idea: there are roughly two ways of increasing the variance

1. **Spreading** the initial indicator function (**smooth** action)
2. **Breaking** the initial indicator function (**non-smooth** action)

↔ the **coercivity estimate determines** each scenario!

Proposition (Variance maximisation)[B. & Rossi'21]

The following statements are equivalent

- (i) (MFCE_N) holds with constant $\rho > 0$ along every triple.
- (ii) It holds $T < \lambda$ and $\rho = \lambda - T > 0$, i.e. $\lambda_{\text{MF}} = T$.
- (iii) The finite-dimensional controls $\{u_i^*(\cdot)\}_{i=1}^N$ satisfy

$$|u_i^*(t) - u_j^*(t)| \leq \frac{1}{\rho} |x_i^*(t) - x_j^*(t)|.$$

Variance maximisation – *The role of coercivity*

Idea: there are roughly two ways of increasing the variance

1. **Spreading** the initial indicator function (**smooth** action)
2. **Breaking** the initial indicator function (**non-smooth** action)

↪ the **coercivity estimate determines** each scenario!

Proposition (Variance maximisation)[B. & Rossi'21]

The following statements are equivalent

- (i) (MFCE_N) holds with constant $\rho > 0$ along every triple.
- (ii) It holds $T < \lambda$ and $\rho = \lambda - T > 0$, i.e. $\lambda_{\text{MF}} = T$.
- (iii) The finite-dimensional controls $\{u_i^*(\cdot)\}_{i=1}^N$ satisfy

$$|u_i^*(t) - u_j^*(t)| \leq \frac{1}{\rho} |x_i^*(t) - x_j^*(t)|.$$

Variance maximisation – *The role of coercivity*

Idea: there are roughly two ways of increasing the variance

1. **Spreading** the initial indicator function (**smooth** action)
2. **Breaking** the initial indicator function (**non-smooth** action)

↪ the **coercivity estimate determines** each scenario!

Proposition (Variance maximisation)[B. & Rossi'21]

The following statements are equivalent

- (i) (MFCE_N) holds with constant $\rho > 0$ along every triple.
- (ii) It holds $T < \lambda$ and $\rho = \lambda - T > 0$, i.e. $\lambda_{\text{MF}} = T$.
- (iii) The finite-dimensional controls $\{u_i^*(\cdot)\}_{i=1}^N$ satisfy

$$|u_i^*(t) - u_j^*(t)| \leq \frac{1}{\rho} |x_i^*(t) - x_j^*(t)|.$$

Variance maximisation – *The role of coercivity*

Idea: there are roughly two ways of increasing the variance

1. **Spreading** the initial indicator function (**smooth** action)
2. **Breaking** the initial indicator function (**non-smooth** action)

↪ the **coercivity estimate determines** each scenario!

Proposition (Variance maximisation)[B. & Rossi'21]

The following statements are equivalent

- (i) (MFCE_N) holds with constant $\rho > 0$ along every triple.
- (ii) It holds $T < \lambda$ and $\rho = \lambda - T > 0$, i.e. $\lambda_{\text{MF}} = T$.
- (iii) The finite-dimensional controls $\{u_i^*(\cdot)\}_{i=1}^N$ satisfy

$$|u_i^*(t) - u_j^*(t)| \leq \frac{1}{\rho} |x_i^*(t) - x_j^*(t)|.$$

Variance maximisation – *The role of coercivity*

Idea: there are roughly two ways of increasing the variance

1. **Spreading** the initial indicator function (**smooth** action)
2. **Breaking** the initial indicator function (**non-smooth** action)

↪ the **coercivity estimate determines** each scenario!

Proposition (Variance maximisation)[B. & Rossi'21]

The following statements are equivalent

- (i) (MFCE_N) holds with constant $\rho > 0$ along every triple.
- (ii) It holds $T < \lambda$ and $\rho = \lambda - T > 0$, i.e. $\lambda_{\text{MF}} = T$.
- (iii) The finite-dimensional controls $\{u_i^*(\cdot)\}_{i=1}^N$ satisfy

$$|u_i^*(t) - u_j^*(t)| \leq \frac{1}{\rho} |x_i^*(t) - x_j^*(t)|.$$

Variance maximisation – *The role of coercivity*

Idea: there are roughly two ways of increasing the variance

1. **Spreading** the initial indicator function (**smooth** action)
2. **Breaking** the initial indicator function (**non-smooth** action)

↪ the **coercivity estimate determines** each scenario!

Proposition (Variance maximisation)[B. & Rossi'21]

The following statements are equivalent

- (i) (MFCE_N) holds with constant $\rho > 0$ along every triple.
- (ii) It holds $T < \lambda$ and $\rho = \lambda - T > 0$, i.e. $\lambda_{\text{MF}} = T$.
- (iii) The finite-dimensional controls $\{u_i^*(\cdot)\}_{i=1}^N$ satisfy

$$|u_i^*(t) - u_j^*(t)| \leq \frac{1}{\rho} |x_i^*(t) - x_j^*(t)|.$$

Outline of the talk

Multi-agent systems and mean-field optimal control

Lipschitz feedbacks in classical OCPs

Lipschitz feedbacks in mean-field OCPs

Example in 1D – Maximisation of the variance

Conclusion and open problems

Lipschitz regularity in (MFOCP) – *Some open perspectives*

We have a “*sharp-ish*” condition for **Lipschitz regularity**... Hence Cauchy-Lipschitz regularity is **very demanding** for (MFOCP)

Open problem (more general well-posedness analysis)

Could we choose another framework for the well-posedness theory ?

↪ **Regular Lagrangian Flows** [DpL'89,A'04]

Idea: Considering rough vector fields on **absolutely continuous** densities & select solutions that **do not concentrate**.

Difficulty: Finding a good definition for the convergence of **particle systems** towards **Lagrangian flows** (still open a priori)

Lipschitz regularity in (MFOCP) – *Some open perspectives*

We have a “*sharp-ish*” condition for **Lipschitz regularity**... Hence Cauchy-Lipschitz regularity is **very demanding** for (MFOCP)

Open problem (more general well-posedness analysis)

Could we choose another framework for the well-posedness theory ?

↪ **Regular Lagrangian Flows** [DpL'89,A'04]

Idea: Considering rough vector fields on **absolutely continuous** densities & select solutions that **do not concentrate**.

Difficulty: Finding a good definition for the convergence of **particle systems** towards **Lagrangian flows** (still open a priori)

Lipschitz regularity in (MFOCP) – *Some open perspectives*

We have a “*sharp-ish*” condition for **Lipschitz regularity**... Hence Cauchy-Lipschitz regularity is **very demanding** for (MFOCP)

Open problem (more general well-posedness analysis)

Could we choose another framework for the well-posedness theory ?

↪ **Regular Lagrangian Flows** [DpL'89,A'04]

Idea: Considering rough vector fields on **absolutely continuous** densities & select solutions that **do not concentrate**.

Difficulty: Finding a good definition for the convergence of **particle systems** towards **Lagrangian flows** (still open a priori)

Lipschitz regularity in (MFOCP) – *Some open perspectives*

We have a “*sharp-ish*” condition for **Lipschitz regularity**... Hence Cauchy-Lipschitz regularity is **very demanding** for (MFOCP)

Open problem (more general well-posedness analysis)

Could we choose another framework for the well-posedness theory ?

↪ **Regular Lagrangian Flows** [DpL'89,A'04]

Idea: Considering rough vector fields on **absolutely continuous** densities & select solutions that **do not concentrate**.

Difficulty: Finding a good definition for the convergence of **particle systems** towards **Lagrangian flows** (still open a priori)

Lipschitz regularity in (MFOCP) – *Some open perspectives*

We have a “*sharp-ish*” condition for **Lipschitz regularity**... Hence Cauchy-Lipschitz regularity is **very demanding** for (MFOCP)

Open problem (more general well-posedness analysis)

Could we choose another framework for the well-posedness theory ?

↪ **Regular Lagrangian Flows** [DpL'89,A'04]

Idea: Considering rough vector fields on **absolutely continuous** densities & select solutions that **do not concentrate**.

Difficulty: Finding a good definition for the convergence of **particle systems** towards **Lagrangian flows** (still open a priori)

Lipschitz regularity in (MFOCP) – *Some open perspectives*

We have a “*sharp-ish*” condition for **Lipschitz regularity**... Hence Cauchy-Lipschitz regularity is **very demanding** for (MFOCP)

Open problem (more general well-posedness analysis)

Could we choose another framework for the well-posedness theory ?

↪ **Regular Lagrangian Flows** [DpL'89,A'04]

Idea: Considering rough vector fields on **absolutely continuous** densities & select solutions that **do not concentrate**.

Difficulty: Finding a good definition for the convergence of **particle systems** towards **Lagrangian flows** (still open a priori)

Lipschitz regularity in (MFOCP) – *Some open perspectives*

We have a “*sharp-ish*” condition for **Lipschitz regularity**... Hence Cauchy-Lipschitz regularity is **very demanding** for (MFOCP)

Open problem (more general well-posedness analysis)

Could we choose another framework for the well-posedness theory ?

↪ **Regular Lagrangian Flows** [DpL'89,A'04]

Idea: Considering rough vector fields on **absolutely continuous** densities & select solutions that **do not concentrate**.

Difficulty: Finding a good definition for the convergence of **particle systems** towards **Lagrangian flows** (still open a priori)

All in all...

Thank you for your attention !

