



Continuity Inclusions and Applications in Mean-Field Optimal Control

Benoît Bonnet

(in collaboration with H. Frankowska)

**Seminar of Analysis, Stochastic Phenomena and
Applications**

February 2, 2021

Outline of the talk

Mean-field control problems

From differential to continuity inclusions

Main structural results in the Cauchy-Lipschitz case

Application : geometric derivation of the PMP

Conclusion

Outline of the talk

Mean-field control problems

From differential to continuity inclusions

Main structural results in the Cauchy-Lipschitz case

Application : geometric derivation of the PMP

Conclusion

Introduction – *Mean-field control*

⇒ Control of **multi-agent** systems in the **mean-field** limit

Introduction – *Mean-field control*

↪ Control of **multi-agent** systems in the **mean-field** limit




Introduction – Mean-field limits of particle systems

First observation (dimensionality-related problems)

The number N of agents is usually **very large** \rightsquigarrow numerical issues

Idea: Approximation procedure using **mean-field** limits!



System of ODEs on N agents $(x_1, \dots, x_N) \in (\mathbb{R}^d)^N$

$$\left(\mu_N = \frac{1}{N} \sum_{i=1}^N \delta_{x_i} \right)$$


Single PDE on the density of agents $\mu : \mathbb{R}^d \rightarrow \mathbb{R}$.

Introduction – Mean-field limits of particle systems

First observation (dimensionality-related problems)

The number N of agents is usually **very large** \rightsquigarrow **numerical issues**

Idea: Approximation procedure using **mean-field** limits!



System of ODEs on N agents $(x_1, \dots, x_N) \in (\mathbb{R}^d)^N$

$$\left(\mu_N = \frac{1}{N} \sum_{i=1}^N \delta_{x_i} \right)$$


Single PDE on the density of agents $\mu : \mathbb{R}^d \rightarrow \mathbb{R}$.

Introduction – Mean-field limits of particle systems

First observation (dimensionality-related problems)

The number N of agents is usually **very large** \rightsquigarrow **numerical issues**

Idea: Approximation procedure using **mean-field** limits!



System of ODEs on N agents $(x_1, \dots, x_N) \in (\mathbb{R}^d)^N$

$$\left(\mu_N = \frac{1}{N} \sum_{i=1}^N \delta_{x_i} \right)$$


Single PDE on the density of agents $\mu : \mathbb{R}^d \rightarrow \mathbb{R}$.

Introduction – Mean-field limits of particle systems

First observation (dimensionality-related problems)

The number N of agents is usually **very large** \rightsquigarrow **numerical issues**

Idea: Approximation procedure using **mean-field** limits!



System of ODEs on N agents $(x_1, \dots, x_N) \in (\mathbb{R}^d)^N$

$$\left(\mu_N = \frac{1}{N} \sum_{i=1}^N \delta_{x_i} \right)$$

Single PDE on the density of agents $\mu : \mathbb{R}^d \rightarrow \mathbb{R}$.

Introduction – *Continuity equations*

Deterministic mean-field dynamics \rightsquigarrow **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), \cdot) \mu(t)) = 0,$$

where $\mu(t) \in \mathcal{P}_c(\mathbb{R}^d)$ and $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times \mathbb{R}^d \rightarrow \mathbb{R}^d$.




Figure: *Solution of a continuity equation with smooth driving field*

Introduction – *Continuity equations*

Deterministic mean-field dynamics \rightsquigarrow **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), \cdot) \mu(t)) = 0,$$

where $\mu(t) \in \mathcal{P}_c(\mathbb{R}^d)$ and $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times \mathbb{R}^d \rightarrow \mathbb{R}^d$.




Figure: *Solution of a continuity equation with smooth driving field*

Introduction – *Continuity equations*

Deterministic mean-field dynamics \rightsquigarrow **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), \cdot) \mu(t)) = 0,$$

where $\mu(t) \in \mathcal{P}_c(\mathbb{R}^d)$ and $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times \mathbb{R}^d \rightarrow \mathbb{R}^d$.




Figure: *Solution of a continuity equation with smooth driving field*

Introduction – Mean-field optimal control

Mean-field control \rightsquigarrow control problems on **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), u(t), \cdot) \mu(t)) = 0,$$

where $u : [0, T] \rightarrow U$ is a **control law**.

- **Existence of optimal trajectories** \rightsquigarrow (B., Frankowska, Fornasier, Marigonda, Quincampoix, Pogodaev, Rossi, Savaré, Solombrino, etc...)
- **1st-order optimality conditions** \rightsquigarrow (B., Carmona, Frankowska, Jannin, Marigonda, Quincampoix, Pataki, Pogodaev, Rossi, etc...)
- **Properties of value functions** \rightsquigarrow (B., Frankowska, Santambrogio, Carmona, Gangbo, Jannin, Marigonda, Quincampoix, etc...)
- **Optimal feedback synthesis** \rightsquigarrow Somewhat open for now

Natural question : General framework to study these problems ?
 \rightsquigarrow Yes! Reformulate controlled dynamics as differential inclusions

Introduction – Mean-field optimal control

Mean-field control \rightsquigarrow control problems on **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), u(t), \cdot) \mu(t)) = 0,$$

where $u : [0, T] \rightarrow U$ is a **control law**.

- ◊ **Existence of optimal trajectories** \rightsquigarrow (B., Frankowska, Fornasier, Marigonda, Quincampoix, Pogodaev, Rossi, Savaré, Solombrino, etc...)
- ◊ **1st-order optimality conditions** \rightsquigarrow (B., Cavagnari, Frankowska, Jimenez, Marigonda, Quincampoix, Piccoli, Pogodaev, Rossi, etc...)
- ◊ **Properties of value functions** \rightsquigarrow (B., Frankowska, Lantieri, Cavagnari, Jimenez, Marigonda, Quincampoix, etc...)
- ◊ **Optimal feedback synthesis** \rightsquigarrow Somewhat open for now

Natural question : General framework to study these problems ?
 \rightsquigarrow Yes! Reformulate controlled dynamics as differential inclusions

Introduction – Mean-field optimal control

Mean-field control \rightsquigarrow control problems on **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), u(t), \cdot) \mu(t)) = 0,$$

where $u : [0, T] \rightarrow U$ is a **control law**.

- ◇ **Existence of optimal trajectories** \rightsquigarrow (B., Frankowska, Fornasier, Marigonda, Quincampoix, Pogodaev, Rossi, Savaré, Solombrino, etc...)
- ◇ **1st-order optimality conditions** \rightsquigarrow (B., Cavagnari, Frankowska, Jimenez, Marigonda, Quincampoix, Piccoli, Pogodaev, Rossi, etc...)
- ◇ **Properties of value functions** \rightsquigarrow (B., Frankowska, Cardaliaguet, Cavagnari, Gangbo, Jimenez, Marigonda, Quincampoix, etc...)
- ◇ **Optimal feedback synthesis** \rightsquigarrow Somewhat open for now

Natural question : General framework to study these problems ?
 \rightsquigarrow Yes! Reformulate controlled dynamics as differential inclusions

Introduction – Mean-field optimal control

Mean-field control \rightsquigarrow control problems on **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), u(t), \cdot) \mu(t)) = 0,$$

where $u : [0, T] \rightarrow U$ is a **control law**.

- ◇ **Existence of optimal trajectories** \rightsquigarrow (B., Frankowska, Fornasier, Marigonda, Quincampoix, Pogodaev, Rossi, Savaré, Solombrino, etc...)
- ◇ **1st-order optimality conditions** \rightsquigarrow (B., Cavagnari, Frankowska, Jimenez, Marigonda, Quincampoix, Piccoli, Pogodaev, Rossi, etc...)
- ◇ **Properties of value functions** \rightsquigarrow (B., Frankowska, Cardaliaguet, Cavagnari, Gangbo, Jimenez, Marigonda, Quincampoix, etc...)
- ◇ **Optimal feedback synthesis** \rightsquigarrow Somewhat open for now

Natural question : General framework to study these problems ?

\rightsquigarrow Yes! Reformulate controlled dynamics as **differential inclusions**

Introduction – *Mean-field optimal control*

Mean-field control \rightsquigarrow control problems on **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), u(t), \cdot) \mu(t)) = 0,$$

where $u : [0, T] \rightarrow U$ is a **control law**.

- ◇ **Existence of optimal trajectories** \rightsquigarrow (B., Frankowska, Fornasier, Marigonda, Quincampoix, Pogodaev, Rossi, Savaré, Solombrino, etc...)
- ◇ **1st-order optimality conditions** \rightsquigarrow (B., Cavagnari, Frankowska, Jimenez, Marigonda, Quincampoix, Piccoli, Pogodaev, Rossi, etc...)
- ◇ **Properties of value functions** \rightsquigarrow (B., Frankowska, Cardaliaguet, Cavagnari, Gangbo, Jimenez, Marigonda, Quincampoix, etc...)
- ◇ **Optimal feedback synthesis** \rightsquigarrow Somewhat open for now

Natural question : General framework to study these problems ?

\rightsquigarrow Yes! Reformulate controlled dynamics as **differential inclusions**

Introduction – *Mean-field optimal control*

Mean-field control \rightsquigarrow control problems on **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), u(t), \cdot) \mu(t)) = 0,$$

where $u : [0, T] \rightarrow U$ is a **control law**.

- ◇ **Existence of optimal trajectories** \rightsquigarrow (B., Frankowska, Fornasier, Marigonda, Quincampoix, Pogodaev, Rossi, Savaré, Solombrino, etc...)
- ◇ **1st-order optimality conditions** \rightsquigarrow (B., Cavagnari, Frankowska, Jimenez, Marigonda, Quincampoix, Piccoli, Pogodaev, Rossi, etc...)
- ◇ **Properties of value functions** \rightsquigarrow (B., Frankowska, Cardaliaguet, Cavagnari, Gangbo, Jimenez, Marigonda, Quincampoix, etc...)
- ◇ **Optimal feedback synthesis** \rightsquigarrow Somewhat open for now

Natural question : General framework to study these problems ?
 \rightsquigarrow Yes! Reformulate controlled dynamics as **differential inclusions**

Introduction – *Mean-field optimal control*

Mean-field control \rightsquigarrow control problems on **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), u(t), \cdot) \mu(t)) = 0,$$

where $u : [0, T] \rightarrow U$ is a **control law**.

- ◇ **Existence of optimal trajectories** \rightsquigarrow (B., Frankowska, Fornasier, Marigonda, Quincampoix, Pogodaev, Rossi, Savaré, Solombrino, etc...)
- ◇ **1st-order optimality conditions** \rightsquigarrow (B., Cavagnari, Frankowska, Jimenez, Marigonda, Quincampoix, Piccoli, Pogodaev, Rossi, etc...)
- ◇ **Properties of value functions** \rightsquigarrow (B., Frankowska, Cardaliaguet, Cavagnari, Gangbo, Jimenez, Marigonda, Quincampoix, etc...)
- ◇ **Optimal feedback synthesis** \rightsquigarrow Somewhat open for now

Natural question : General framework to study these problems ?

\hookrightarrow Yes! Reformulate controlled dynamics as **differential inclusions**

Introduction – *Mean-field optimal control*

Mean-field control \rightsquigarrow control problems on **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), u(t), \cdot) \mu(t)) = 0,$$

where $u : [0, T] \rightarrow U$ is a **control law**.

- ◇ **Existence of optimal trajectories** \rightsquigarrow (B., Frankowska, Fornasier, Marigonda, Quincampoix, Pogodaev, Rossi, Savaré, Solombrino, etc...)
- ◇ **1st-order optimality conditions** \rightsquigarrow (B., Cavagnari, Frankowska, Jimenez, Marigonda, Quincampoix, Piccoli, Pogodaev, Rossi, etc...)
- ◇ **Properties of value functions** \rightsquigarrow (B., Frankowska, Cardaliaguet, Cavagnari, Gangbo, Jimenez, Marigonda, Quincampoix, etc...)
- ◇ **Optimal feedback synthesis** \rightsquigarrow Somewhat open for now

Natural question : General framework to study these problems ?

\hookrightarrow Yes! Reformulate controlled dynamics as **differential inclusions**

Outline of the talk

Mean-field control problems

From differential to continuity inclusions

Main structural results in the Cauchy-Lipschitz case

Application : geometric derivation of the PMP

Conclusion

Differential inclusions – *The classical case*

Intuition: Differential inclusions \Longleftrightarrow ODEs with **set-valued** r.h.s

Definition (differential inclusion)

Let \mathcal{M} be a smooth manifold and $F : \mathcal{M} \rightrightarrows T\mathcal{M}$. Then

$$\dot{x}(t) \in F(x(t)) \iff \begin{cases} x(0) = x_0, \\ x(\cdot) \in C^1([0, T]), \\ \dot{x}(t) \in F(x(t)). \end{cases}$$



Differential inclusions – *The classical case*

Intuition: Differential inclusions \iff ODEs with **set-valued** r.h.s

Definition (differential inclusion)

Let \mathcal{M} be a smooth manifold and $F : \mathcal{M} \rightrightarrows T\mathcal{M}$. Then

$$\dot{x}(t) \in F(x(t)) \iff \begin{cases} \dot{x}(t) = f(t), \\ f(t) \in F(x(t)), \end{cases} \quad \mathcal{L}^1\text{-a.e. in } [0, T]$$




Differential inclusions – *The classical case*

Intuition: Differential inclusions \Longleftrightarrow ODEs with **set-valued** r.h.s

Definition (differential inclusion)

Let \mathcal{M} be a smooth manifold and $F : \mathcal{M} \rightrightarrows T\mathcal{M}$. Then

$$\dot{x}(t) \in F(x(t)) \Longleftrightarrow \begin{cases} \dot{x}(t) = f(t), \\ f(t) \in F(x(t)), \end{cases} \quad \mathcal{L}^1\text{-a.e. in } [0, T]$$




Differential inclusions – *The classical case*

Intuition: Differential inclusions \Longleftrightarrow ODEs with **set-valued** r.h.s

Definition (differential inclusion)

Let \mathcal{M} be a smooth manifold and $F : \mathcal{M} \rightrightarrows T\mathcal{M}$. Then

$$\dot{x}(t) \in F(x(t)) \Longleftrightarrow \begin{cases} \dot{x}(t) = f(t), \\ f(t) \in F(x(t)), \end{cases} \quad \mathcal{L}^1\text{-a.e. in } [0, T]$$



Differential inclusions – *Link with control system*

Key point: Control systems are **equivalent** to diff. inclusions

Equivalence (heuristic statement)

If $f : [0, T] \times \mathcal{M} \times U \rightarrow T\mathcal{M}$ is **Carathéodory**, then

$$\dot{x}(t) = f(t, x(t), u(t)) \iff \dot{x}(t) \in F(t, x(t)),$$

where

$$F(t, x) := \left\{ f(t, x, u) \text{ s.t. } u \in U \right\}.$$

Recap': Building differential inclusions

- 1) Check that the ambient space has a **manifold-like** structure,
- 2) Identify an ODE structure,
- 3) Make the velocities **set-valued** !

Differential inclusions – *Link with control system*

Key point: Control systems are **equivalent** to diff. inclusions

Equivalence (heuristic statement)

If $f : [0, T] \times \mathcal{M} \times U \rightarrow T\mathcal{M}$ is **Carathéodory**, then

$$\dot{x}(t) = f(t, x(t), u(t)) \iff \dot{x}(t) \in F(t, x(t)),$$

where

$$F(t, x) := \left\{ f(t, x, u) \text{ s.t. } u \in U \right\}.$$

Recap': Building differential inclusions

- 1) Check that the ambient space has a manifold-like structure,
- 2) Identify an ODE structure,
- 3) Make the velocities set-valued !

Differential inclusions – *Link with control system*

Key point: Control systems are **equivalent** to diff. inclusions

Equivalence (heuristic statement)

If $f : [0, T] \times \mathcal{M} \times U \rightarrow T\mathcal{M}$ is **Carathéodory**, then

$$\dot{x}(t) = f(t, x(t), u(t)) \iff \dot{x}(t) \in F(t, x(t)),$$

where

$$F(t, x) := \left\{ f(t, x, u) \text{ s.t. } u \in U \right\}.$$

Recap': Building differential inclusions

- 1) Check that the ambient space has a manifold-like structure,
- 2) Identify an ODE structure,
- 3) Make the velocities **set-valued** !

Differential inclusions – *Link with control system*

Key point: Control systems are **equivalent** to diff. inclusions

Equivalence (heuristic statement)

If $f : [0, T] \times \mathcal{M} \times U \rightarrow T\mathcal{M}$ is **Carathéodory**, then

$$\dot{x}(t) = f(t, x(t), u(t)) \iff \dot{x}(t) \in F(t, x(t)),$$

where

$$F(t, x) := \left\{ f(t, x, u) \text{ s.t. } u \in U \right\}.$$

Recap': Building differential inclusions

- 1) Check that the ambient space has a **manifold-like** structure,
- 2) Identify an ODE structure,
- 3) Make the velocities **set-valued** !

Differential inclusions – *Link with control system*

Key point: Control systems are **equivalent** to diff. inclusions

Equivalence (heuristic statement)

If $f : [0, T] \times \mathcal{M} \times U \rightarrow T\mathcal{M}$ is **Carathéodory**, then

$$\dot{x}(t) = f(t, x(t), u(t)) \iff \dot{x}(t) \in F(t, x(t)),$$

where

$$F(t, x) := \left\{ f(t, x, u) \text{ s.t. } u \in U \right\}.$$

Recap': Building differential inclusions

- 1) Check that the ambient space has a **manifold-like** structure,
- 2) Identify an ODE structure,
- 3) Make the velocities **set-valued** !

Differential inclusions – *Link with control system*

Key point: Control systems are **equivalent** to diff. inclusions

Equivalence (heuristic statement)

If $f : [0, T] \times \mathcal{M} \times U \rightarrow T\mathcal{M}$ is **Carathéodory**, then

$$\dot{x}(t) = f(t, x(t), u(t)) \iff \dot{x}(t) \in F(t, x(t)),$$

where

$$F(t, x) := \left\{ f(t, x, u) \text{ s.t. } u \in U \right\}.$$

Recap': Building differential inclusions

- 1) Check that the ambient space has a **manifold-like** structure,
- 2) Identify an ODE structure,
- 3) Make the velocities **set-valued** !

Differential inclusions – *Link with control system*

Key point: Control systems are **equivalent** to diff. inclusions

Equivalence (heuristic statement)

If $f : [0, T] \times \mathcal{M} \times U \rightarrow T\mathcal{M}$ is **Carathéodory**, then

$$\dot{x}(t) = f(t, x(t), u(t)) \iff \dot{x}(t) \in F(t, x(t)),$$

where


$$F(t, x) := \left\{ f(t, x, u) \text{ s.t. } u \in U \right\}.$$

Recap': Building differential inclusions

- 1) Check that the ambient space has a **manifold-like** structure,
- 2) Identify an ODE structure,
- 3) Make the velocities **set-valued** !

Differential inclusions – *What about continuity equations ?*

Idea: Endow $\mathcal{P}_c(\mathbb{R}^d)$ with **Wasserstein** metrics W_p

$$W_p(\mu, \nu) := \min_{\gamma \in \Gamma(\mu, \nu)} \left(\int_{\mathbb{R}^{2d}} |x-y|^p d\gamma(x, y) \right)^{1/p}$$


Wasserstein geometry & ODEs [Ambrosio, Gigli, McCann, Otto, Savaré]

- ◇ $(\mathcal{P}_2(\mathbb{R}^d), W_2)$ is a “**manifold**” with $T_\mu \mathcal{P}_2(\mathbb{R}^d) \subset L^2(\mu)$
- ◇ Continuity equations


$$\partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), \cdot) \mu(t)) = 0,$$

are **ODEs** over $\mathcal{P}_2(\mathbb{R}^d)$ with velocities $v(t, \mu(t), \cdot) \in L^2(\mu(t))$

Conclusion: Make $(t, \mu) \mapsto v(t, \mu, \cdot) \in L^2(\mu)$ **set-valued** !

Differential inclusions – *What about continuity equations ?*

Idea: Endow $\mathcal{P}_c(\mathbb{R}^d)$ with **Wasserstein** metrics W_p

$$W_p(\mu, \nu) := \min_{\gamma \in \Gamma(\mu, \nu)} \left(\int_{\mathbb{R}^{2d}} |x-y|^p d\gamma(x, y) \right)^{1/p}$$


Wasserstein geometry & ODEs [Ambrosio, Gigli, McCann, Otto, Savaré]

- ◇ $(\mathcal{P}_2(\mathbb{R}^d), W_2)$ is a “**manifold**” with $T_\mu \mathcal{P}_2(\mathbb{R}^d) \subset L^2(\mu)$
- ◇ Continuity equations


$$\partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), \cdot) \mu(t)) = 0,$$

are **ODEs** over $\mathcal{P}_2(\mathbb{R}^d)$ with velocities $v(t, \mu(t), \cdot) \in L^2(\mu(t))$

Conclusion: Make $(t, \mu) \mapsto v(t, \mu, \cdot) \in L^2(\mu)$ **set-valued** !

Differential inclusions – *What about continuity equations ?*

Idea: Endow $\mathcal{P}_c(\mathbb{R}^d)$ with **Wasserstein** metrics W_p

$$W_p(\mu, \nu) := \min_{\gamma \in \Gamma(\mu, \nu)} \left(\int_{\mathbb{R}^{2d}} |x-y|^p d\gamma(x, y) \right)^{1/p}$$


Wasserstein geometry & ODEs [Ambrosio, Gigli, McCann, Otto, Savaré]

- ◇ $(\mathcal{P}_2(\mathbb{R}^d), W_2)$ is a “**manifold**” with $T_\mu \mathcal{P}_2(\mathbb{R}^d) \subset L^2(\mu)$
- ◇ Continuity equations


$$\partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), \cdot) \mu(t)) = 0,$$

are **ODEs** over $\mathcal{P}_2(\mathbb{R}^d)$ with velocities $v(t, \mu(t), \cdot) \in L^2(\mu(t))$

Conclusion: Make $(t, \mu) \mapsto v(t, \mu, \cdot) \in L^2(\mu)$ **set-valued** !

Differential inclusions – *What about continuity equations ?*

Idea: Endow $\mathcal{P}_c(\mathbb{R}^d)$ with **Wasserstein** metrics W_p

$$W_p(\mu, \nu) := \min_{\gamma \in \Gamma(\mu, \nu)} \left(\int_{\mathbb{R}^{2d}} |x-y|^p d\gamma(x, y) \right)^{1/p}$$


Wasserstein geometry & ODEs [Ambrosio, Gigli, McCann, Otto, Savaré]

- ◇ $(\mathcal{P}_2(\mathbb{R}^d), W_2)$ is a “**manifold**” with $T_\mu \mathcal{P}_2(\mathbb{R}^d) \subset L^2(\mu)$
- ◇ Continuity equations


$$\partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), \cdot) \mu(t)) = 0,$$

are **ODEs** over $\mathcal{P}_2(\mathbb{R}^d)$ with velocities $v(t, \mu(t), \cdot) \in L^2(\mu(t))$

Conclusion: Make $(t, \mu) \mapsto v(t, \mu, \cdot) \in L^2(\mu)$ **set-valued** !

Differential inclusions – *What about continuity equations ?*

Idea: Endow $\mathcal{P}_c(\mathbb{R}^d)$ with **Wasserstein** metrics W_p

$$W_p(\mu, \nu) := \min_{\gamma \in \Gamma(\mu, \nu)} \left(\int_{\mathbb{R}^{2d}} |x-y|^p d\gamma(x, y) \right)^{1/p}$$


Wasserstein geometry & ODEs [Ambrosio, Gigli, McCann, Otto, Savaré]

- ◇ $(\mathcal{P}_2(\mathbb{R}^d), W_2)$ is a “**manifold**” with $T_\mu \mathcal{P}_2(\mathbb{R}^d) \subset L^2(\mu)$
- ◇ Continuity equations


$$\partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), \cdot) \mu(t)) = 0,$$

are **ODEs** over $\mathcal{P}_2(\mathbb{R}^d)$ with velocities $v(t, \mu(t), \cdot) \in L^2(\mu(t))$

Conclusion: Make $(t, \mu) \mapsto v(t, \mu, \cdot) \in L^2(\mu)$ **set-valued** !

Differential inclusions – *What about continuity equations ?*

Idea: Endow $\mathcal{P}_c(\mathbb{R}^d)$ with **Wasserstein** metrics W_p

$$W_p(\mu, \nu) := \min_{\gamma \in \Gamma(\mu, \nu)} \left(\int_{\mathbb{R}^{2d}} |x-y|^p d\gamma(x, y) \right)^{1/p}$$


Wasserstein geometry & ODEs [Ambrosio, Gigli, McCann, Otto, Savaré]

- ◇ $(\mathcal{P}_2(\mathbb{R}^d), W_2)$ is a “**manifold**” with $T_\mu \mathcal{P}_2(\mathbb{R}^d) \subset L^2(\mu)$
- ◇ Continuity equations

$$\partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), \cdot) \mu(t)) = 0,$$

are **ODEs** over $\mathcal{P}_2(\mathbb{R}^d)$ with velocities $v(t, \mu(t), \cdot) \in L^2(\mu(t))$

Conclusion: Make $(t, \mu) \mapsto v(t, \mu, \cdot) \in L^2(\mu)$ **set-valued** !

Continuity inclusions – *Definition*

Definition (Continuity inclusions) [B., Frankowska '21]

Let $(t, \mu) \in [0, T] \times \mathcal{P}_2(\mathbb{R}^d) \Rightarrow V(t, \mu) \subset L^2(\mu)$. Then, $\mu(\cdot)$ solves

$$\partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \quad (\text{CI})$$

if there exists a meas. selection $t \in [0, T] \mapsto \mathbf{v}(t) \in V(t, \mu(t))$ s.t.

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t)\mu(t)) = 0. \quad (\text{CE})$$

Literature comparison (Gradient and Hamiltonian flows)

- ◊ $V(t, \mu(t)) = -\partial\phi(\mu(t)) \rightsquigarrow$ **Gradient flow** [AGS'08 & many]
- ◊ $V(t, \mu(t)) = \mathbb{J}_{2d}\partial\mathbb{H}(\mu(t)) \rightsquigarrow$ **Hamiltonian flow** [AG'08 & PMP]

Continuity inclusions – Definition

Definition (Continuity inclusions) [B., Frankowska '21]

Let $(t, \mu) \in [0, T] \times \mathcal{P}_2(\mathbb{R}^d) \Rightarrow V(t, \mu) \subset L^2(\mu)$. Then, $\mu(\cdot)$ solves

$$\partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \quad (\text{CI})$$

if there exists a meas. selection $t \in [0, T] \mapsto \mathbf{v}(t) \in V(t, \mu(t))$ s.t.

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t)\mu(t)) = 0. \quad (\text{CE})$$

Literature comparison (Gradient and Hamiltonian flows)

- ◊ $V(t, \mu(t)) = -\partial\phi(\mu(t)) \rightsquigarrow$ Gradient flow [AGS'08 & many]
- ◊ $V(t, \mu(t)) = \mathbb{J}_{2d}\partial\mathbb{H}(\mu(t)) \rightsquigarrow$ Hamiltonian flow [AG'08 & PMP]

Continuity inclusions – Definition

Definition (Continuity inclusions) [B., Frankowska '21]

Let $(t, \mu) \in [0, T] \times \mathcal{P}_2(\mathbb{R}^d) \Rightarrow V(t, \mu) \subset L^2(\mu)$. Then, $\mu(\cdot)$ solves

$$\partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \quad (\text{CI})$$

if there exists a meas. selection $t \in [0, T] \mapsto \mathbf{v}(t) \in V(t, \mu(t))$ s.t.

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t)\mu(t)) = 0. \quad (\text{CE})$$

Literature comparison (Gradient and Hamiltonian flows)

◊ $V(t, \mu(t)) = -\partial\phi(\mu(t)) \rightsquigarrow$ Gradient flow [AGS'08 & many]

◊ $V(t, \mu(t)) = \mathbb{J}_{2d}\partial\mathbb{H}(\mu(t)) \rightsquigarrow$ Hamiltonian flow [AG'08 & PMP]

Continuity inclusions – *Definition*

Definition (Continuity inclusions) [B., Frankowska '21]

Let $(t, \mu) \in [0, T] \times \mathcal{P}_2(\mathbb{R}^d) \Rightarrow V(t, \mu) \subset L^2(\mu)$. Then, $\mu(\cdot)$ solves

$$\partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \quad (\text{CI})$$

if there exists a meas. selection $t \in [0, T] \mapsto \mathbf{v}(t) \in V(t, \mu(t))$ s.t.

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t)\mu(t)) = 0. \quad (\text{CE})$$

Literature comparison (Gradient and Hamiltonian flows)

- ◇ $V(t, \mu(t)) = -\partial\phi(\mu(t)) \rightsquigarrow$ **Gradient flow** [AGS'08 & many]
- ◇ $V(t, \mu(t)) = \mathbb{J}_{2d}\partial\mathbb{H}(\mu(t)) \rightsquigarrow$ **Hamiltonian flow** [AG'08 & PMP]

Continuity inclusions – *Definition*

Definition (Continuity inclusions) [B., Frankowska '21]

Let $(t, \mu) \in [0, T] \times \mathcal{P}_2(\mathbb{R}^d) \Rightarrow V(t, \mu) \subset L^2(\mu)$. Then, $\mu(\cdot)$ solves

$$\partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \quad (\text{CI})$$

if there exists a meas. selection $t \in [0, T] \mapsto \mathbf{v}(t) \in V(t, \mu(t))$ s.t.

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t)\mu(t)) = 0. \quad (\text{CE})$$

Literature comparison (Gradient and Hamiltonian flows)

◇ $V(t, \mu(t)) = -\partial\phi(\mu(t)) \rightsquigarrow$ **Gradient flow** [AGS'08 & many]

◇ $V(t, \mu(t)) = \mathbb{J}_{2d}\partial\mathcal{H}(\mu(t)) \rightsquigarrow$ **Hamiltonian flow** [AG'08 & PMP]

Continuity inclusions – *Definition*

Definition (Continuity inclusions) [B., Frankowska '21]

Let $(t, \mu) \in [0, T] \times \mathcal{P}_2(\mathbb{R}^d) \Rightarrow V(t, \mu) \subset L^2(\mu)$. Then, $\mu(\cdot)$ solves

$$\partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \quad (\text{CI})$$

if there exists a meas. selection $t \in [0, T] \mapsto \mathbf{v}(t) \in V(t, \mu(t))$ s.t.

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t)\mu(t)) = 0. \quad (\text{CE})$$

Literature comparison (Gradient and Hamiltonian flows)

- ◇ $V(t, \mu(t)) = -\partial\phi(\mu(t)) \rightsquigarrow$ **Gradient flow** [AGS'08 & many]
- ◇ $V(t, \mu(t)) = \mathbb{J}_{2d}\partial\mathbb{H}(\mu(t)) \rightsquigarrow$ **Hamiltonian flow** [AG'08 & PMP]

Continuity inclusions – *Link with control systems*

Certificate: Is this notion suited to control systems ? \rightsquigarrow **Yes!**

Theorem (Correspondence with control systems) [B., Frankowska '21]

Take $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times U \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is **Cauchy-Lip** and set

$$V(t, \mu) := v(t, \mu, U) \subset C^0(\mathbb{R}^d, \mathbb{R}^d).$$

Then $\mu(\cdot)$ solves (CI) $\iff \mu(\cdot)$ solves (CE) driven by

$$t \in [0, T] \mapsto \mathbf{v}(t) := v(t, \mu(t), u(t)) \in C^0(\mathbb{R}^d, \mathbb{R}^d).$$

Remark (Closed-loop controls)

For closed-loop controls, the previous result holds with

$$t \in [0, T] \mapsto \hat{\mathbf{v}}(t) := v(t, \mu(t), u(t, \cdot), \cdot) \in C^0(\mathbb{R}^d, \mathbb{R}^d),$$

by identifying $(t, x) \mapsto u(t, x) \in U$ and $t \mapsto u(t, \cdot) \in C^0(\mathbb{R}^d, U)$.

Continuity inclusions – *Link with control systems*

Certificate: Is this notion suited to control systems ? \rightsquigarrow **Yes!**

Theorem (Correspondence with control systems) [B., Frankowska '21]

Take $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times U \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is **Cauchy-Lip** and set

$$V(t, \mu) := v(t, \mu, U) \subset C^0(\mathbb{R}^d, \mathbb{R}^d).$$

Then $\mu(\cdot)$ solves (CI) $\iff \mu(\cdot)$ solves (CE) driven by

$$t \in [0, T] \mapsto \mathbf{v}(t) := v(t, \mu(t), u(t)) \in C^0(\mathbb{R}^d, \mathbb{R}^d).$$

Remark (Closed-loop controls)

For closed-loop controls, the previous result holds with

$$t \in [0, T] \mapsto \hat{\mathbf{v}}(t) := v(t, \mu(t), u(t, \cdot), \cdot) \in C^0(\mathbb{R}^d, \mathbb{R}^d),$$

by identifying $(t, x) \mapsto u(t, x) \in U$ and $t \mapsto u(t, \cdot) \in C^0(\mathbb{R}^d, U)$.

Continuity inclusions – *Link with control systems*

Certificate: Is this notion suited to control systems ? \rightsquigarrow **Yes!**

Theorem (Correspondence with control systems) [B., Frankowska '21]

Take $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times U \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is **Cauchy-Lip** and set

$$V(t, \mu) := v(t, \mu, U) \subset C^0(\mathbb{R}^d, \mathbb{R}^d).$$

Then $\mu(\cdot)$ solves (CI) $\iff \mu(\cdot)$ solves (CE) driven by

$$t \in [0, T] \mapsto \mathbf{v}(t) := v(t, \mu(t), u(t)) \in C^0(\mathbb{R}^d, \mathbb{R}^d).$$

Remark (Closed-loop controls)

For closed-loop controls, the previous result holds with

$$t \in [0, T] \mapsto \hat{\mathbf{v}}(t) := v(t, \mu(t), u(t, \cdot), \cdot) \in C^0(\mathbb{R}^d, \mathbb{R}^d),$$

by identifying $(t, x) \mapsto u(t, x) \in U$ and $t \mapsto u(t, \cdot) \in C^0(\mathbb{R}^d, U)$.

Continuity inclusions – *Link with control systems*

Certificate: Is this notion suited to control systems ? \rightsquigarrow **Yes!**

Theorem (Correspondence with control systems) [B., Frankowska '21]

Take $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times U \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is **Cauchy-Lip** and set

$$V(t, \mu) := v(t, \mu, U) \subset C^0(\mathbb{R}^d, \mathbb{R}^d).$$

Then $\mu(\cdot)$ solves (CI) $\iff \mu(\cdot)$ solves (CE) driven by

$$t \in [0, T] \mapsto \mathbf{v}(t) := v(t, \mu(t), u(t)) \in C^0(\mathbb{R}^d, \mathbb{R}^d).$$

Remark (Closed-loop controls)

For closed-loop controls, the previous result holds with

$$t \in [0, T] \mapsto \hat{\mathbf{v}}(t) := v(t, \mu(t), u(t, \cdot), \cdot) \in C^0(\mathbb{R}^d, \mathbb{R}^d),$$

by identifying $(t, x) \mapsto u(t, x) \in U$ and $t \mapsto u(t, \cdot) \in C^0(\mathbb{R}^d, U)$.

Continuity inclusions – *Link with control systems*

Certificate: Is this notion suited to control systems ? \rightsquigarrow **Yes!**

Theorem (Correspondence with control systems) [B., Frankowska '21]

Take $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times U \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is **Cauchy-Lip** and set

$$V(t, \mu) := v(t, \mu, U) \subset C^0(\mathbb{R}^d, \mathbb{R}^d).$$

Then $\mu(\cdot)$ solves (CI) $\iff \mu(\cdot)$ solves (CE) driven by

$$t \in [0, T] \mapsto \mathbf{v}(t) := v(t, \mu(t), u(t)) \in C^0(\mathbb{R}^d, \mathbb{R}^d).$$

Remark (Closed-loop controls)

For closed-loop controls, the previous result holds with

$$t \in [0, T] \mapsto \hat{\mathbf{v}}(t) := v(t, \mu(t), u(t, \cdot), \cdot) \in C^0(\mathbb{R}^d, \mathbb{R}^d),$$

by identifying $(t, x) \mapsto u(t, x) \in U$ and $t \mapsto u(t, \cdot) \in C^0(\mathbb{R}^d, U)$.

Outline of the talk

Mean-field control problems

From differential to continuity inclusions

Main structural results in the Cauchy-Lipschitz case

Application : geometric derivation of the PMP

Conclusion

Continuity inclusions – *Filippov estimates (1)*

Theorem (Filippov)[B., Frankowska '21]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip** with **compact** images, let $\nu(\cdot)$ solve

$$\partial_t \nu(t) + \operatorname{div}(\mathbf{w}(t)\nu(t)) = 0,$$

with $\mathbf{w} : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ Carathéodory, and assume that

$$\eta : t \in [0, T] \mapsto \operatorname{dist}(\mathbf{w}(t); V(t, \nu(t))),$$

is integrable. Then for any μ^0 , there exists $\mu(\cdot)$ sol. of (CI) s.t.

$$\sup_{t \in [0, T]} W_p(\mu(t), \nu(t)) \leq C_p \left(W_p(\mu^0, \nu(0)) + \int_0^T \eta(t) dt \right).$$

Continuity inclusions – *Filippov estimates (1)*

Theorem (Filippov)[B., Frankowska '21]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip** with **compact** images, let $\nu(\cdot)$ solve

$$\partial_t \nu(t) + \operatorname{div}(\mathbf{w}(t)\nu(t)) = 0,$$

with $\mathbf{w} : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ Carathéodory, and assume that

$$\eta : t \in [0, T] \mapsto \operatorname{dist}(\mathbf{w}(t); V(t, \nu(t))),$$

is integrable. Then for any μ^0 , there exists $\mu(\cdot)$ sol. of (CI) s.t.

$$\sup_{t \in [0, T]} W_p(\mu(t), \nu(t)) \leq C_p \left(W_p(\mu^0, \nu(0)) + \int_0^T \eta(t) dt \right).$$

Continuity inclusions – *Filippov estimates (1)*

Theorem (Filippov)[B., Frankowska '21]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip** with **compact** images, let $\nu(\cdot)$ solve

$$\partial_t \nu(t) + \operatorname{div}(\mathbf{w}(t)\nu(t)) = 0,$$

with $\mathbf{w} : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ Carathéodory, and assume that

$$\eta : t \in [0, T] \mapsto \operatorname{dist}(\mathbf{w}(t); V(t, \nu(t))),$$

is integrable. Then for any μ^0 , there exists $\mu(\cdot)$ sol. of (CI) s.t.

$$\sup_{t \in [0, T]} W_p(\mu(t), \nu(t)) \leq C_p \left(W_p(\mu^0, \nu(0)) + \int_0^T \eta(t) dt \right).$$

Continuity inclusions – *Filippov estimates (1)*

Theorem (Filippov)[B., Frankowska '21]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip** with **compact** images, let $\nu(\cdot)$ solve

$$\partial_t \nu(t) + \operatorname{div}(\mathbf{w}(t)\nu(t)) = 0,$$

with $\mathbf{w} : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ Carathéodory, and assume that

$$\eta : t \in [0, T] \mapsto \operatorname{dist}(\mathbf{w}(t); V(t, \nu(t))),$$

is integrable. Then for any μ^0 , there exists $\mu(\cdot)$ sol. of (CI) s.t.

$$\sup_{t \in [0, T]} W_p(\mu(t), \nu(t)) \leq C_p \left(W_p(\mu^0, \nu(0)) + \int_0^T \eta(t) dt \right).$$

Continuity inclusions – *Filippov estimates (1)*

Theorem (Filippov)[B., Frankowska '21]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip** with **compact** images, let $\nu(\cdot)$ solve

$$\partial_t \nu(t) + \operatorname{div}(\mathbf{w}(t)\nu(t)) = 0,$$

with $\mathbf{w} : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ Carathéodory, and assume that

$$\eta : t \in [0, T] \mapsto \operatorname{dist}(\mathbf{w}(t); V(t, \nu(t))),$$

is integrable. Then for any μ^0 , there exists $\mu(\cdot)$ sol. of (CI) s.t.

$$\sup_{t \in [0, T]} W_p(\mu(t), \nu(t)) \leq C_p \left(W_p(\mu^0, \nu(0)) + \int_0^T \eta(t) dt \right).$$

Continuity inclusions – *Filippov estimates (1)*

Theorem (Filippov)[B., Frankowska '21]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip** with **compact** images, let $\nu(\cdot)$ solve

$$\partial_t \nu(t) + \operatorname{div}(\mathbf{w}(t)\nu(t)) = 0,$$

with $\mathbf{w} : [0, T] \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ Carathéodory, and assume that

$$\eta : t \in [0, T] \mapsto \operatorname{dist}(\mathbf{w}(t); V(t, \nu(t))),$$

is integrable. Then for any μ^0 , there exists $\mu(\cdot)$ sol. of (CI) s.t.

$$\sup_{t \in [0, T]} W_p(\mu(t), \nu(t)) \leq C_p \left(W_p(\mu^0, \nu(0)) + \int_0^T \eta(t) dt \right).$$

Continuity inclusions – Filippov's estimates (2)




Figure: Illustration of Filippov's estimate

Proof (Main ideas)

1) Build a sequence $(\mu_k(\cdot), v_k(\cdot))$ s.t.

$$\partial_t \mu_k(t) + \operatorname{div}(v_k(t) \mu_k(t)) = 0, \quad v_k(t) \in V(t, \mu_{k-1}(t)),$$

2) Cauchy in $C^0([0, T], \mathcal{P}_p(K)) \times L^1([0, T], C^0(K, \mathbb{R}^d))$,

3) Conclude since $\operatorname{Graph}(V(t, \cdot))$ is closed for \mathcal{L}^1 -a.e. $t \in [0, T]$.

Continuity inclusions – Filippov's estimates (2)




Figure: Illustration of Filippov's estimate

Proof (Main ideas)

1) Build a sequence $(\mu_k(\cdot), v_k(\cdot))$ s.t.

$$\partial_t \mu_k(t) + \operatorname{div}(v_k(t) \mu_k(t)) = 0, \quad v_k(t) \in V(t, \mu_{k-1}(t)),$$

2) Cauchy in $C^0([0, T], \mathcal{P}_p(K)) \times L^1([0, T], C^0(K, \mathbb{R}^d))$,

3) Conclude since $\operatorname{Graph}(V(t, \cdot))$ is closed for \mathcal{L}^1 -a.e. $t \in [0, T]$.

Continuity inclusions – Filippov's estimates (2)




Figure: Illustration of Filippov's estimate

Proof (Main ideas)

1) Build a sequence $(\mu_k(\cdot), v_k(\cdot))$ s.t.

$$\partial_t \mu_k(t) + \operatorname{div}(v_k(t) \mu_k(t)) = 0, \quad v_k(t) \in V(t, \mu_{k-1}(t)),$$

2) Cauchy in $C^0([0, T], \mathcal{P}_p(K)) \times L^1([0, T], C^0(K, \mathbb{R}^d))$,

3) Conclude since $\operatorname{Graph}(V(t, \cdot))$ is closed for \mathcal{L}^1 -a.e. $t \in [0, T]$.

Continuity inclusions – Filippov's estimates (2)




Figure: Illustration of Filippov's estimate

Proof (Main ideas)

1) Build a sequence $(\mu_k(\cdot), v_k(\cdot))$ s.t.

$$\partial_t \mu_k(t) + \operatorname{div}(v_k(t) \mu_k(t)) = 0, \quad v_k(t) \in V(t, \mu_{k-1}(t)),$$

2) Cauchy in $C^0([0, T], \mathcal{P}_p(K)) \times L^1([0, T], C^0(K, \mathbb{R}^d))$,

3) Conclude since $\operatorname{Graph}(V(t, \cdot))$ is closed for \mathcal{L}^1 -a.e. $t \in [0, T]$.

Continuity inclusions – Filippov's estimates (2)




Figure: Illustration of Filippov's estimate

Proof (Main ideas)

1) Build a sequence $(\mu_k(\cdot), \mathbf{v}_k(\cdot))$ s.t.

$$\partial_t \mu_k(t) + \operatorname{div}(\mathbf{v}_k(t) \mu_k(t)) = 0, \quad \mathbf{v}_k(t) \in V(t, \mu_{k-1}(t)),$$

2) Cauchy in $C^0([0, T], \mathcal{P}_p(K)) \times L^1([0, T], C^0(K, \mathbb{R}^d))$,

3) Conclude since $\operatorname{Graph}(V(t, \cdot))$ is closed for \mathcal{L}^1 -a.e. $t \in [0, T]$.

Continuity inclusions – Filippov's estimates (2)




Figure: Illustration of Filippov's estimate

Proof (Main ideas)

1) Build a sequence $(\mu_k(\cdot), \mathbf{v}_k(\cdot))$ s.t.

$$\partial_t \mu_k(t) + \operatorname{div}(\mathbf{v}_k(t) \mu_k(t)) = 0, \quad \mathbf{v}_k(t) \in V(t, \mu_{k-1}(t)),$$

2) Cauchy in $C^0([0, T], \mathcal{P}_p(K)) \times L^1([0, T], C^0(K, \mathbb{R}^d))$,

3) Conclude since $\operatorname{Graph}(V(t, \cdot))$ is closed for \mathcal{L}^1 -a.e. $t \in [0, T]$.

Continuity inclusions – Filippov's estimates (2)




Figure: Illustration of Filippov's estimate

Proof (Main ideas)

- 1) Build a sequence $(\mu_k(\cdot), \mathbf{v}_k(\cdot))$ s.t.

$$\partial_t \mu_k(t) + \operatorname{div}(\mathbf{v}_k(t) \mu_k(t)) = 0, \quad \mathbf{v}_k(t) \in V(t, \mu_{k-1}(t)),$$

- 2) Cauchy in $C^0([0, T], \mathcal{P}_p(K)) \times L^1([0, T], C^0(K, \mathbb{R}^d))$,

- 3) Conclude since $\operatorname{Graph}(V(t, \cdot))$ is closed for \mathcal{L}^1 -a.e. $t \in [0, T]$.

Continuity inclusions – Filippov's estimates (2)




Figure: Illustration of Filippov's estimate

Proof (Main ideas)

- 1) Build a sequence $(\mu_k(\cdot), \mathbf{v}_k(\cdot))$ s.t.

$$\partial_t \mu_k(t) + \operatorname{div}(\mathbf{v}_k(t) \mu_k(t)) = 0, \quad \mathbf{v}_k(t) \in V(t, \mu_{k-1}(t)),$$

- 2) Cauchy in $C^0([0, T], \mathcal{P}_p(K)) \times L^1([0, T], C^0(K, \mathbb{R}^d))$,
- 3) Conclude since $\operatorname{Graph}(V(t, \cdot))$ is closed for \mathcal{L}^1 -a.e. $t \in [0, T]$.

Continuity inclusions – Compactness & Relaxation (1)

Theorem (Compactness of solutions)[B., Frankowska '21]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **compact convex** images. Then for any μ^0 , the solution set

$$\mathcal{S}(\mu^0) := \left\{ \mu(\cdot) \text{ s.t. } \partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \mu(0) = \mu^0 \right\},$$

is **compact** in the $C^0 W_p$ -topology \rightsquigarrow [FS'14] \oplus **Mazur** on $V(\cdot, \cdot)$.

Theorem (Relaxation)[B., Frankowska '21]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **compact** images. Then, $\mathcal{S}(\mu^0)$ is dense in the solution set of

$$\partial_t \mu(t) \in -\operatorname{div}(\overline{\operatorname{co}} V(t, \mu(t))\mu(t)), \quad \mu(0) = \mu^0,$$

with $\overline{\operatorname{co}} V(t, \mu)$ the **closed convex hull** of $V(t, \mu) \subset C^0(\mathbb{R}^d, \mathbb{R}^d)$.

Continuity inclusions – Compactness & Relaxation (1)

Theorem (Compactness of solutions)[B., Frankowska '21]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **compact convex** images. Then for any μ^0 , the solution set

$$\mathcal{S}(\mu^0) := \left\{ \mu(\cdot) \text{ s.t. } \partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \mu(0) = \mu^0 \right\},$$

is **compact** in the $C^0 W_p$ -topology \rightsquigarrow [FS'14] \oplus Mazur on $V(\cdot, \cdot)$.

Theorem (Relaxation)[B., Frankowska '21]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **compact** images. Then, $\mathcal{S}(\mu^0)$ is dense in the solution set of

$$\partial_t \mu(t) \in -\operatorname{div}(\overline{\operatorname{co}} V(t, \mu(t))\mu(t)), \quad \mu(0) = \mu^0,$$

with $\overline{\operatorname{co}} V(t, \mu)$ the closed convex hull of $V(t, \mu) \subset C^0(\mathbb{R}^d, \mathbb{R}^d)$.

Continuity inclusions – Compactness & Relaxation (1)

Theorem (Compactness of solutions)[B., Frankowska '21]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **compact convex** images. Then for any μ^0 , the solution set

$$\mathcal{S}(\mu^0) := \left\{ \mu(\cdot) \text{ s.t. } \partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \mu(0) = \mu^0 \right\},$$

is **compact** in the $C^0 W_p$ -topology \rightsquigarrow [FS'14] \oplus Mazur on $V(\cdot, \cdot)$.

Theorem (Relaxation)[B., Frankowska '21]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **compact** images. Then, $\mathcal{S}(\mu^0)$ is dense in the solution set of

$$\partial_t \mu(t) \in -\operatorname{div}(\overline{\operatorname{co}} V(t, \mu(t))\mu(t)), \quad \mu(0) = \mu^0,$$

with $\overline{\operatorname{co}} V(t, \mu)$ the closed convex hull of $V(t, \mu) \subset C^0(\mathbb{R}^d, \mathbb{R}^d)$.

Continuity inclusions – Compactness & Relaxation (1)

Theorem (Compactness of solutions)[B., Frankowska '21]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **compact convex** images. Then for any μ^0 , the solution set

$$\mathcal{S}(\mu^0) := \left\{ \mu(\cdot) \text{ s.t. } \partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \mu(0) = \mu^0 \right\},$$

is **compact** in the $C^0 W_p$ -topology \rightsquigarrow [FS'14] \oplus **Mazur** on $V(\cdot, \cdot)$.

Theorem (Relaxation)[B., Frankowska '21]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **compact** images. Then, $\mathcal{S}(\mu^0)$ is dense in the solution set of

$$\partial_t \mu(t) \in -\operatorname{div}(\overline{\operatorname{co}} V(t, \mu(t))\mu(t)), \quad \mu(0) = \mu^0,$$

with $\overline{\operatorname{co}} V(t, \mu)$ the **closed convex hull** of $V(t, \mu) \subset C^0(\mathbb{R}^d, \mathbb{R}^d)$.

Continuity inclusions – Compactness & Relaxation (1)

Theorem (Compactness of solutions)[B., Frankowska '21]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **compact convex** images. Then for any μ^0 , the solution set

$$\mathcal{S}(\mu^0) := \left\{ \mu(\cdot) \text{ s.t. } \partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \mu(0) = \mu^0 \right\},$$

is **compact** in the $C^0 W_p$ -topology \rightsquigarrow [FS'14] \oplus **Mazur** on $V(\cdot, \cdot)$.

Theorem (Relaxation)[B., Frankowska '21]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **compact** images. Then, $\mathcal{S}(\mu^0)$ is dense in the solution set of

$$\partial_t \mu(t) \in -\operatorname{div}(\overline{\operatorname{co}} V(t, \mu(t))\mu(t)), \quad \mu(0) = \mu^0,$$

with $\overline{\operatorname{co}} V(t, \mu)$ the **closed convex hull** of $V(t, \mu) \subset C^0(\mathbb{R}^d, \mathbb{R}^d)$.

Continuity inclusions – Compactness & Relaxation (1)

Theorem (Compactness of solutions)[B., Frankowska '21]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **compact convex** images. Then for any μ^0 , the solution set

$$\mathcal{S}(\mu^0) := \left\{ \mu(\cdot) \text{ s.t. } \partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \mu(0) = \mu^0 \right\},$$

is **compact** in the $C^0 W_p$ -topology \rightsquigarrow [FS'14] \oplus **Mazur** on $V(\cdot, \cdot)$.

Theorem (Relaxation)[B., Frankowska '21]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **compact** images. Then, $\mathcal{S}(\mu^0)$ is dense in the solution set of

$$\partial_t \mu(t) \in -\operatorname{div}(\overline{\operatorname{co}} V(t, \mu(t))\mu(t)), \quad \mu(0) = \mu^0,$$

with $\overline{\operatorname{co}} V(t, \mu)$ the **closed convex hull** of $V(t, \mu) \subset C^0(\mathbb{R}^d, \mathbb{R}^d)$.

Continuity inclusions – Compactness & Relaxation (1)

Theorem (Compactness of solutions)[B., Frankowska '21]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **compact convex** images. Then for any μ^0 , the solution set

$$\mathcal{S}(\mu^0) := \left\{ \mu(\cdot) \text{ s.t. } \partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \mu(0) = \mu^0 \right\},$$

is **compact** in the $C^0 W_p$ -topology \rightsquigarrow [FS'14] \oplus **Mazur** on $V(\cdot, \cdot)$.

Theorem (Relaxation)[B., Frankowska '21]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **compact** images. Then, $\mathcal{S}(\mu^0)$ is dense in the solution set of

$$\partial_t \mu(t) \in -\operatorname{div}(\overline{\operatorname{co}} V(t, \mu(t))\mu(t)), \quad \mu(0) = \mu^0,$$

with $\overline{\operatorname{co}} V(t, \mu)$ the **closed convex hull** of $V(t, \mu) \subset C^0(\mathbb{R}^d, \mathbb{R}^d)$.

Continuity inclusions – Compactness & Relaxation (2)


Proof (Main ideas)

1) Let $\mu(\cdot)$ be a relaxed traj. By **Aumann**, there exists $\nu(\cdot)$ s.t.

$$\begin{cases} \partial_t \nu(t) + \operatorname{div}(w(t)\nu(t)) = 0 & \text{and } w(t) \in V(t, \mu(t)), \\ \sup_{t \in [0, T]} W_p(\mu(t), \nu(t)) \leq \frac{\delta}{2} & \text{for an arbitrary } \delta > 0. \end{cases}$$

2) By **Filippov**, there exists $\mu_\delta(\cdot)$ solution of (CI) from μ^0 s.t.

$$\sup_{t \in [0, T]} W_p(\nu(t), \mu_\delta(t)) \leq \frac{\delta}{2}.$$



Continuity inclusions – Compactness & Relaxation (2)


Proof (Main ideas)

1) Let $\mu(\cdot)$ be a relaxed traj. By **Aumann**, there exists $\nu(\cdot)$ s.t.

$$\begin{cases} \partial_t \nu(t) + \operatorname{div}(\mathbf{w}(t)\nu(t)) = 0 & \text{and } \mathbf{w}(t) \in V(t, \mu(t)), \\ \sup_{t \in [0, T]} W_p(\mu(t), \nu(t)) \leq \frac{\delta}{2} & \text{for an arbitrary } \delta > 0. \end{cases}$$

2) By **Filippov**, there exists $\mu_\delta(\cdot)$ solution of (CI) from μ^0 s.t.

$$\sup_{t \in [0, T]} W_p(\nu(t), \mu_\delta(t)) \leq \frac{\delta}{2}.$$



Continuity inclusions – Compactness & Relaxation (2)


Proof (Main ideas)

1) Let $\mu(\cdot)$ be a relaxed traj. By **Aumann**, there exists $\nu(\cdot)$ s.t.

$$\begin{cases} \partial_t \nu(t) + \operatorname{div}(\mathbf{w}(t)\nu(t)) = 0 & \text{and } \mathbf{w}(t) \in V(t, \mu(t)), \\ \sup_{t \in [0, T]} W_p(\mu(t), \nu(t)) \leq \frac{\delta}{2} & \text{for an arbitrary } \delta > 0. \end{cases}$$

2) By **Filippov**, there exists $\mu_\delta(\cdot)$ solution of (CI) from μ^0 s.t.

$$\sup_{t \in [0, T]} W_p(\nu(t), \mu_\delta(t)) \leq \frac{\delta}{2}.$$



Continuity inclusions – Compactness & Relaxation (2)


Proof (Main ideas)

1) Let $\mu(\cdot)$ be a relaxed traj. By **Aumann**, there exists $\nu(\cdot)$ s.t.

$$\begin{cases} \partial_t \nu(t) + \operatorname{div}(\mathbf{w}(t)\nu(t)) = 0 & \text{and } \mathbf{w}(t) \in V(t, \mu(t)), \\ \sup_{t \in [0, T]} W_p(\mu(t), \nu(t)) \leq \frac{\delta}{2} & \text{for an arbitrary } \delta > 0. \end{cases}$$

2) By **Filippov**, there exists $\mu_\delta(\cdot)$ solution of (CI) from μ^0 s.t.

$$\sup_{t \in [0, T]} W_p(\nu(t), \mu_\delta(t)) \leq \frac{\delta}{2}.$$



Continuity inclusions – Compactness & Relaxation (2)


Proof (Main ideas)

1) Let $\mu(\cdot)$ be a relaxed traj. By **Aumann**, there exists $\nu(\cdot)$ s.t.

$$\begin{cases} \partial_t \nu(t) + \operatorname{div}(\mathbf{w}(t)\nu(t)) = 0 & \text{and } \mathbf{w}(t) \in V(t, \mu(t)), \\ \sup_{t \in [0, T]} W_p(\mu(t), \nu(t)) \leq \frac{\delta}{2} & \text{for an arbitrary } \delta > 0. \end{cases}$$

2) By **Filippov**, there exists $\mu_\delta(\cdot)$ solution of (CI) from μ^0 s.t.

$$\sup_{t \in [0, T]} W_p(\nu(t), \mu_\delta(t)) \leq \frac{\delta}{2}.$$



Outline of the talk

Mean-field control problems

From differential to continuity inclusions

Main structural results in the Cauchy-Lipschitz case

Application : geometric derivation of the PMP

Conclusion

Mean-field OCP – Definition and existence

Consider the Mayer problem

$$(\mathbf{OCP}) \quad \left\{ \begin{array}{l} \min_{u(\cdot) \in \mathcal{U}} [\varphi(\mu(T))], \\ \text{s.t.} \quad \left\{ \begin{array}{l} \partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), u(t))\mu(t)) = 0, \\ \mu(0) = \mu^0, \\ \Psi_i(\mu(T)) \leq 0 \quad \text{for } i \in \{1, \dots, m\}, \end{array} \right. \end{array} \right.$$

where

- ◇ $u : [0, T] \rightarrow U$ is \mathcal{L}^1 -meas. with (U, d_U) comp. metric space.
- ◇ $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times U \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is \mathcal{L}^1 -meas. in $t \in [0, T]$, continuous in $u \in U$ and \mathcal{C}_{loc}^1 w.r.t. $(\mu, x) \in \mathcal{P}_c(\mathbb{R}^d) \times \mathbb{R}^d$.
- ◇ $\varphi : \mathcal{P}_c(\mathbb{R}^d) \rightarrow \mathbb{R}$ and $\Psi_i : \mathcal{P}_c(\mathbb{R}^d) \rightarrow \mathbb{R}$ are \mathcal{C}_{loc}^1 .

Theorem (Existence of solutions)[B., Frankowska '21]

If $v(t, \mu, U) \subset C^0(\mathbb{R}^d, \mathbb{R}^d)$ is convex, then (\mathbf{OCP}) admits a solution.

Mean-field OCP – Definition and existence

Consider the Mayer problem

$$(\mathbf{OCP}) \quad \begin{cases} \min_{u(\cdot) \in \mathcal{U}} [\varphi(\mu(T))], \\ \text{s.t.} \quad \begin{cases} \partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), u(t))\mu(t)) = 0, \\ \mu(0) = \mu^0, \\ \Psi_i(\mu(T)) \leq 0 \quad \text{for } i \in \{1, \dots, m\}, \end{cases} \end{cases}$$

where

- ◇ $u : [0, T] \rightarrow U$ is \mathcal{L}^1 -meas. with (U, d_U) comp. metric space.
- ◇ $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times U \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is \mathcal{L}^1 -meas. in $t \in [0, T]$, continuous in $u \in U$ and \mathcal{C}_{loc}^1 w.r.t. $(\mu, x) \in \mathcal{P}_c(\mathbb{R}^d) \times \mathbb{R}^d$.
- ◇ $\varphi : \mathcal{P}_c(\mathbb{R}^d) \rightarrow \mathbb{R}$ and $\Psi_i : \mathcal{P}_c(\mathbb{R}^d) \rightarrow \mathbb{R}$ are \mathcal{C}_{loc}^1 .

Theorem (Existence of solutions)[B., Frankowska '21]

If $v(t, \mu, U) \subset C^0(\mathbb{R}^d, \mathbb{R}^d)$ is convex, then (\mathbf{OCP}) admits a solution.

Mean-field OCP – Definition and existence

Consider the Mayer problem

$$(\mathbf{OCP}) \quad \left\{ \begin{array}{l} \min_{u(\cdot) \in \mathcal{U}} [\varphi(\mu(T))], \\ \text{s.t.} \quad \left\{ \begin{array}{l} \partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), u(t))\mu(t)) = 0, \\ \mu(0) = \mu^0, \\ \Psi_i(\mu(T)) \leq 0 \quad \text{for } i \in \{1, \dots, m\}, \end{array} \right. \end{array} \right.$$

where

- ◇ $u : [0, T] \rightarrow U$ is \mathcal{L}^1 -meas. with (U, d_U) comp. metric space.
- ◇ $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times U \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is \mathcal{L}^1 -meas. in $t \in [0, T]$, continuous in $u \in U$ and \mathcal{C}_{loc}^1 w.r.t. $(\mu, x) \in \mathcal{P}_c(\mathbb{R}^d) \times \mathbb{R}^d$.
- ◇ $\varphi : \mathcal{P}_c(\mathbb{R}^d) \rightarrow \mathbb{R}$ and $\Psi_i : \mathcal{P}_c(\mathbb{R}^d) \rightarrow \mathbb{R}$ are \mathcal{C}_{loc}^1 .

Theorem (Existence of solutions)[B., Frankowska '21]

If $v(t, \mu, U) \subset C^0(\mathbb{R}^d, \mathbb{R}^d)$ is convex, then (\mathbf{OCP}) admits a solution.

Mean-field OCP – Definition and existence

Consider the Mayer problem

$$(\mathbf{OCP}) \quad \left\{ \begin{array}{l} \min_{u(\cdot) \in \mathcal{U}} [\varphi(\mu(T))], \\ \text{s.t.} \quad \left\{ \begin{array}{l} \partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), u(t))\mu(t)) = 0, \\ \mu(0) = \mu^0, \\ \Psi_i(\mu(T)) \leq 0 \quad \text{for } i \in \{1, \dots, m\}, \end{array} \right. \end{array} \right.$$

where

- ◇ $u : [0, T] \rightarrow U$ is \mathcal{L}^1 -meas. with (U, d_U) comp. metric space.
- ◇ $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times U \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is \mathcal{L}^1 -meas. in $t \in [0, T]$, continuous in $u \in U$ and \mathcal{C}_{loc}^1 w.r.t. $(\mu, x) \in \mathcal{P}_c(\mathbb{R}^d) \times \mathbb{R}^d$.
- ◇ $\varphi : \mathcal{P}_c(\mathbb{R}^d) \rightarrow \mathbb{R}$ and $\Psi_i : \mathcal{P}_c(\mathbb{R}^d) \rightarrow \mathbb{R}$ are \mathcal{C}_{loc}^1 .

Theorem (Existence of solutions)[B., Frankowska '21]

If $v(t, \mu, U) \subset C^0(\mathbb{R}^d, \mathbb{R}^d)$ is convex, then (\mathbf{OCP}) admits a solution.

Mean-field OCP – Definition and existence

Consider the Mayer problem

$$(\mathbf{OCP}) \quad \left\{ \begin{array}{l} \min_{u(\cdot) \in \mathcal{U}} [\varphi(\mu(T))], \\ \text{s.t.} \quad \left\{ \begin{array}{l} \partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), u(t))\mu(t)) = 0, \\ \mu(0) = \mu^0, \\ \Psi_i(\mu(T)) \leq 0 \quad \text{for } i \in \{1, \dots, m\}, \end{array} \right. \end{array} \right.$$

where

- ◇ $u : [0, T] \rightarrow U$ is \mathcal{L}^1 -meas. with (U, d_U) comp. metric space.
- ◇ $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times U \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is \mathcal{L}^1 -meas. in $t \in [0, T]$, continuous in $u \in U$ and C_{loc}^1 w.r.t. $(\mu, x) \in \mathcal{P}_c(\mathbb{R}^d) \times \mathbb{R}^d$.
- ◇ $\varphi : \mathcal{P}_c(\mathbb{R}^d) \rightarrow \mathbb{R}$ and $\Psi_i : \mathcal{P}_c(\mathbb{R}^d) \rightarrow \mathbb{R}$ are C_{loc}^1 .

Theorem (Existence of solutions)[B., Frankowska '21]

If $v(t, \mu, U) \subset C^0(\mathbb{R}^d, \mathbb{R}^d)$ is convex, then (\mathbf{OCP}) admits a solution.

Mean-field OCP – Definition and existence

Consider the Mayer problem

$$(\mathbf{OCP}) \quad \left\{ \begin{array}{l} \min_{u(\cdot) \in \mathcal{U}} [\varphi(\mu(T))], \\ \text{s.t.} \quad \left\{ \begin{array}{l} \partial_t \mu(t) + \operatorname{div}(v(t, \mu(t), u(t))\mu(t)) = 0, \\ \mu(0) = \mu^0, \\ \Psi_i(\mu(T)) \leq 0 \quad \text{for } i \in \{1, \dots, m\}, \end{array} \right. \end{array} \right.$$

where

- ◇ $u : [0, T] \rightarrow U$ is \mathcal{L}^1 -meas. with (U, d_U) comp. metric space.
- ◇ $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times U \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is \mathcal{L}^1 -meas. in $t \in [0, T]$, continuous in $u \in U$ and $\mathcal{C}_{\text{loc}}^1$ w.r.t. $(\mu, x) \in \mathcal{P}_c(\mathbb{R}^d) \times \mathbb{R}^d$.
- ◇ $\varphi : \mathcal{P}_c(\mathbb{R}^d) \rightarrow \mathbb{R}$ and $\Psi_i : \mathcal{P}_c(\mathbb{R}^d) \rightarrow \mathbb{R}$ are $\mathcal{C}_{\text{loc}}^1$.

Theorem (Existence of solutions)[B., Frankowska '21]

If $v(t, \mu, U) \subset C^0(\mathbb{R}^d, \mathbb{R}^d)$ is convex, then **(OCP)** admits a solution.

Mean-field OCP – C_{loc}^1 -functionals

Definition (Local W_2 -sub and superdifferentials)[B., Frankowska '21]

Let $\phi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R} \cup \{\pm\infty\}$ be s.t. $\mathcal{P}_c(\mathbb{R}^d) \subset D(\phi)$ and $\mu \in \mathcal{P}_c(\mathbb{R}^d)$. Then, $\xi \in \partial_{\text{loc}}^- \phi(\mu)$ if and only if

$$\phi(\nu) - \phi(\mu) \geq \inf_{\gamma \in \Gamma_o(\mu, \nu)} \int_{\mathbb{R}^{2d}} \langle \xi(x), y - x \rangle d\gamma(x, y) + o_R(W_2(\mu, \nu)),$$

for any $R > 0$ and all $\nu \in \mathcal{P}(B_R(\mu)) \rightsquigarrow R$ -fattening of $\text{supp}(\mu)$.

Theorem (Wasserstein gradients)[GT'18 & B., Frankowska '21]

If $\partial_{\text{loc}}^- \phi(\mu)$ and $\partial_{\text{loc}}^+ \phi(\mu)$ are both **non-empty**, then necessarily $\partial_{\text{loc}}^- \phi(\mu) \cap \partial_{\text{loc}}^+ \phi(\mu) = \{\nabla \phi(\mu)\}$. Moreover, it holds

$$\phi(\nu) = \phi(\mu) + \int_{\mathbb{R}^{2d}} \langle \nabla \phi(\mu), y - x \rangle d\mu(x, y) + o(W_{2,\mu}(\mu, \nu)),$$

for all $\mu \in \Gamma(\mu, \nu) \rightsquigarrow$ Nice for **control theory** !

Mean-field OCP – C_{loc}^1 -functionals

Definition (Local W_2 -sub and superdifferentials)[B., Frankowska '21]

Let $\phi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R} \cup \{\pm\infty\}$ be s.t. $\mathcal{P}_c(\mathbb{R}^d) \subset D(\phi)$ and $\mu \in \mathcal{P}_c(\mathbb{R}^d)$. Then, $\xi \in \partial_{\text{loc}}^- \phi(\mu)$ if and only if

$$\phi(\nu) - \phi(\mu) \geq \inf_{\gamma \in \Gamma_o(\mu, \nu)} \int_{\mathbb{R}^{2d}} \langle \xi(x), y - x \rangle d\gamma(x, y) + o_R(W_2(\mu, \nu)),$$

for any $R > 0$ and all $\nu \in \mathcal{P}(B_R(\mu)) \rightsquigarrow R$ -fattening of $\text{supp}(\mu)$.

Theorem (Wasserstein gradients)[GT'18 & B., Frankowska '21]

If $\partial_{\text{loc}}^- \phi(\mu)$ and $\partial_{\text{loc}}^+ \phi(\mu)$ are both **non-empty**, then necessarily $\partial_{\text{loc}}^- \phi(\mu) \cap \partial_{\text{loc}}^+ \phi(\mu) = \{\nabla \phi(\mu)\}$. Moreover, it holds

$$\phi(\nu) = \phi(\mu) + \int_{\mathbb{R}^{2d}} \langle \nabla \phi(\mu), y - x \rangle d\mu(x, y) + o(W_{2,\mu}(\mu, \nu)),$$

for all $\mu \in \Gamma(\mu, \nu) \rightsquigarrow$ Nice for **control theory** !

Mean-field OCP – C_{loc}^1 -functionals

Definition (Local W_2 -sub and superdifferentials)[B., Frankowska '21]

Let $\phi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R} \cup \{\pm\infty\}$ be s.t. $\mathcal{P}_c(\mathbb{R}^d) \subset D(\phi)$ and $\mu \in \mathcal{P}_c(\mathbb{R}^d)$. Then, $\xi \in \partial_{\text{loc}}^- \phi(\mu)$ if and only if

$$\phi(\nu) - \phi(\mu) \geq \inf_{\gamma \in \Gamma_o(\mu, \nu)} \int_{\mathbb{R}^{2d}} \langle \xi(x), y - x \rangle d\gamma(x, y) + o_R(W_2(\mu, \nu)),$$

for any $R > 0$ and all $\nu \in \mathcal{P}(B_R(\mu)) \rightsquigarrow R$ -fattening of $\text{supp}(\mu)$.

Theorem (Wasserstein gradients)[GT'18 & B., Frankowska '21]

If $\partial_{\text{loc}}^- \phi(\mu)$ and $\partial_{\text{loc}}^+ \phi(\mu)$ are both **non-empty**, then necessarily $\partial_{\text{loc}}^- \phi(\mu) \cap \partial_{\text{loc}}^+ \phi(\mu) = \{\nabla \phi(\mu)\}$. Moreover, it holds

$$\phi(\nu) = \phi(\mu) + \int_{\mathbb{R}^{2d}} \langle \nabla \phi(\mu), y - x \rangle d\mu(x, y) + o(W_{2,\mu}(\mu, \nu)),$$

for all $\mu \in \Gamma(\mu, \nu) \rightsquigarrow$ Nice for **control theory** !

Mean-field OCP – C_{loc}^1 -functionals

Definition (Local W_2 -sub and superdifferentials)[B., Frankowska '21]

Let $\phi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R} \cup \{\pm\infty\}$ be s.t. $\mathcal{P}_c(\mathbb{R}^d) \subset D(\phi)$ and $\mu \in \mathcal{P}_c(\mathbb{R}^d)$. Then, $\xi \in \partial_{\text{loc}}^- \phi(\mu)$ if and only if

$$\phi(\nu) - \phi(\mu) \geq \inf_{\gamma \in \Gamma_o(\mu, \nu)} \int_{\mathbb{R}^{2d}} \langle \xi(x), y - x \rangle d\gamma(x, y) + o_R(W_2(\mu, \nu)),$$

for any $R > 0$ and all $\nu \in \mathcal{P}(B_R(\mu)) \rightsquigarrow R$ -fattening of $\text{supp}(\mu)$.

Theorem (Wasserstein gradients)[GT'18 & B., Frankowska '21]

If $\partial_{\text{loc}}^- \phi(\mu)$ and $\partial_{\text{loc}}^+ \phi(\mu)$ are both **non-empty**, then necessarily $\partial_{\text{loc}}^- \phi(\mu) \cap \partial_{\text{loc}}^+ \phi(\mu) = \{\nabla \phi(\mu)\}$. Moreover, it holds

$$\phi(\nu) = \phi(\mu) + \int_{\mathbb{R}^{2d}} \langle \nabla \phi(\mu), y - x \rangle d\mu(x, y) + o(W_{2,\mu}(\mu, \nu)),$$

for all $\mu \in \Gamma(\mu, \nu) \rightsquigarrow$ Nice for **control theory** !

Mean-field OCP – C_{loc}^1 -functionals

Definition (Local W_2 -sub and superdifferentials)[B., Frankowska '21]

Let $\phi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R} \cup \{\pm\infty\}$ be s.t. $\mathcal{P}_c(\mathbb{R}^d) \subset D(\phi)$ and $\mu \in \mathcal{P}_c(\mathbb{R}^d)$. Then, $\xi \in \partial_{\text{loc}}^- \phi(\mu)$ if and only if

$$\phi(\nu) - \phi(\mu) \geq \inf_{\gamma \in \Gamma_o(\mu, \nu)} \int_{\mathbb{R}^{2d}} \langle \xi(x), y - x \rangle d\gamma(x, y) + o_R(W_2(\mu, \nu)),$$

for any $R > 0$ and all $\nu \in \mathcal{P}(B_R(\mu)) \rightsquigarrow R$ -fattening of $\text{supp}(\mu)$.

Theorem (Wasserstein gradients)[GT'18 & B., Frankowska '21]

If $\partial_{\text{loc}}^- \phi(\mu)$ and $\partial_{\text{loc}}^+ \phi(\mu)$ are both **non-empty**, then necessarily $\partial_{\text{loc}}^- \phi(\mu) \cap \partial_{\text{loc}}^+ \phi(\mu) = \{\nabla \phi(\mu)\}$. Moreover, it holds

$$\phi(\nu) = \phi(\mu) + \int_{\mathbb{R}^{2d}} \langle \nabla \phi(\mu), y - x \rangle d\mu(x, y) + o(W_{2,\mu}(\mu, \nu)),$$

for all $\mu \in \Gamma(\mu, \nu) \rightsquigarrow$ Nice for **control theory** !

Mean-field OCP – C_{loc}^1 -functionals

Definition (Local W_2 -sub and superdifferentials)[B., Frankowska '21]

Let $\phi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R} \cup \{\pm\infty\}$ be s.t. $\mathcal{P}_c(\mathbb{R}^d) \subset D(\phi)$ and $\mu \in \mathcal{P}_c(\mathbb{R}^d)$. Then, $\xi \in \partial_{\text{loc}}^- \phi(\mu)$ if and only if

$$\phi(\nu) - \phi(\mu) \geq \inf_{\gamma \in \Gamma_o(\mu, \nu)} \int_{\mathbb{R}^{2d}} \langle \xi(x), y - x \rangle d\gamma(x, y) + o_R(W_2(\mu, \nu)),$$

for any $R > 0$ and all $\nu \in \mathcal{P}(B_R(\mu)) \rightsquigarrow R$ -fattening of $\text{supp}(\mu)$.

Theorem (Wasserstein gradients)[GT'18 & B., Frankowska '21]

If $\partial_{\text{loc}}^- \phi(\mu)$ and $\partial_{\text{loc}}^+ \phi(\mu)$ are both **non-empty**, then necessarily $\partial_{\text{loc}}^- \phi(\mu) \cap \partial_{\text{loc}}^+ \phi(\mu) = \{\nabla \phi(\mu)\}$. Moreover, it holds

$$\phi(\nu) = \phi(\mu) + \int_{\mathbb{R}^{2d}} \langle \nabla \phi(\mu), y - x \rangle d\mu(x, y) + o(W_{2, \mu}(\mu, \nu)),$$

for all $\mu \in \Gamma(\mu, \nu) \rightsquigarrow$ Nice for **control theory** !

Mean-field OCP – C_{loc}^1 -functionals

Definition (Local W_2 -sub and superdifferentials)[B., Frankowska '21]

Let $\phi : \mathcal{P}_2(\mathbb{R}^d) \rightarrow \mathbb{R} \cup \{\pm\infty\}$ be s.t. $\mathcal{P}_c(\mathbb{R}^d) \subset D(\phi)$ and $\mu \in \mathcal{P}_c(\mathbb{R}^d)$. Then, $\xi \in \partial_{\text{loc}}^- \phi(\mu)$ if and only if

$$\phi(\nu) - \phi(\mu) \geq \inf_{\gamma \in \Gamma_o(\mu, \nu)} \int_{\mathbb{R}^{2d}} \langle \xi(x), y - x \rangle d\gamma(x, y) + o_R(W_2(\mu, \nu)),$$

for any $R > 0$ and all $\nu \in \mathcal{P}(B_R(\mu)) \rightsquigarrow R$ -fattening of $\text{supp}(\mu)$.

Theorem (Wasserstein gradients)[GT'18 & B., Frankowska '21]

If $\partial_{\text{loc}}^- \phi(\mu)$ and $\partial_{\text{loc}}^+ \phi(\mu)$ are both **non-empty**, then necessarily $\partial_{\text{loc}}^- \phi(\mu) \cap \partial_{\text{loc}}^+ \phi(\mu) = \{\nabla \phi(\mu)\}$. Moreover, it holds

$$\phi(\nu) = \phi(\mu) + \int_{\mathbb{R}^{2d}} \langle \nabla \phi(\mu), y - x \rangle d\mu(x, y) + o(W_{2, \mu}(\mu, \nu)),$$

for all $\mu \in \Gamma(\mu, \nu) \rightsquigarrow$ Nice for **control theory** !

Mean-field OCP – PMP for smooth Mayer problems (1)

Pontryagin Maximum Principle for **(OCP)** [B., Frankowska '21]

Let $(\mu^*(\cdot), u^*(\cdot))$ be a strong local min. for **(OCP)**. Then, there exist **non-trivial multipliers** $(\lambda_0, \lambda_1, \dots, \lambda_n) \in \{0, 1\} \times \mathbb{R}_+^n$ and a **state-costate curve** $\nu^*(\cdot) \in \text{AC}([0, T], \mathcal{P}_c(\mathbb{R}^{2d}))$ s.t.

$$\begin{cases} \partial_t \nu^*(t) + \text{div} \left(\mathbb{J}_{2d} \nabla_{\nu} \mathbb{H}(t, \nu^*(t), u^*(t)) \nu^*(t) \right) = 0, \\ \pi_{\#}^1 \nu^*(t) = \mu^*(t) & \text{for all times } t \in [0, T], \\ \nu^*(T) = \left(\text{Id}, -\lambda_0 \nabla \varphi(\mu^*(T)) - \sum_{i=1}^n \lambda_i \nabla \Psi_i(\mu^*(T)) \right)_{\#} \mu^*(T), \end{cases}$$

and

$$\mathbb{H}(t, \nu^*(t), u^*(t)) = \max_{u \in U} \mathbb{H}(t, \nu^*(t), u),$$

for a.e. $t \in [0, T]$, where the **Hamiltonian** is defined by

$$\mathbb{H}(t, \nu, u) := \int_{\mathbb{R}^{2d}} \langle r, v(t, \pi_{\#}^1 \nu, u, x) \rangle d\nu(x, r).$$

Mean-field OCP – PMP for smooth Mayer problems (1)

Pontryagin Maximum Principle for **(OCP)** [B., Frankowska '21]

Let $(\mu^*(\cdot), u^*(\cdot))$ be a strong local min. for **(OCP)**. Then, there exist **non-trivial multipliers** $(\lambda_0, \lambda_1, \dots, \lambda_n) \in \{0, 1\} \times \mathbb{R}_+^n$ and a **state-costate curve** $\nu^*(\cdot) \in \text{AC}([0, T], \mathcal{P}_c(\mathbb{R}^{2d}))$ s.t.

$$\begin{cases} \partial_t \nu^*(t) + \text{div} \left(\mathbb{J}_{2d} \nabla_{\nu} \mathbb{H}(t, \nu^*(t), u^*(t)) \nu^*(t) \right) = 0, \\ \pi_{\#}^1 \nu^*(t) = \mu^*(t) & \text{for all times } t \in [0, T], \\ \nu^*(T) = \left(\text{Id}, -\lambda_0 \nabla \varphi(\mu^*(T)) - \sum_{i=1}^n \lambda_i \nabla \Psi_i(\mu^*(T)) \right)_{\#} \mu^*(T), \end{cases}$$

and

$$\mathbb{H}(t, \nu^*(t), u^*(t)) = \max_{u \in U} \mathbb{H}(t, \nu^*(t), u),$$

for a.e. $t \in [0, T]$, where the **Hamiltonian** is defined by

$$\mathbb{H}(t, \nu, u) := \int_{\mathbb{R}^{2d}} \langle r, v(t, \pi_{\#}^1 \nu, u, x) \rangle d\nu(x, r).$$

Mean-field OCP – PMP for smooth Mayer problems (1)

Pontryagin Maximum Principle for **(OCP)** [B., Frankowska '21]

Let $(\mu^*(\cdot), u^*(\cdot))$ be a strong local min. for **(OCP)**. Then, there exist **non-trivial multipliers** $(\lambda_0, \lambda_1, \dots, \lambda_n) \in \{0, 1\} \times \mathbb{R}_+^n$ and a **state-costate curve** $\nu^*(\cdot) \in \text{AC}([0, T], \mathcal{P}_c(\mathbb{R}^{2d}))$ s.t.

$$\begin{cases} \partial_t \nu^*(t) + \text{div} \left(\mathbb{J}_{2d} \nabla_\nu \mathbb{H}(t, \nu^*(t), u^*(t)) \nu^*(t) \right) = 0, \\ \pi_{\#}^1 \nu^*(t) = \mu^*(t) & \text{for all times } t \in [0, T], \\ \nu^*(T) = \left(\text{Id}, -\lambda_0 \nabla \varphi(\mu^*(T)) - \sum_{i=1}^n \lambda_i \nabla \Psi_i(\mu^*(T)) \right)_{\#} \mu^*(T), \end{cases}$$

and

$$\mathbb{H}(t, \nu^*(t), u^*(t)) = \max_{u \in U} \mathbb{H}(t, \nu^*(t), u),$$

for a.e. $t \in [0, T]$, where the **Hamiltonian** is defined by

$$\mathbb{H}(t, \nu, u) := \int_{\mathbb{R}^{2d}} \langle r, v(t, \pi_{\#}^1 \nu, u, x) \rangle d\nu(x, r).$$

Mean-field OCP – PMP for smooth Mayer problems (1)

Pontryagin Maximum Principle for **(OCP)** [B., Frankowska '21]

Let $(\mu^*(\cdot), u^*(\cdot))$ be a strong local min. for **(OCP)**. Then, there exist **non-trivial multipliers** $(\lambda_0, \lambda_1, \dots, \lambda_n) \in \{0, 1\} \times \mathbb{R}_+^n$ and a **state-costate curve** $\nu^*(\cdot) \in \text{AC}([0, T], \mathcal{P}_c(\mathbb{R}^{2d}))$ s.t.

$$\begin{cases} \partial_t \nu^*(t) + \text{div} \left(\mathbb{J}_{2d} \nabla_{\nu} \mathbb{H}(t, \nu^*(t), u^*(t)) \nu^*(t) \right) = 0, \\ \pi_{\#}^1 \nu^*(t) = \mu^*(t) & \text{for all times } t \in [0, T], \\ \nu^*(T) = \left(\text{Id}, -\lambda_0 \nabla \varphi(\mu^*(T)) - \sum_{i=1}^n \lambda_i \nabla \Psi_i(\mu^*(T)) \right)_{\#} \mu^*(T), \end{cases}$$

and

$$\mathbb{H}(t, \nu^*(t), u^*(t)) = \max_{u \in U} \mathbb{H}(t, \nu^*(t), u),$$

for a.e. $t \in [0, T]$, where the **Hamiltonian** is defined by

$$\mathbb{H}(t, \nu, u) := \int_{\mathbb{R}^{2d}} \langle r, v(t, \pi_{\#}^1 \nu, u, x) \rangle d\nu(x, r).$$

Mean-field OCP – PMP for smooth Mayer problems (1)

Pontryagin Maximum Principle for **(OCP)** [B., Frankowska '21]

Let $(\mu^*(\cdot), u^*(\cdot))$ be a strong local min. for **(OCP)**. Then, there exist **non-trivial multipliers** $(\lambda_0, \lambda_1, \dots, \lambda_n) \in \{0, 1\} \times \mathbb{R}_+^n$ and a **state-costate curve** $\nu^*(\cdot) \in \text{AC}([0, T], \mathcal{P}_c(\mathbb{R}^{2d}))$ s.t.

$$\begin{cases} \partial_t \nu^*(t) + \text{div} \left(\mathbb{J}_{2d} \nabla_\nu \mathbb{H}(t, \nu^*(t), u^*(t)) \nu^*(t) \right) = 0, \\ \pi_{\#}^1 \nu^*(t) = \mu^*(t) & \text{for all times } t \in [0, T], \\ \nu^*(T) = \left(\text{Id}, -\lambda_0 \nabla \varphi(\mu^*(T)) - \sum_{i=1}^n \lambda_i \nabla \Psi_i(\mu^*(T)) \right)_{\#} \mu^*(T), \end{cases}$$

and

$$\mathbb{H}(t, \nu^*(t), u^*(t)) = \max_{u \in U} \mathbb{H}(t, \nu^*(t), u),$$

for a.e. $t \in [0, T]$, where the **Hamiltonian** is defined by

$$\mathbb{H}(t, \nu, u) := \int_{\mathbb{R}^{2d}} \langle r, v(t, \pi_{\#}^1 \nu, u, x) \rangle d\nu(x, r).$$

Mean-field OCP – PMP for smooth Mayer problems (1)

Pontryagin Maximum Principle for **(OCP)** [B., Frankowska '21]

Let $(\mu^*(\cdot), u^*(\cdot))$ be a strong local min. for **(OCP)**. Then, there exist **non-trivial multipliers** $(\lambda_0, \lambda_1, \dots, \lambda_n) \in \{0, 1\} \times \mathbb{R}_+^n$ and a **state-costate curve** $\nu^*(\cdot) \in \text{AC}([0, T], \mathcal{P}_c(\mathbb{R}^{2d}))$ s.t.

$$\begin{cases} \partial_t \nu^*(t) + \text{div} \left(\mathbb{J}_{2d} \nabla_\nu \mathbb{H}(t, \nu^*(t), u^*(t)) \nu^*(t) \right) = 0, \\ \pi_{\#}^1 \nu^*(t) = \mu^*(t) & \text{for all times } t \in [0, T], \\ \nu^*(T) = \left(\text{Id}, -\lambda_0 \nabla \varphi(\mu^*(T)) - \sum_{i=1}^n \lambda_i \nabla \Psi_i(\mu^*(T)) \right)_{\#} \mu^*(T), \end{cases}$$

and

$$\mathbb{H}(t, \nu^*(t), u^*(t)) = \max_{u \in U} \mathbb{H}(t, \nu^*(t), u),$$

for a.e. $t \in [0, T]$, where the **Hamiltonian** is defined by

$$\mathbb{H}(t, \nu, u) := \int_{\mathbb{R}^{2d}} \langle r, v(t, \pi_{\#}^1 \nu, u, x) \rangle d\nu(x, r).$$

Mean-field OCP – *PMP for smooth Mayer problems (2)*

Pontryagin Maximum Principle (Literature overview)

Previous derivations of the PMP:

- 1) by mean-field limits on **leader-follower** dynamics [BFRS'17],
- 2) on a particular problem inspired by fluids dynamics [P'16],
- 3) by **needle-variations** with closed-loop controls [BR'19],
- 4) by **packages of variations** for constrained problems [B'19],
- 5) for unconstrained **impulsive** problems [PS'20].

Only results for **constrained** Bolza problems \rightsquigarrow 4), but...

- Extremely technical proof (countable variations, limits, etc...)
- No discrimination between normal and abnormal cases

\rightsquigarrow Geometric approach with **set-valued** linearisations!

Mean-field OCP – PMP for smooth Mayer problems (2)

Pontryagin Maximum Principle (Literature overview)

Previous derivations of the PMP:

- 1) by mean-field limits on **leader-follower** dynamics [BFRS'17],
- 2) on a particular problem inspired by fluids dynamics [P'16],
- 3) by **needle-variations** with closed-loop controls [BR'19],
- 4) by **packages of variations** for constrained problems [B'19],
- 5) for unconstrained **impulsive** problems [PS'20].

Only results for **constrained** Bolza problems \rightsquigarrow 4), but...

- ◇ Extremely technical proof (countable variations, limits, etc...)
- ◇ No discrimination between normal and abnormal cases.

\rightsquigarrow Geometric approach with **set-valued linearisations**!

Mean-field OCP – PMP for smooth Mayer problems (2)

Pontryagin Maximum Principle (Literature overview)

Previous derivations of the PMP:

- 1) by mean-field limits on **leader-follower** dynamics [BFRS'17],
- 2) on a particular problem inspired by fluids dynamics [P'16],
- 3) by **needle-variations** with closed-loop controls [BR'19],
- 4) by **packages of variations** for constrained problems [B'19],
- 5) for unconstrained **impulsive** problems [PS'20].

Only results for **constrained** Bolza problems \rightsquigarrow 4), but...

- ◇ Extremely technical proof (countable variations, limits, etc...)
- ◇ No discrimination between normal and abnormal cases.

\rightsquigarrow Geometric approach with **set-valued linearisations**!

Mean-field OCP – PMP for smooth Mayer problems (2)

Pontryagin Maximum Principle (Literature overview)

Previous derivations of the PMP:

- 1) by mean-field limits on **leader-follower** dynamics [BFRS'17],
- 2) on a particular problem inspired by fluids dynamics [P'16],
- 3) by **needle-variations** with closed-loop controls [BR'19],
- 4) by **packages of variations** for constrained problems [B'19],
- 5) for unconstrained **impulsive** problems [PS'20].

Only results for **constrained** Bolza problems \rightsquigarrow 4), but...

- ◇ Extremely technical proof (countable variations, limits, etc...)
- ◇ No discrimination between normal and abnormal cases.

\rightsquigarrow Geometric approach with **set-valued linearisations**!

Mean-field OCP – PMP for smooth Mayer problems (2)

Pontryagin Maximum Principle (Literature overview)

Previous derivations of the PMP:

- 1) by mean-field limits on **leader-follower** dynamics [BFRS'17],
- 2) on a particular problem inspired by fluids dynamics [P'16],
- 3) by **needle-variations** with closed-loop controls [BR'19],
- 4) by **packages of variations** for constrained problems [B'19],
- 5) for unconstrained **impulsive** problems [PS'20].

Only results for **constrained** Bolza problems \rightsquigarrow 4), but...

- ◇ Extremely technical proof (countable variations, limits, etc...)
- ◇ No discrimination between normal and abnormal cases.

\rightsquigarrow Geometric approach with **set-valued** linearisations!

Mean-field OCP – PMP for smooth Mayer problems (2)

Pontryagin Maximum Principle (Literature overview)

Previous derivations of the PMP:

- 1) by mean-field limits on **leader-follower** dynamics [BFRS'17],
- 2) on a particular problem inspired by fluids dynamics [P'16],
- 3) by **needle-variations** with closed-loop controls [BR'19],
- 4) by **packages of variations** for constrained problems [B'19],
- 5) for unconstrained **impulsive** problems [PS'20].

Only results for **constrained** Bolza problems \rightsquigarrow 4), but...

- ◇ Extremely technical proof (countable variations, limits, etc...)
- ◇ No discrimination between normal and abnormal cases.

\rightsquigarrow Geometric approach with **set-valued linearisations**!

Mean-field OCP – PMP for smooth Mayer problems (2)

Pontryagin Maximum Principle (Literature overview)

Previous derivations of the PMP:

- 1) by mean-field limits on **leader-follower** dynamics [BFRS'17],
- 2) on a particular problem inspired by fluids dynamics [P'16],
- 3) by **needle-variations** with closed-loop controls [BR'19],
- 4) by **packages of variations** for constrained problems [B'19],
- 5) for unconstrained **impulsive** problems [PS'20].

Only results for **constrained** Bolza problems \rightsquigarrow 4), but...

- ◇ Extremely technical proof (countable variations, limits, etc...)
- ◇ No discrimination between normal and abnormal cases.

\rightsquigarrow Geometric approach with **set-valued** linearisations!

Mean-field OCP – PMP for smooth Mayer problems (2)

Pontryagin Maximum Principle (Literature overview)

Previous derivations of the PMP:

- 1) by mean-field limits on **leader-follower** dynamics [BFRS'17],
- 2) on a particular problem inspired by fluids dynamics [P'16],
- 3) by **needle-variations** with closed-loop controls [BR'19],
- 4) by **packages of variations** for constrained problems [B'19],
- 5) for unconstrained **impulsive** problems [PS'20].

Only results for **constrained** Bolza problems \rightsquigarrow 4), but...

- ◇ Extremely technical proof (countable variations, limits, etc...)
- ◇ No discrimination between normal and abnormal cases.

\rightsquigarrow Geometric approach with **set-valued** linearisations!

Mean-field OCP – PMP for smooth Mayer problems (2)

Pontryagin Maximum Principle (Literature overview)

Previous derivations of the PMP:

- 1) by mean-field limits on **leader-follower** dynamics [BFRS'17],
- 2) on a particular problem inspired by fluids dynamics [P'16],
- 3) by **needle-variations** with closed-loop controls [BR'19],
- 4) by **packages of variations** for constrained problems [B'19],
- 5) for unconstrained **impulsive** problems [PS'20].

Only results for **constrained** Bolza problems \rightsquigarrow 4), but...

- ◇ Extremely technical proof (countable variations, limits, etc...)
- ◇ No discrimination between normal and abnormal cases.

\rightsquigarrow Geometric approach with **set-valued** linearisations!

Mean-field OCP – PMP for smooth Mayer problems (2)

Pontryagin Maximum Principle (Literature overview)

Previous derivations of the PMP:

- 1) by mean-field limits on **leader-follower** dynamics [BFRS'17],
- 2) on a particular problem inspired by fluids dynamics [P'16],
- 3) by **needle-variations** with closed-loop controls [BR'19],
- 4) by **packages of variations** for constrained problems [B'19],
- 5) for unconstrained **impulsive** problems [PS'20].

Only results for **constrained** Bolza problems \rightsquigarrow 4), but...

- ◇ Extremely technical proof (countable variations, limits, etc...)
- ◇ No discrimination between normal and abnormal cases.

\rightsquigarrow Geometric approach with **set-valued linearisations**!

Mean-field OCP – Proof of the PMP

Proof of the PMP (Main steps)

- 1) Reformulate the dynamics as (CI) with $V(t, \mu) := v(t, \mu, U)$.
- 2) For all $\epsilon > 0$ and $t \mapsto \mathbf{w}(t) \in T_{\text{co}V(t, \mu^*(t))}(v(t, \mu^*(t), u^*(t)))$, there exists $\tilde{\mu}_\epsilon(\cdot)$ sol. of (CI) s.t.

$$\sup_{t \in [0, T]} W_1\left(\tilde{\mu}_\epsilon(t), (\text{Id} + \epsilon \mathcal{F}_{\mathbf{w}}(t, \cdot))_{\#} \mu^*(t)\right) = o(\epsilon),$$

where $t \mapsto \mathcal{F}_{\mathbf{w}}(t, x) \in \mathbb{R}^d$ “solves” a lin. system [B’19, BR’19].

- 3) Contradiction \oplus Separation $\implies \exists (\lambda_0, \lambda_1, \dots, \lambda_n) \neq 0$ s.t.

$$\left\langle \lambda_0 \nabla \varphi(\mu^*(T)) + \sum_{i=1}^n \lambda_i \nabla \Psi_i(\mu^*(T)), \mathcal{F}_{\mathbf{w}}(T) \right\rangle_{L^2(\mu^*(T))} \geq 0,$$

for every $t \mapsto \mathbf{w}(t) \in T_{\text{co}V(t, \mu^*(t))}(v(t, \mu^*(t), u^*(t)))$.

- 4) Backward prop. along $\nu^*(\cdot) \rightsquigarrow$ **maximisation condition!**

Mean-field OCP – Proof of the PMP

Proof of the PMP (Main steps)

- 1) Reformulate the dynamics as (CI) with $V(t, \mu) := v(t, \mu, U)$.
- 2) For all $\epsilon > 0$ and $t \mapsto \mathbf{w}(t) \in T_{\text{co}V(t, \mu^*(t))}(v(t, \mu^*(t), \mathbf{u}^*(t)))$, there exists $\tilde{\mu}_\epsilon(\cdot)$ sol. of (CI) s.t.

$$\sup_{t \in [0, T]} W_1\left(\tilde{\mu}_\epsilon(t), (\text{Id} + \epsilon \mathcal{F}_{\mathbf{w}}(t, \cdot))_{\#} \mu^*(t)\right) = o(\epsilon),$$

where $t \mapsto \mathcal{F}_{\mathbf{w}}(t, x) \in \mathbb{R}^d$ “solves” a lin. system [B’19, BR’19].

- 3) Contradiction \oplus Separation $\implies \exists (\lambda_0, \lambda_1, \dots, \lambda_n) \neq 0$ s.t.

$$\left\langle \lambda_0 \nabla \varphi(\mu^*(T)) + \sum_{i=1}^n \lambda_i \nabla \Psi_i(\mu^*(T)), \mathcal{F}_{\mathbf{w}}(T) \right\rangle_{L^2(\mu^*(T))} \geq 0,$$

for every $t \mapsto \mathbf{w}(t) \in T_{\text{co}V(t, \mu^*(t))}(v(t, \mu^*(t), \mathbf{u}^*(t)))$.

- 4) Backward prop. along $\nu^*(\cdot) \rightsquigarrow$ **maximisation condition!**

Mean-field OCP – Proof of the PMP

Proof of the PMP (Main steps)

- 1) Reformulate the dynamics as (CI) with $V(t, \mu) := v(t, \mu, U)$.
- 2) For all $\epsilon > 0$ and $t \mapsto \mathbf{w}(t) \in T_{\text{co}V(t, \mu^*(t))}(v(t, \mu^*(t), \mathbf{u}^*(t)))$, there exists $\tilde{\mu}_\epsilon(\cdot)$ sol. of (CI) s.t.

$$\sup_{t \in [0, T]} W_1\left(\tilde{\mu}_\epsilon(t), (\text{Id} + \epsilon \mathcal{F}_{\mathbf{w}}(t, \cdot))_{\#} \mu^*(t)\right) = o(\epsilon),$$

where $t \mapsto \mathcal{F}_{\mathbf{w}}(t, x) \in \mathbb{R}^d$ “solves” a lin. system [B’19, BR’19].

- 3) Contradiction \oplus Separation $\implies \exists (\lambda_0, \lambda_1, \dots, \lambda_n) \neq 0$ s.t.

$$\left\langle \lambda_0 \nabla \varphi(\mu^*(T)) + \sum_{i=1}^n \lambda_i \nabla \Psi_i(\mu^*(T)), \mathcal{F}_{\mathbf{w}}(T) \right\rangle_{L^2(\mu^*(T))} \geq 0,$$

for every $t \mapsto \mathbf{w}(t) \in T_{\text{co}V(t, \mu^*(t))}(v(t, \mu^*(t), \mathbf{u}^*(t)))$.

- 4) Backward prop. along $\nu^*(\cdot) \rightsquigarrow$ **maximisation condition!**

Mean-field OCP – Proof of the PMP

Proof of the PMP (Main steps)

- 1) Reformulate the dynamics as (CI) with $V(t, \mu) := v(t, \mu, U)$.
- 2) For all $\epsilon > 0$ and $t \mapsto \mathbf{w}(t) \in T_{\text{co}V(t, \mu^*(t))}(v(t, \mu^*(t), \mathbf{u}^*(t)))$, there exists $\tilde{\mu}_\epsilon(\cdot)$ sol. of (CI) s.t.

$$\sup_{t \in [0, T]} W_1\left(\tilde{\mu}_\epsilon(t), (\text{Id} + \epsilon \mathcal{F}_{\mathbf{w}}(t, \cdot))_{\#} \mu^*(t)\right) = o(\epsilon),$$

where $t \mapsto \mathcal{F}_{\mathbf{w}}(t, x) \in \mathbb{R}^d$ “solves” a lin. system [B’19, BR’19].

- 3) Contradiction \oplus Separation $\implies \exists (\lambda_0, \lambda_1, \dots, \lambda_n) \neq 0$ s.t.

$$\left\langle \lambda_0 \nabla \varphi(\mu^*(T)) + \sum_{i=1}^n \lambda_i \nabla \Psi_i(\mu^*(T)), \mathcal{F}_{\mathbf{w}}(T) \right\rangle_{L^2(\mu^*(T))} \geq 0,$$

for every $t \mapsto \mathbf{w}(t) \in T_{\text{co}V(t, \mu^*(t))}(v(t, \mu^*(t), \mathbf{u}^*(t)))$.

- 4) Backward prop. along $\nu^*(\cdot) \rightsquigarrow$ **maximisation condition!**

Mean-field OCP – Proof of the PMP

Proof of the PMP (Main steps)

- 1) Reformulate the dynamics as (CI) with $V(t, \mu) := v(t, \mu, U)$.
- 2) For all $\epsilon > 0$ and $t \mapsto \mathbf{w}(t) \in T_{\text{co}V(t, \mu^*(t))}(v(t, \mu^*(t), \mathbf{u}^*(t)))$, there exists $\tilde{\mu}_\epsilon(\cdot)$ sol. of (CI) s.t.

$$\sup_{t \in [0, T]} W_1\left(\tilde{\mu}_\epsilon(t), (\text{Id} + \epsilon \mathcal{F}_{\mathbf{w}}(t, \cdot))_{\#} \mu^*(t)\right) = o(\epsilon),$$

where $t \mapsto \mathcal{F}_{\mathbf{w}}(t, x) \in \mathbb{R}^d$ “solves” a lin. system [B’19, BR’19].

- 3) Contradiction \oplus Separation $\implies \exists (\lambda_0, \lambda_1, \dots, \lambda_n) \neq 0$ s.t.

$$\left\langle \lambda_0 \nabla \varphi(\mu^*(T)) + \sum_{i=1}^n \lambda_i \nabla \Psi_i(\mu^*(T)), \mathcal{F}_{\mathbf{w}}(T) \right\rangle_{L^2(\mu^*(T))} \geq 0,$$

for every $t \mapsto \mathbf{w}(t) \in T_{\text{co}V(t, \mu^*(t))}(v(t, \mu^*(t), \mathbf{u}^*(t)))$.

- 4) Backward prop. along $\nu^*(\cdot) \rightsquigarrow$ **maximisation condition!**

Mean-field OCP – Proof of the PMP

Proof of the PMP (Main steps)

- 1) Reformulate the dynamics as (CI) with $V(t, \mu) := v(t, \mu, U)$.
- 2) For all $\epsilon > 0$ and $t \mapsto \mathbf{w}(t) \in T_{\text{co}V(t, \mu^*(t))}(v(t, \mu^*(t), \mathbf{u}^*(t)))$, there exists $\tilde{\mu}_\epsilon(\cdot)$ sol. of (CI) s.t.

$$\sup_{t \in [0, T]} W_1\left(\tilde{\mu}_\epsilon(t), (\text{Id} + \epsilon \mathcal{F}_{\mathbf{w}}(t, \cdot))_{\#} \mu^*(t)\right) = o(\epsilon),$$

where $t \mapsto \mathcal{F}_{\mathbf{w}}(t, x) \in \mathbb{R}^d$ “solves” a lin. system [B’19, BR’19].

- 3) Contradiction \oplus Separation $\implies \exists (\lambda_0, \lambda_1, \dots, \lambda_n) \neq 0$ s.t.

$$\left\langle \lambda_0 \nabla \varphi(\mu^*(T)) + \sum_{i=1}^n \lambda_i \nabla \Psi_i(\mu^*(T)), \mathcal{F}_{\mathbf{w}}(T) \right\rangle_{L^2(\mu^*(T))} \geq 0,$$

for every $t \mapsto \mathbf{w}(t) \in T_{\text{co}V(t, \mu^*(t))}(v(t, \mu^*(t), \mathbf{u}^*(t)))$.

- 4) Backward prop. along $\nu^*(\cdot) \rightsquigarrow$ **maximisation condition!**

Mean-field OCP – Proof of the PMP

Proof of the PMP (Main steps)

- 1) Reformulate the dynamics as (CI) with $V(t, \mu) := v(t, \mu, U)$.
- 2) For all $\epsilon > 0$ and $t \mapsto \mathbf{w}(t) \in T_{\text{co}V(t, \mu^*(t))}(v(t, \mu^*(t), \mathbf{u}^*(t)))$, there exists $\tilde{\mu}_\epsilon(\cdot)$ sol. of (CI) s.t.

$$\sup_{t \in [0, T]} W_1\left(\tilde{\mu}_\epsilon(t), (\text{Id} + \epsilon \mathcal{F}_{\mathbf{w}}(t, \cdot))_{\#} \mu^*(t)\right) = o(\epsilon),$$

where $t \mapsto \mathcal{F}_{\mathbf{w}}(t, x) \in \mathbb{R}^d$ “solves” a lin. system [B’19, BR’19].

- 3) Contradiction \oplus Separation $\implies \exists (\lambda_0, \lambda_1, \dots, \lambda_n) \neq 0$ s.t.

$$\left\langle \lambda_0 \nabla \varphi(\mu^*(T)) + \sum_{i=1}^n \lambda_i \nabla \Psi_i(\mu^*(T)), \mathcal{F}_{\mathbf{w}}(T) \right\rangle_{L^2(\mu^*(T))} \geq 0,$$

for every $t \mapsto \mathbf{w}(t) \in T_{\text{co}V(t, \mu^*(t))}(v(t, \mu^*(t), \mathbf{u}^*(t)))$.

- 4) **Backward** prop. along $\nu^*(\cdot) \rightsquigarrow$ **maximisation condition!**

Mean-field OCP – Proof of the PMP

Proof of the PMP (Main steps)

- 1) Reformulate the dynamics as (CI) with $V(t, \mu) := v(t, \mu, U)$.
- 2) For all $\epsilon > 0$ and $t \mapsto \mathbf{w}(t) \in T_{\text{co}V(t, \mu^*(t))}(v(t, \mu^*(t), \mathbf{u}^*(t)))$, there exists $\tilde{\mu}_\epsilon(\cdot)$ sol. of (CI) s.t.

$$\sup_{t \in [0, T]} W_1\left(\tilde{\mu}_\epsilon(t), (\text{Id} + \epsilon \mathcal{F}_{\mathbf{w}}(t, \cdot))_{\#} \mu^*(t)\right) = o(\epsilon),$$

where $t \mapsto \mathcal{F}_{\mathbf{w}}(t, x) \in \mathbb{R}^d$ “solves” a lin. system [B’19, BR’19].

- 3) Contradiction \oplus Separation $\implies \exists (\lambda_0, \lambda_1, \dots, \lambda_n) \neq 0$ s.t.

$$\left\langle \lambda_0 \nabla \varphi(\mu^*(T)) + \sum_{i=1}^n \lambda_i \nabla \Psi_i(\mu^*(T)), \mathcal{F}_{\mathbf{w}}(T) \right\rangle_{L^2(\mu^*(T))} \geq 0,$$

for every $t \mapsto \mathbf{w}(t) \in T_{\text{co}V(t, \mu^*(t))}(v(t, \mu^*(t), \mathbf{u}^*(t)))$.

- 4) **Backward** prop. along $\nu^*(\cdot) \rightsquigarrow$ **maximisation condition!**

Outline of the talk

Mean-field control problems

From differential to continuity inclusions

Main structural results in the Cauchy-Lipschitz case

Application : geometric derivation of the PMP

Conclusion

Conclusion – *Recap' and open perspectives*

Recap' of the presentation

- ◇ **Geometric** notion of **set-valued** dynamics for continuity equations \rightsquigarrow **Continuity inclusions** [B., Frankowska JDE'21]
- ◇ **Localised** differentiability and **variational** linearisations \rightsquigarrow **Pontryagin Principle** [B., Frankowska AMO'21 (soon)]

Work (in progress and to come)

- 1) **Semiconcavity** and **sensitivity** value function \rightsquigarrow submitted !
- 2) General **viability** theorems \rightsquigarrow in progress
- 3) PMP for problems with **general constraints** & second-order **necessary and sufficient conditions** \rightsquigarrow to do later...

Conclusion – *Recap' and open perspectives*

Recap' of the presentation

- ◇ **Geometric** notion of **set-valued** dynamics for continuity equations \rightsquigarrow **Continuity inclusions** [B., Frankowska JDE'21]
- ◇ **Localised** differentiability and **variational** linearisations \rightsquigarrow **Pontryagin Principle** [B., Frankowska AMO'21 (soon)]

Work (in progress and to come)

- 1) **Semiconcavity** and **sensitivity** value function \rightsquigarrow submitted !
- 2) General **viability** theorems \rightsquigarrow in progress
- 3) PMP for problems with **general constraints** & second-order **necessary and sufficient conditions** \rightsquigarrow to do later...

Conclusion – *Recap' and open perspectives*

Recap' of the presentation

- ◇ **Geometric** notion of **set-valued** dynamics for continuity equations \rightsquigarrow **Continuity inclusions** [B., Frankowska JDE'21]
- ◇ **Localised** differentiability and **variational** linearisations \rightsquigarrow **Pontryagin Principle** [B., Frankowska AMO'21 (soon)]

Work (in progress and to come)

- 1) **Semiconcavity** and **sensitivity** value function \rightsquigarrow submitted !
- 2) General **viability** theorems \rightsquigarrow in progress
- 3) PMP for problems with **general constraints** & second-order **necessary** and **sufficient** conditions \rightsquigarrow to do later...

Conclusion – *Recap' and open perspectives*

Recap' of the presentation

- ◇ **Geometric** notion of **set-valued** dynamics for continuity equations \rightsquigarrow **Continuity inclusions** [B., Frankowska JDE'21]
- ◇ **Localised** differentiability and **variational** linearisations \rightsquigarrow **Pontryagin Principle** [B., Frankowska AMO'21 (soon)]

Work (in progress and to come)

- 1) **Semiconcavity** and **sensitivity** value function \rightsquigarrow submitted !
- 2) General **viability** theorems \rightsquigarrow in progress
- 3) PMP for problems with **general constraints** & second-order **necessary** and **sufficient** conditions \rightsquigarrow to do later...

Conclusion – *Recap' and open perspectives*

Recap' of the presentation

- ◇ **Geometric** notion of **set-valued** dynamics for continuity equations \rightsquigarrow **Continuity inclusions** [B., Frankowska JDE'21]
- ◇ **Localised** differentiability and **variational** linearisations \rightsquigarrow **Pontryagin Principle** [B., Frankowska AMO'21 (soon)]

Work (in progress and to come)

- 1) **Semiconcavity** and **sensitivity** value function \rightsquigarrow submitted !
- 2) General **viability** theorems \rightsquigarrow in progress
- 3) PMP for problems with **general constraints** & second-order **necessary** and **sufficient** conditions \rightsquigarrow to do later...

Conclusion – *Recap' and open perspectives*

Recap' of the presentation

- ◇ **Geometric** notion of **set-valued** dynamics for continuity equations \rightsquigarrow **Continuity inclusions** [B., Frankowska JDE'21]
- ◇ **Localised** differentiability and **variational** linearisations \rightsquigarrow **Pontryagin Principle** [B., Frankowska AMO'21 (soon)]

Work (in progress and to come)

- 1) **Semiconcavity** and **sensitivity** value function \rightsquigarrow submitted !
- 2) General **viability** theorems \rightsquigarrow in progress
- 3) PMP for problems with **general constraints** & second-order **necessary** and **sufficient** conditions \rightsquigarrow to do later...

Conclusion – *That's all folks!*

Thank you for your attention !

