

# Optimal Control Problems in Wasserstein Spaces

## *A PDE Approach to Multi-Agent Systems*

Benoît Bonnet

LIS, Aix-Marseille Université, FRANCE

*LabEx Archimède*

04/07/2018

**14th Viennese Conference on Optimal Control and  
Dynamic Games**

# Outline of the talk

- 1 Introduction and Motivations
- 2 Optimal Transport Theory & Wasserstein Spaces
- 3 Optimal Control Problems and Pontryagin Maximum Principle

# Outline of the talk


- 1 Introduction and Motivations
- 2 Optimal Transport Theory & Wasserstein Spaces
- 3 Optimal Control Problems and Pontryagin Maximum Principle

**Idea :** Understand and control the global behaviour of large systems of **interacting** agents




# Introduction and motivation - *Multi-Agent Systems*

**Idea** : Understand and control the global behaviour of large systems of **interacting** agents



# Introduction and motivation - *Multi-Agent Systems*

**Idea :** Understand and control the global behaviour of large systems of **interacting** agents



# Introduction and motivation - *Multi-Agent Systems*

**Idea :** Understand and control the global behaviour of large systems of **interacting** agents



## Example (Finite dim. Drift + Convolution + control)

Consider  $N$  agents  $(x_1, \dots, x_N) \in (\mathbb{R}^d)^N$  evolving according to

$$\dot{x}_i(t) = v_d(x_i) + \frac{1}{N} \sum_{j=1}^N K(x_i - x_j) + u_i(t), \quad (S_N)$$

**Objective :** design a set of control laws  $t \mapsto (u_1(t), \dots, u_N(t))$  achieving a certain goal, e.g. forming a consensus, etc...

## Issues related to the formulation $(S_N)$

- ◇ No a priori knowledge of the system (number of agents, exact positions, etc...)
- ◇ Not so relevant to consider a discrete model for very large systems of interacting agents
- ◇ Extremely demanding computationally speaking



## Example (Finite dim. Drift + Convolution + control)

Consider  $N$  agents  $(x_1, \dots, x_N) \in (\mathbb{R}^d)^N$  evolving according to

$$\dot{x}_i(t) = v_d(x_i) + \frac{1}{N} \sum_{j=1}^N K(x_i - x_j) + u_i(t), \quad (S_N)$$

**Objective :** design a set of control laws  $t \mapsto (u_1(t), \dots, u_N(t))$  achieving a certain goal, e.g. forming a consensus, etc...

## Issues related to the formulation $(S_N)$

- ◇ No a priori knowledge of the system (number of agents, exact positions, etc...)
- ◇ Not so relevant to consider a discrete model for very large systems of interacting agents
- ◇ Extremely demanding computationally speaking

## Example (Finite dim. Drift + Convolution + control)

Consider  $N$  agents  $(x_1, \dots, x_N) \in (\mathbb{R}^d)^N$  evolving according to

$$\dot{x}_i(t) = v_d(x_i) + \frac{1}{N} \sum_{j=1}^N K(x_i - x_j) + u_i(t), \quad (S_N)$$

**Objective :** design a set of control laws  $t \mapsto (u_1(t), \dots, u_N(t))$  achieving a certain goal, e.g. forming a consensus, etc...

## Issues related to the formulation $(S_N)$

- ◇ No a priori knowledge of the system (number of agents, exact positions, etc...)
- ◇ Not so relevant to consider a discrete model for very large systems of interacting agents
- ◇ Extremely demanding computationally speaking

## Example (Finite dim. Drift + Convolution + control)

Consider  $N$  agents  $(x_1, \dots, x_N) \in (\mathbb{R}^d)^N$  evolving according to

$$\dot{x}_i(t) = v_d(x_i) + \frac{1}{N} \sum_{j=1}^N K(x_i - x_j) + u_i(t), \quad (S_N)$$

**Objective :** design a set of control laws  $t \mapsto (u_1(t), \dots, u_N(t))$  achieving a certain goal, e.g. forming a consensus, etc...

## Issues related to the formulation $(S_N)$

- ◇ No a priori knowledge of the system (number of agents, exact positions, etc...)
- ◇ Not so relevant to consider a discrete model for very large systems of interacting agents
- ◇ Extremely demanding computationally speaking

## Example (Finite dim. Drift + Convolution + control)

Consider  $N$  agents  $(x_1, \dots, x_N) \in (\mathbb{R}^d)^N$  evolving according to

$$\dot{x}_i(t) = v_d(x_i) + \frac{1}{N} \sum_{j=1}^N K(x_i - x_j) + u_i(t), \quad (S_N)$$


**Objective :** design a set of control laws  $t \mapsto (u_1(t), \dots, u_N(t))$  achieving a certain goal, e.g. forming a consensus, etc...

## Issues related to the formulation $(S_N)$

- ◇ No a priori knowledge of the system (number of agents, exact positions, etc...)
- ◇ Not so relevant to consider a discrete model for very large systems of interacting agents
- ◇ Extremely demanding computationally speaking

# Introduction and motivation - *Mean-field approximation*


**Idea :** Approximate the large crowd by a single PDE through a *mean-field process*



$N$  ODEs on the agents  $(x_1, \dots, x_N) \rightsquigarrow 1$  PDE on the density  $\mu$

# Introduction and motivation - *Mean-field approximation*

**Idea :** Approximate the large crowd by a single PDE through a *mean-field process*



$N$  ODEs on the agents  $(x_1, \dots, x_N) \rightsquigarrow 1$  PDE on the density  $\mu$

**Mean field controlled system** described by *transport equations with non-local velocities*

$$\partial_t \mu(t) + \nabla \cdot ((v[\mu(t)](\cdot) + u(t, \cdot))\mu(t)) = 0. \quad (S_\infty)$$

where the controls *depend on time and space*.

Question : Choice of the state space ?

Discrete  $\oplus$  continuous objects  $\rightsquigarrow$  '*distributional spaces*'

**Problem** : Distributional topologies are not very nice...

$\hookrightarrow$  Space of probability measures *endowed with an optimal transportation metric* !

**Mean field controlled system** described by *transport equations with non-local velocities*

$$\partial_t \mu(t) + \nabla \cdot ((v[\mu(t)](\cdot) + u(t, \cdot))\mu(t)) = 0. \quad (S_\infty)$$

where the controls *depend on time and space*.

**Question :** Choice of the state space ?

Discrete  $\oplus$  continuous objects  $\rightsquigarrow$  '*distributional spaces*'

**Problem :** Distributional topologies are not very nice...

$\hookrightarrow$  Space of probability measures *endowed with an optimal transportation metric !*



**Mean field controlled system** described by *transport equations with non-local velocities*

$$\partial_t \mu(t) + \nabla \cdot ((v[\mu(t)](\cdot) + u(t, \cdot))\mu(t)) = 0. \quad (S_\infty)$$

where the controls *depend on time and space*.

**Question :** Choice of the state space ?

Discrete  $\oplus$  continuous objects  $\rightsquigarrow$  '*distributional spaces*'

**Problem :** Distributional topologies are not very nice...

$\hookrightarrow$  Space of probability measures *endowed with an optimal transportation metric !*

**Mean field controlled system** described by *transport equations with non-local velocities*

$$\partial_t \mu(t) + \nabla \cdot ((v[\mu(t)](\cdot) + u(t, \cdot))\mu(t)) = 0. \quad (S_\infty)$$

where the controls *depend on time and space*.

**Question :** Choice of the state space ?

Discrete  $\oplus$  continuous objects  $\rightsquigarrow$  '*distributional spaces*'

**Problem :** Distributional topologies are not very nice...

$\hookrightarrow$  Space of probability measures *endowed with an optimal transportation metric !*

**Mean field controlled system** described by *transport equations with non-local velocities*

$$\partial_t \mu(t) + \nabla \cdot ((v[\mu(t)](\cdot) + u(t, \cdot))\mu(t)) = 0. \quad (S_\infty)$$

where the controls *depend on time and space*.

**Question :** Choice of the state space ?

Discrete  $\oplus$  continuous objects  $\rightsquigarrow$  '*distributional spaces*'

**Problem :** Distributional topologies are not very nice...

$\hookrightarrow$  Space of probability measures *endowed with an optimal transportation metric !*

**Mean field controlled system** described by *transport equations with non-local velocities*

$$\partial_t \mu(t) + \nabla \cdot ((v[\mu(t)](\cdot) + u(t, \cdot))\mu(t)) = 0. \quad (S_\infty)$$

where the controls *depend on time and space*.

**Question :** Choice of the state space ?

Discrete  $\oplus$  continuous objects  $\rightsquigarrow$  '*distributional spaces*'

**Problem :** Distributional topologies are not very nice...

$\hookrightarrow$  Space of probability measures *endowed with an optimal transportation metric !*

# Outline of the talk


- 1 Introduction and Motivations
- 2 Optimal Transport Theory & Wasserstein Spaces
- 3 Optimal Control Problems and Pontryagin Maximum Principle

## Kantorovich problem (1942) - $(OT_K)$

Given two probability measures  $\mu, \nu \in \mathcal{P}_c(\mathbb{R}^d)$  and a cost function  $c : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ , **find a measure**  $\gamma \in \mathcal{P}_c(\mathbb{R}^{2d})$  such that

$$(i) \quad \gamma \in \Gamma(\mu, \nu) = \left\{ \gamma' \in \mathcal{P}_c(\mathbb{R}^{2d}) \text{ s.t. } \pi_{\#}^1 \gamma' = \mu, \pi_{\#}^2 \gamma' = \nu \right\},$$

$$(ii) \quad \int_{\mathbb{R}^{2d}} c(x, y) d\gamma(x, y) = \min_{\gamma' \in \Gamma(\mu, \nu)} \left[ \int_{\mathbb{R}^{2d}} c(x, y) d\gamma'(x, y) \right].$$




## Kantorovich problem (1942) - ( $\text{OT}_K$ )

Given two probability measures  $\mu, \nu \in \mathcal{P}_c(\mathbb{R}^d)$  and a cost function  $c : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ , **find a measure**  $\gamma \in \mathcal{P}_c(\mathbb{R}^{2d})$  such that

$$(i) \quad \gamma \in \Gamma(\mu, \nu) = \left\{ \gamma' \in \mathcal{P}_c(\mathbb{R}^{2d}) \text{ s.t. } \pi_{\#}^1 \gamma' = \mu, \pi_{\#}^2 \gamma' = \nu \right\},$$

$$(ii) \quad \int_{\mathbb{R}^{2d}} c(x, y) d\gamma(x, y) = \min_{\gamma' \in \Gamma(\mu, \nu)} \left[ \int_{\mathbb{R}^{2d}} c(x, y) d\gamma'(x, y) \right].$$




## Kantorovich problem (1942) - $(OT_K)$

Given two probability measures  $\mu, \nu \in \mathcal{P}_c(\mathbb{R}^d)$  and a cost function  $c : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ , **find a measure**  $\gamma \in \mathcal{P}_c(\mathbb{R}^{2d})$  such that

$$(i) \quad \gamma \in \Gamma(\mu, \nu) = \left\{ \gamma' \in \mathcal{P}_c(\mathbb{R}^{2d}) \text{ s.t. } \pi_{\#}^1 \gamma' = \mu, \pi_{\#}^2 \gamma' = \nu \right\},$$

$$(ii) \quad \int_{\mathbb{R}^{2d}} c(x, y) d\gamma(x, y) = \min_{\gamma' \in \Gamma(\mu, \nu)} \left[ \int_{\mathbb{R}^{2d}} c(x, y) d\gamma'(x, y) \right].$$






## Kantorovich problem (1942) - $(OT_K)$

Given two probability measures  $\mu, \nu \in \mathcal{P}_c(\mathbb{R}^d)$  and a cost function  $c : \mathbb{R}^d \times \mathbb{R}^d \rightarrow \mathbb{R}$ , **find a measure**  $\gamma \in \mathcal{P}_c(\mathbb{R}^{2d})$  such that

$$(i) \quad \gamma \in \Gamma(\mu, \nu) = \left\{ \gamma' \in \mathcal{P}_c(\mathbb{R}^{2d}) \text{ s.t. } \pi_{\#}^1 \gamma' = \mu, \pi_{\#}^2 \gamma' = \nu \right\},$$

$$(ii) \quad \int_{\mathbb{R}^{2d}} c(x, y) d\gamma(x, y) = \min_{\gamma' \in \Gamma(\mu, \nu)} \left[ \int_{\mathbb{R}^{2d}} c(x, y) d\gamma'(x, y) \right].$$



## Definition (Wasserstein distance)

Taking  $c(x, y) = |x - y|^2$ , the quantity

$$W_2(\mu, \nu) = \min_{\gamma \in \Gamma(\mu, \nu)} \left\{ \left( \int_{\mathbb{R}^{2d}} |x - y|^2 d\gamma(x, y) \right)^{1/2} \right\}$$

defines a *distance* over  $\mathcal{P}_c(\mathbb{R}^d)$ .

*~ Infimum of  $L^2$ -distances over the couplings  $\gamma \in \Gamma(\mu, \nu)$ .*

## Proposition (Some useful properties)

- ◇  $W_2$  metrizes the usual weak- $*$  topology of measures.*
- ◇  $(\mathcal{P}_c(\mathbb{R}^d), W_2)$  is a complete and separable metric space.*
- ◇  $(\mathcal{P}_c(\mathbb{R}^d), W_2)$  can be endowed with a differential structure*

## Definition (Wasserstein distance)

Taking  $c(x, y) = |x - y|^2$ , the quantity

$$W_2(\mu, \nu) = \min_{\gamma \in \Gamma(\mu, \nu)} \left\{ \left( \int_{\mathbb{R}^{2d}} |x - y|^2 d\gamma(x, y) \right)^{1/2} \right\}$$

defines a *distance* over  $\mathcal{P}_c(\mathbb{R}^d)$ .

$\sim$  Infimum of  $L^2$ -distances over the couplings  $\gamma \in \Gamma(\mu, \nu)$ .

## Proposition (Some useful properties)

- $\diamond$   $W_2$  metrizes the usual weak-\* topology of measures.
- $\diamond$   $(\mathcal{P}_c(\mathbb{R}^d), W_2)$  is a complete and separable metric space.
- $\diamond$   $(\mathcal{P}_c(\mathbb{R}^d), W_2)$  can be endowed with a differential structure

## Definition (Wasserstein distance)

Taking  $c(x, y) = |x - y|^2$ , the quantity

$$W_2(\mu, \nu) = \min_{\gamma \in \Gamma(\mu, \nu)} \left\{ \left( \int_{\mathbb{R}^{2d}} |x - y|^2 d\gamma(x, y) \right)^{1/2} \right\}$$

defines a *distance* over  $\mathcal{P}_c(\mathbb{R}^d)$ .

$\sim$  Infimum of  $L^2$ -distances over the couplings  $\gamma \in \Gamma(\mu, \nu)$ .

## Proposition (Some useful properties)

- ◇  $W_2$  metrizes the usual weak-\* topology of measures.
- ◇  $(\mathcal{P}_c(\mathbb{R}^d), W_2)$  is a complete and separable metric space.
- ◇  $(\mathcal{P}_c(\mathbb{R}^d), W_2)$  can be endowed with a differential structure

## Definition (Wasserstein distance)

Taking  $c(x, y) = |x - y|^2$ , the quantity

$$W_2(\mu, \nu) = \min_{\gamma \in \Gamma(\mu, \nu)} \left\{ \left( \int_{\mathbb{R}^{2d}} |x - y|^2 d\gamma(x, y) \right)^{1/2} \right\}$$

defines a *distance* over  $\mathcal{P}_c(\mathbb{R}^d)$ .

$\sim$  Infimum of  $L^2$ -distances over the couplings  $\gamma \in \Gamma(\mu, \nu)$ .

## Proposition (Some useful properties)

- ◇  $W_2$  metrizes the usual weak- $*$  topology of measures.
- ◇  $(\mathcal{P}_c(\mathbb{R}^d), W_2)$  is a complete and separable metric space.
- ◇  $(\mathcal{P}_c(\mathbb{R}^d), W_2)$  can be endowed with a differential structure

## Definition (Wasserstein distance)

Taking  $c(x, y) = |x - y|^2$ , the quantity

$$W_2(\mu, \nu) = \min_{\gamma \in \Gamma(\mu, \nu)} \left\{ \left( \int_{\mathbb{R}^{2d}} |x - y|^2 d\gamma(x, y) \right)^{1/2} \right\}$$

defines a *distance* over  $\mathcal{P}_c(\mathbb{R}^d)$ .

$\sim$  Infimum of  $L^2$ -distances over the couplings  $\gamma \in \Gamma(\mu, \nu)$ .

## Proposition (Some useful properties)

- ◇  $W_2$  metrizes the usual weak- $*$  topology of measures.
- ◇  $(\mathcal{P}_c(\mathbb{R}^d), W_2)$  is a complete and separable metric space.
- ◇  $(\mathcal{P}_c(\mathbb{R}^d), W_2)$  can be endowed with a differential structure

Theorem (Ambrosio, Gangbo '07 - Piccoli, Rossi '13)

*Under Cauchy-Lipschitz assumptions on  $v$ , the Cauchy problem*

$$\partial_t \mu(t) + \nabla \cdot (v[\mu(t)](t, \cdot) \mu(t)) = 0, \quad \mu(0) = \mu^0 \in \mathcal{P}_c(\mathbb{R}^d)$$

*has a unique Lipschitz solution which is continuous w.r.t. its initial datum.*

*The solution  $t \mapsto \mu(t)$  is given explicitly by*

$$\mu(t) = \Phi_{(0,t)}^v(\cdot)_{\#} \mu^0 \quad (\text{image measure})$$

*with  $(\Phi_{(0,t)}^v(\cdot))_t$  the geometric flow generated by  $v[\mu(t)](t, \cdot)$ .*

$\hookrightarrow$  **Continuity Equations are Wasserstein ODEs !**

Theorem (Ambrosio, Gangbo '07 - Piccoli, Rossi '13)

*Under Cauchy-Lipschitz assumptions on  $v$ , the Cauchy problem*

$$\partial_t \mu(t) + \nabla \cdot (v[\mu(t)](t, \cdot) \mu(t)) = 0, \quad \mu(0) = \mu^0 \in \mathcal{P}_c(\mathbb{R}^d)$$

*has a **unique Lipschitz solution** which is **continuous w.r.t. its initial datum**.*

*The solution  $t \mapsto \mu(t)$  is given explicitly by*

$$\mu(t) = \Phi_{(0,t)}^v(\cdot)_{\#} \mu^0 \quad (\text{image measure})$$

*with  $(\Phi_{(0,t)}^v(\cdot))_t$  the **geometric flow** generated by  $v[\mu(t)](t, \cdot)$ .*

$\hookrightarrow$  **Continuity Equations are Wasserstein ODEs !**



Theorem (Ambrosio, Gangbo '07 - Piccoli, Rossi '13)

*Under Cauchy-Lipschitz assumptions on  $v$ , the Cauchy problem*

$$\partial_t \mu(t) + \nabla \cdot (v[\mu(t)](t, \cdot) \mu(t)) = 0, \quad \mu(0) = \mu^0 \in \mathcal{P}_c(\mathbb{R}^d)$$

*has a **unique Lipschitz solution** which is **continuous w.r.t. its initial datum**.*

*The solution  $t \mapsto \mu(t)$  is given explicitly by*

$$\mu(t) = \Phi_{(0,t)}^v(\cdot) \# \mu^0 \quad (\text{image measure})$$

*with  $(\Phi_{(0,t)}^v(\cdot))_t$  the **geometric flow generated by  $v[\mu(t)](t, \cdot)$** .*

$\hookrightarrow$  **Continuity Equations are Wasserstein ODEs !**

Theorem (Ambrosio, Gangbo '07 - Piccoli, Rossi '13)

*Under Cauchy-Lipschitz assumptions on  $v$ , the Cauchy problem*

$$\partial_t \mu(t) + \nabla \cdot (v[\mu(t)](t, \cdot) \mu(t)) = 0, \quad \mu(0) = \mu^0 \in \mathcal{P}_c(\mathbb{R}^d)$$

*has a **unique Lipschitz solution** which is **continuous w.r.t. its initial datum**.*

*The solution  $t \mapsto \mu(t)$  is given explicitly by*

$$\mu(t) = \Phi_{(0,t)}^v(\cdot) \# \mu^0 \quad (\text{image measure})$$

*with  $(\Phi_{(0,t)}^v(\cdot))_t$  the **geometric flow generated by  $v[\mu(t)](t, \cdot)$** .*

$\hookrightarrow$  **Continuity Equations are Wasserstein ODEs !**

# Outline of the talk

- 1 Introduction and Motivations
- 2 Optimal Transport Theory & Wasserstein Spaces
- 3 Optimal Control Problems and Pontryagin Maximum Principle

Optimal control problem in  $(\mathcal{P}_c(\mathbb{R}^d), W_2)$

$$\begin{cases} \min_{u \in \mathcal{U}} \left[ \int_0^T L(\mu(t), u(t)) dt + \varphi(\mu(T)) \right] \\ \text{s.t.} \begin{cases} \partial_t \mu(t) + \nabla \cdot ((v[\mu(t)](t, \cdot) + u(t, \cdot))\mu(t)) = 0, \\ \mu(0) = \mu^0 \in \mathcal{P}_c(\mathbb{R}^d), \end{cases} \end{cases}$$

where  $\mathcal{U}$  is defined by

$$\mathcal{U} = \left\{ u \in L^\infty([0, T], C^1(\mathbb{R}^d, \mathbb{R}^d)) \text{ s.t. } \|u(t)\|_{C^1(\mathbb{R}^d, \mathbb{R}^d)} \leq M \right\}$$

Question : Choice of the control set  $\mathcal{U}$

$\hookrightarrow$  Cauchy-Lipschitz requires  $u(t) \in \text{Lip}_{\text{loc}}(\mathbb{R}^d, \mathbb{R}^d) \oplus \text{sublinear}$ .

$\hookrightarrow$  PMP requires  $C^1(\mathbb{R}^d, \mathbb{R}^d) \oplus \text{uniformly Lipschitz}$

Optimal control problem in  $(\mathcal{P}_c(\mathbb{R}^d), W_2)$

$$\begin{cases} \min_{u \in \mathcal{U}} \left[ \int_0^T L(\mu(t), u(t)) dt + \varphi(\mu(T)) \right] \\ \text{s.t.} \begin{cases} \partial_t \mu(t) + \nabla \cdot ((v[\mu(t)](t, \cdot) + u(t, \cdot))\mu(t)) = 0, \\ \mu(0) = \mu^0 \in \mathcal{P}_c(\mathbb{R}^d), \end{cases} \end{cases}$$

where  $\mathcal{U}$  is defined by

$$\mathcal{U} = \left\{ u \in L^\infty([0, T], C^1(\mathbb{R}^d, \mathbb{R}^d)) \text{ s.t. } \|u(t)\|_{C^1(\mathbb{R}^d, \mathbb{R}^d)} \leq M \right\}$$

Question : Choice of the control set  $\mathcal{U}$

↪ Cauchy-Lipschitz requires  $u(t) \in \text{Lip}_{\text{loc}}(\mathbb{R}^d, \mathbb{R}^d) \oplus \text{sublinear}$ .

↪ PMP requires  $C^1(\mathbb{R}^d, \mathbb{R}^d) \oplus \text{uniformly Lipschitz}$

Optimal control problem in  $(\mathcal{P}_c(\mathbb{R}^d), W_2)$

$$\begin{cases} \min_{u \in \mathcal{U}} \left[ \int_0^T L(\mu(t), u(t)) dt + \varphi(\mu(T)) \right] \\ \text{s.t.} \begin{cases} \partial_t \mu(t) + \nabla \cdot ((v[\mu(t)](t, \cdot) + u(t, \cdot))\mu(t)) = 0, \\ \mu(0) = \mu^0 \in \mathcal{P}_c(\mathbb{R}^d), \end{cases} \end{cases}$$

where  $\mathcal{U}$  is defined by

$$\mathcal{U} = \left\{ u \in L^\infty([0, T], C^1(\mathbb{R}^d, \mathbb{R}^d)) \text{ s.t. } \|u(t)\|_{C^1(\mathbb{R}^d, \mathbb{R}^d)} \leq M \right\}$$

Question : Choice of the control set  $\mathcal{U}$

$\hookrightarrow$  Cauchy-Lipschitz requires  $u(t) \in \text{Lip}_{\text{loc}}(\mathbb{R}^d, \mathbb{R}^d) \oplus \text{sublinear}$ .

$\hookrightarrow$  PMP requires  $C^1(\mathbb{R}^d, \mathbb{R}^d) \oplus \text{uniformly Lipschitz}$

Optimal control problem in  $(\mathcal{P}_c(\mathbb{R}^d), W_2)$

$$\begin{cases} \min_{u \in \mathcal{U}} \left[ \int_0^T L(\mu(t), u(t)) dt + \varphi(\mu(T)) \right] \\ \text{s.t.} \begin{cases} \partial_t \mu(t) + \nabla \cdot ((v[\mu(t)](t, \cdot) + u(t, \cdot))\mu(t)) = 0, \\ \mu(0) = \mu^0 \in \mathcal{P}_c(\mathbb{R}^d), \end{cases} \end{cases}$$

where  $\mathcal{U}$  is defined by

$$\mathcal{U} = \left\{ u \in L^\infty([0, T], C^1(\mathbb{R}^d, \mathbb{R}^d)) \text{ s.t. } \|u(t)\|_{C^1(\mathbb{R}^d, \mathbb{R}^d)} \leq M \right\}$$

Question : Choice of the control set  $\mathcal{U}$

↪ Cauchy-Lipschitz requires  $u(t) \in \text{Lip}_{\text{loc}}(\mathbb{R}^d, \mathbb{R}^d) \oplus \text{sublinear}$ .

↪ PMP requires  $C^1(\mathbb{R}^d, \mathbb{R}^d) \oplus \text{uniformly Lipschitz}$

- ◇ **Existence results** : Mean-field and  $\Gamma$ -convergence approaches (*Bongini, Fornasier, Piccoli, Rossi, Solombrino*) or classical PDE methods (*Achdou, Laurière*).
- ◇ **Numerical methods** : Good methods in the transport  $\oplus$  diffusion case (*Albi, Pareschi, Toscani, Zanella,...*).  
 $\rightsquigarrow$  More ad hoc and difficult for pure transport.
- ◇ **Optimality conditions** : Hamilton-Jacobi Optimality conditions (*Cavagnari, Marigonda*)  
 $\hookrightarrow$  Recent results on Pontryagin-type conditions (*Bongini, Fornasier, Rossi, Solombrino '17 & B., Rossi '18*).



- ◇ **Existence results** : Mean-field and  $\Gamma$ -convergence approaches (*Bongini, Fornasier, Piccoli, Rossi, Solombrino*) or classical PDE methods (*Achdou, Laurière*).
- ◇ **Numerical methods** : Good methods in the transport  $\oplus$  diffusion case (*Albi, Pareschi, Toscani, Zanella,...*).  
 $\rightsquigarrow$  More ad hoc and difficult for pure transport.

- ◇ **Optimality conditions** : Hamilton-Jacobi Optimality conditions (*Cavagnari, Marigonda*)  
 $\hookrightarrow$  Recent results on Pontryagin-type conditions (*Bongini, Fornasier, Rossi, Solombrino '17 & B., Rossi '18*).

- ◇ **Existence results** : Mean-field and  $\Gamma$ -convergence approaches (*Bongini, Fornasier, Piccoli, Rossi, Solombrino*) or classical PDE methods (*Achdou, Laurière*).
- ◇ **Numerical methods** : Good methods in the transport  $\oplus$  diffusion case (*Albi, Pareschi, Toscani, Zanella,...*).  
 $\rightsquigarrow$  More ad hoc and difficult for pure transport.
- ◇ **Optimality conditions** : Hamilton-Jacobi Optimality conditions (*Cavagnari, Marigonda*)  
 $\hookrightarrow$  Recent results on Pontryagin-type conditions (*Bongini, Fornasier, Rossi, Solombrino '17 & B., Rossi '18*).

- ◇ **Existence results** : Mean-field and  $\Gamma$ -convergence approaches (*Bongini, Fornasier, Piccoli, Rossi, Solombrino*) or classical PDE methods (*Achdou, Laurière*).
- ◇ **Numerical methods** : Good methods in the transport  $\oplus$  diffusion case (*Albi, Pareschi, Toscani, Zanella,...*).  
 $\rightsquigarrow$  More ad hoc and difficult for pure transport.
- ◇ **Optimality conditions** : Hamilton-Jacobi Optimality conditions (*Cavagnari, Marigonda*)  
 $\hookrightarrow$  Recent results on Pontryagin-type conditions (*Bongini, Fornasier, Rossi, Solombrino '17 & B., Rossi '18*).

- ◇ **Existence results** : Mean-field and  $\Gamma$ -convergence approaches (*Bongini, Fornasier, Piccoli, Rossi, Solombrino*) or classical PDE methods (*Achdou, Laurière*).
- ◇ **Numerical methods** : Good methods in the transport  $\oplus$  diffusion case (*Albi, Pareschi, Toscani, Zanella,...*).  
 $\rightsquigarrow$  More ad hoc and difficult for pure transport.
- ◇ **Optimality conditions** : Hamilton-Jacobi Optimality conditions (*Cavagnari, Marigonda*)  
 $\hookrightarrow$  Recent results on Pontryagin-type conditions (*Bongini, Fornasier, Rossi, Solombrino '17 & B., Rossi '18*).

## Theorem (Unconstrained and smooth PMP on $\mathbb{R}^d$ )

Let  $(u^*(\cdot), x^*(\cdot))$  be an optimal pair control-trajectory for the problem

$$\begin{cases} \min_{u \in \mathcal{U}} \left[ \int_0^T L(t, x(t), u(t)) dt + \varphi(x(T)) \right] \\ \dot{x}(t) = f(t, x(t), u(t)) , \quad x(0) = x^0. \end{cases}$$

Then, there exists a curve  $p^*(\cdot)$  called *costate* such that

$$\begin{cases} \dot{x}^*(t) = \nabla_p \mathcal{H}(t, x^*(t), p^*(t)) , \quad x^*(0) = x^0, \\ \dot{p}^*(t) = -\nabla_x \mathcal{H}(t, x^*(t), p^*(t)) , \quad p^*(T) = -\nabla \varphi(x^*(T)), \end{cases}$$

where

$$\mathcal{H}(t, x^*(t), p^*(t)) = \max_{\omega \in \mathcal{U}} [\langle p^*(t), f(t, x^*(t), \omega) \rangle - L(t, x^*(t), \omega)] .$$

## Theorem (Unconstrained and smooth PMP on $\mathbb{R}^d$ )

Let  $(u^*(\cdot), x^*(\cdot))$  be an optimal pair control-trajectory for the problem

$$\begin{cases} \min_{u \in \mathcal{U}} \left[ \int_0^T L(t, x(t), u(t)) dt + \varphi(x(T)) \right] \\ \dot{x}(t) = f(t, x(t), u(t)) , \quad x(0) = x^0. \end{cases}$$

Then, there exists a curve  $p^*(\cdot)$  called *costate* such that

$$\begin{cases} \dot{x}^*(t) = \nabla_p \mathcal{H}(t, x^*(t), p^*(t)) & , \quad x^*(0) = x^0, \\ \dot{p}^*(t) = -\nabla_x \mathcal{H}(t, x^*(t), p^*(t)) & , \quad p^*(T) = -\nabla \varphi(x^*(T)), \end{cases}$$

where

$$\mathcal{H}(t, x^*(t), p^*(t)) = \max_{\omega \in \mathcal{U}} [\langle p^*(t), f(t, x^*(t), \omega) \rangle - L(t, x^*(t), \omega)] .$$

## Analogies with the finite dimensional setting

◇ Couple state-costate  $(x^*(\cdot), p^*(\cdot)) \in \text{Lip}([0, T], \mathbb{R}^{2d})$

↪ measures  $\nu^*(\cdot) \in \text{Lip}([0, T], \mathcal{P}_c(\mathbb{R}^{2d}))$  on the product space.

◇ Euclidean scalar product  $\langle p^*(t), f(t, x^*(t), u^*(t)) \rangle$

↪ Hilbertian  $L^2(\nu^*(t))$ -scalar product

◇ Classical Hamiltonian flow of  $\mathcal{H}(t, x, p)$  with respect to  $|\cdot|_2$ .

↪ Metric Hamiltonian flow of  $\mathbb{H}(t, \nu)$  with respect to  $W_2(\cdot, \cdot)$ .

## Analogies with the finite dimensional setting

◇ Couple state-costate  $(x^*(\cdot), p^*(\cdot)) \in \text{Lip}([0, T], \mathbb{R}^{2d})$

↪ measures  $\nu^*(\cdot) \in \text{Lip}([0, T], \mathcal{P}_c(\mathbb{R}^{2d}))$  on the product space.

◇ Euclidean scalar product  $\langle p^*(t), f(t, x^*(t), u^*(t)) \rangle$

↪ Hilbertian  $L^2(\nu^*(t))$ -scalar product

◇ Classical Hamiltonian flow of  $\mathcal{H}(t, x, p)$  with respect to  $|\cdot|_2$ .

↪ Metric Hamiltonian flow of  $\mathbb{H}(t, \nu)$  with respect to  $W_2(\cdot, \cdot)$ .



Analogies with the finite dimensional setting

◇ Couple state-costate  $(x^*(\cdot), p^*(\cdot)) \in \text{Lip}([0, T], \mathbb{R}^{2d})$

↪ *measures  $\nu^*(\cdot) \in \text{Lip}([0, T], \mathcal{P}_c(\mathbb{R}^{2d}))$  on the product space.*

◇ Euclidean scalar product  $\langle p^*(t), f(t, x^*(t), u^*(t)) \rangle$

↪ *Hilbertian  $L^2(\nu^*(t))$ -scalar product*

◇ Classical Hamiltonian flow of  $\mathcal{H}(t, x, p)$  with respect to  $|\cdot|_2$ .

↪ *Metric Hamiltonian flow of  $\mathbb{H}(t, \nu)$  with respect to  $W_2(\cdot, \cdot)$ .*

Analogies with the finite dimensional setting

◇ Couple state-costate  $(x^*(\cdot), p^*(\cdot)) \in \text{Lip}([0, T], \mathbb{R}^{2d})$

↪ measures  $\nu^*(\cdot) \in \text{Lip}([0, T], \mathcal{P}_c(\mathbb{R}^{2d}))$  on the product space.

◇ Euclidean scalar product  $\langle p^*(t), f(t, x^*(t), u^*(t)) \rangle$

↪ Hilbertian  $L^2(\nu^*(t))$ -scalar product

◇ Classical Hamiltonian flow of  $\mathcal{H}(t, x, p)$  with respect to  $|\cdot|_2$ .

↪ Metric Hamiltonian flow of  $\mathbb{H}(t, \nu)$  with respect to  $W_2(\cdot, \cdot)$ .

Analogies with the finite dimensional setting

◇ Couple state-costate  $(x^*(\cdot), p^*(\cdot)) \in \text{Lip}([0, T], \mathbb{R}^{2d})$

↪ measures  $\nu^*(\cdot) \in \text{Lip}([0, T], \mathcal{P}_c(\mathbb{R}^{2d}))$  on the product space.

◇ Euclidean scalar product  $\langle p^*(t), f(t, x^*(t), u^*(t)) \rangle$

↪ Hilbertian  $L^2(\nu^*(t))$ -scalar product

◇ Classical Hamiltonian flow of  $\mathcal{H}(t, x, p)$  with respect to  $|\cdot|_2$ .

↪ Metric Hamiltonian flow of  $\mathbb{H}(t, \nu)$  with respect to  $W_2(\cdot, \cdot)$ .

Analogies with the finite dimensional setting

◇ Couple state-costate  $(x^*(\cdot), p^*(\cdot)) \in \text{Lip}([0, T], \mathbb{R}^{2d})$

↪ measures  $\nu^*(\cdot) \in \text{Lip}([0, T], \mathcal{P}_c(\mathbb{R}^{2d}))$  on the product space.

◇ Euclidean scalar product  $\langle p^*(t), f(t, x^*(t), u^*(t)) \rangle$

↪ Hilbertian  $L^2(\nu^*(t))$ -scalar product

◇ Classical Hamiltonian flow of  $\mathcal{H}(t, x, p)$  with respect to  $|\cdot|_2$ .

↪ Metric Hamiltonian flow of  $\mathbb{H}(t, \nu)$  with respect to  $W_2(\cdot, \cdot)$ .

Analogies with the finite dimensional setting

◇ Couple state-costate  $(x^*(\cdot), p^*(\cdot)) \in \text{Lip}([0, T], \mathbb{R}^{2d})$

↪ *measures  $\nu^*(\cdot) \in \text{Lip}([0, T], \mathcal{P}_c(\mathbb{R}^{2d}))$  on the product space.*

◇ Euclidean scalar product  $\langle p^*(t), f(t, x^*(t), u^*(t)) \rangle$

↪ *Hilbertian  $L^2(\nu^*(t))$ -scalar product*

◇ Classical Hamiltonian flow of  $\mathcal{H}(t, x, p)$  with respect to  $|\cdot|_2$ .

↪ *Metric Hamiltonian flow of  $\mathbb{H}(t, \nu)$  with respect to  $W_2(\cdot, \cdot)$ .*

## Theorem (B., Rossi '18)

If  $(u^*(\cdot), \mu^*(\cdot))$  is an optimal pair for  $(\mathcal{P}_{OC})$ , there exists a curve  $\nu^*(\cdot) \in \text{Lip}([0, T], \mathcal{P}_c(\mathbb{R}^{2d}))$  solution of

$$\begin{cases} \partial_t \nu^*(t) + \nabla \cdot (\mathbb{J}_{2d} \nabla_\nu \mathbb{H}(t, \nu^*(t)) \nu^*(t)) = 0, \\ \pi_{\#}^1 \nu^*(0) = \mu^0, \\ \pi_{\#}^2 \nu^*(T) = (-\nabla_\mu \varphi(\mu^*(T))_{\#} \mu^*(T), \end{cases}$$

where  $(x, r) \mapsto \nabla_\nu \mathbb{H}(t, \nu^*(t))(x, r)$  is the **Wasserstein gradient** of the maximized Hamiltonian.

**Idea of proof :** needle-variations  $\oplus$  geometric structure of solutions & differential structure of  $(\mathcal{P}_c(\mathbb{R}^d), W_2)$

## Theorem (B., Rossi '18)

If  $(u^*(\cdot), \mu^*(\cdot))$  is an optimal pair for  $(\mathcal{P}_{OC})$ , there exists a curve  $\nu^*(\cdot) \in \text{Lip}([0, T], \mathcal{P}_c(\mathbb{R}^{2d}))$  solution of

$$\begin{cases} \partial_t \nu^*(t) + \nabla \cdot (\mathbb{J}_{2d} \nabla_\nu \mathbb{H}(t, \nu^*(t)) \nu^*(t)) = 0, \\ \pi_{\#}^1 \nu^*(0) = \mu^0, \\ \pi_{\#}^2 \nu^*(T) = (-\nabla_\mu \varphi(\mu^*(T)))_{\#} \mu^*(T), \end{cases}$$

where  $(x, r) \mapsto \nabla_\nu \mathbb{H}(t, \nu^*(t))(x, r)$  is the **Wasserstein gradient** of the maximized Hamiltonian.

**Idea of proof :** *needle-variations  $\oplus$  geometric structure of solutions & differential structure of  $(\mathcal{P}_c(\mathbb{R}^d), W_2)$*

## Theorem (B., Rossi '18)

If  $(u^*(\cdot), \mu^*(\cdot))$  is an optimal pair for  $(\mathcal{P}_{OC})$ , there exists a curve  $\nu^*(\cdot) \in \text{Lip}([0, T], \mathcal{P}_c(\mathbb{R}^{2d}))$  solution of

$$\begin{cases} \partial_t \nu^*(t) + \nabla \cdot (\mathbb{J}_{2d} \nabla_\nu \mathbb{H}(t, \nu^*(t)) \nu^*(t)) = 0, \\ \pi_\#^1 \nu^*(0) = \mu^0, \\ \pi_\#^2 \nu^*(T) = (-\nabla_\mu \varphi(\mu^*(T))_\# \mu^*(T), \end{cases}$$

where  $(x, r) \mapsto \nabla_\nu \mathbb{H}(t, \nu^*(t))(x, r)$  is the **Wasserstein gradient** of the maximized Hamiltonian.

**Idea of proof :** needle-variations  $\oplus$  geometric structure of solutions & differential structure of  $(\mathcal{P}_c(\mathbb{R}^d), W_2)$



## Developments with (and without) PMP flavours

- ◇ Design of a shooting method
- ◇ Study gradient type methods with the same kind of duality involved in this PMP (measure on the product space)
- ◇ Intrinsic Lipschitz regularity for mean field optimal controls

## Developments with (and without) PMP flavours

- ◇ Design of a shooting method
- ◇ Study gradient type methods with the same kind of duality involved in this PMP (measure on the product space)
- ◇ Intrinsic Lipschitz regularity for mean field optimal controls

## Developments with (and without) PMP flavours

- ◇ Design of a shooting method
- ◇ Study gradient type methods with the same kind of duality involved in this PMP (measure on the product space)
- ◇ Intrinsic Lipschitz regularity for mean field optimal controls

# Thank you for your attention !

