

Mean-Field Control and Continuity Inclusions

Benoît Bonnet

(in collaboration with H. Frankowska)

59th Conference on Decision and Control

From my kitchen to Jeju Island

November 5, 2020

Outline of the talk

Introduction to mean-field control

From differential inclusions to continuity inclusions

Main contributions

Outline of the talk

Introduction to mean-field control

From differential inclusions to continuity inclusions

Main contributions

Introduction – *Multi-agent systems*

A multi-agent system is a large ensemble of interacting things




Introduction – Mean-field limits of particle systems

First observation (dimensionality-related problems)

The number N of agents is usually **very large** \rightsquigarrow numerical issues

Idea: Approximation procedure using **mean-field limits!**



System of ODEs on N agents $(x_1, \dots, x_N) \in (\mathbb{R}^d)^N$

$$\left(\mu_N = \frac{1}{N} \sum_{i=1}^N \delta_{x_i} \right)$$


Single PDE on the density of agents $\mu : \mathbb{R}^d \rightarrow \mathbb{R}$.

Introduction – Mean-field limits of particle systems

First observation (dimensionality-related problems)

The number N of agents is usually **very large** \rightsquigarrow **numerical issues**

Idea: Approximation procedure using **mean-field limits!**



System of ODEs on N agents $(x_1, \dots, x_N) \in (\mathbb{R}^d)^N$

$$\left(\mu_N = \frac{1}{N} \sum_{i=1}^N \delta_{x_i} \right)$$


Single PDE on the density of agents $\mu : \mathbb{R}^d \rightarrow \mathbb{R}$.

Introduction – Mean-field limits of particle systems

First observation (dimensionality-related problems)

The number N of agents is usually **very large** \rightsquigarrow **numerical issues**

Idea: Approximation procedure using **mean-field limits!**



System of ODEs on N agents $(x_1, \dots, x_N) \in (\mathbb{R}^d)^N$

$$\left(\mu_N = \frac{1}{N} \sum_{i=1}^N \delta_{x_i} \right)$$


Single PDE on the density of agents $\mu : \mathbb{R}^d \rightarrow \mathbb{R}$.

Introduction – Mean-field limits of particle systems

First observation (dimensionality-related problems)

The number N of agents is usually **very large** \rightsquigarrow **numerical issues**

Idea: Approximation procedure using **mean-field limits!**



System of ODEs on N agents $(x_1, \dots, x_N) \in (\mathbb{R}^d)^N$

$$\left(\mu_N = \frac{1}{N} \sum_{i=1}^N \delta_{x_i} \right)$$


Single PDE on the density of agents $\mu : \mathbb{R}^d \rightarrow \mathbb{R}$.

Introduction – *Continuity equations*

Multi-agent dynamics are modelled via **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t, \mu(t), \cdot) \mu(t)) = 0,$$

where $\mu(t) \in \mathcal{P}_c(\mathbb{R}^d)$ and $\mathbf{v} : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times \mathbb{R}^d \rightarrow \mathbb{R}^d$.




Introduction – *Continuity equations*

Multi-agent dynamics are modelled via **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t, \mu(t), \cdot) \mu(t)) = 0,$$

where $\mu(t) \in \mathcal{P}_c(\mathbb{R}^d)$ and $\mathbf{v} : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times \mathbb{R}^d \rightarrow \mathbb{R}^d$.




Introduction – *Continuity equations*

Multi-agent dynamics are modelled via **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t, \mu(t), \cdot) \mu(t)) = 0,$$

where $\mu(t) \in \mathcal{P}_c(\mathbb{R}^d)$ and $\mathbf{v} : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times \mathbb{R}^d \rightarrow \mathbb{R}^d$.



Introduction – Mean-field optimal control

Mean-field control \rightsquigarrow control problems on **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t, \mu(t), u(t), \cdot) \mu(t)) = 0,$$

where $u : [0, T] \rightarrow U$ is a **control law**.

- **Existence of optimal trajectories** \rightsquigarrow (B., Frankowska, Fornasier, Marigonda, Quincampoix, Pogodaev, Rossi, Savaré, Solombrino, etc...)
- **1st-order optimality conditions** \rightsquigarrow (B., Carlier, Frankowska, Gomes, Marigonda, Quincampoix, Piccoli, Pogodaev, Rossi, etc...)
- **Properties of value functions** \rightsquigarrow (B., Frankowska, Santambrogio, Carlier, Crippa, Gomes, Marigonda, Quincampoix, etc...)
- **Optimal feedback synthesis** \rightsquigarrow Open for now

Natural question : General framework to study these problems ?
 \rightsquigarrow Yes! Reformulate controlled dynamics as differential inclusions

Introduction – Mean-field optimal control

Mean-field control \rightsquigarrow control problems on **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t, \mu(t), u(t), \cdot) \mu(t)) = 0,$$

where $u : [0, T] \rightarrow U$ is a **control law**.

- ◊ **Existence of optimal trajectories** \rightsquigarrow (B., Frankowska, Fornasier, Marigonda, Quincampoix, Pogodaev, Rossi, Savaré, Solombrino, etc...)
- ◊ **1st-order optimality conditions** \rightsquigarrow (B., Cavagnari, Frankowska, Gimenez, Marigonda, Quincampoix, Piccoli, Pogodaev, Rossi, etc...)
- ◊ **Properties of value functions** \rightsquigarrow (B., Frankowska, Lemaire, Cavagnari, Piccoli, Gimenez, Marigonda, Quincampoix, etc...)
- ◊ **Optimal feedback synthesis** \rightsquigarrow Open for now

Natural question : General framework to study these problems ?
 \rightsquigarrow Yet Reformulate controlled dynamics as differential inclusions

Introduction – Mean-field optimal control

Mean-field control \rightsquigarrow control problems on **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t, \mu(t), u(t), \cdot) \mu(t)) = 0,$$

where $u : [0, T] \rightarrow U$ is a **control law**.

- ◇ **Existence of optimal trajectories** \rightsquigarrow (B., Frankowska, Fornasier, Marigonda, Quincampoix, Pogodaev, Rossi, Savaré, Solombrino, etc...)
- ◇ **1st-order optimality conditions** \rightsquigarrow (B., Cavagnari, Frankowska, Gimenez, Marigonda, Quincampoix, Piccoli, Pogodaev, Rossi, etc...)
- ◇ **Properties of value functions** \rightsquigarrow (B., Frankowska, Cardaliaguet, Cavagnari, Gangbo, Gimenez, Marigonda, Quincampoix, etc...)
- ◇ **Optimal feedback synthesis** \rightsquigarrow Open for now

Natural question : General framework to study these problems ?
 \rightsquigarrow Yet reformulate controlled dynamics as differential inclusions

Introduction – *Mean-field optimal control*

Mean-field control \rightsquigarrow control problems on **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t, \mu(t), u(t), \cdot) \mu(t)) = 0,$$

where $u : [0, T] \rightarrow U$ is a **control law**.

- ◇ **Existence of optimal trajectories** \rightsquigarrow (B., Frankowska, Fornasier, Marigonda, Quincampoix, Pogodaev, Rossi, Savaré, Solombrino, etc...)
- ◇ **1st-order optimality conditions** \rightsquigarrow (B., Cavagnari, Frankowska, Gimenez, Marigonda, Quincampoix, Piccoli, Pogodaev, Rossi, etc...)
- ◇ **Properties of value functions** \rightsquigarrow (B., Frankowska, Cardaliaguet, Cavagnari, Gangbo, Gimenez, Marigonda, Quincampoix, etc...)
- ◇ **Optimal feedback synthesis** \rightsquigarrow Open for now

Natural question : General framework to study these problems ?
 \rightsquigarrow Yes! Reformulate controlled dynamics as **differential inclusions**

Introduction – Mean-field optimal control

Mean-field control \rightsquigarrow control problems on **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t, \mu(t), u(t), \cdot) \mu(t)) = 0,$$

where $u : [0, T] \rightarrow U$ is a **control law**.

- ◇ **Existence of optimal trajectories** \rightsquigarrow (B., Frankowska, Fornasier, Marigonda, Quincampoix, Pogodaev, Rossi, Savaré, Solombrino, etc...)
- ◇ **1st-order optimality conditions** \rightsquigarrow (B., Cavagnari, Frankowska, Gimenez, Marigonda, Quincampoix, Piccoli, Pogodaev, Rossi, etc...)
- ◇ **Properties of value functions** \rightsquigarrow (B., Frankowska, Cardaliaguet, Cavagnari, Gangbo, Gimenez, Marigonda, Quincampoix, etc...)
- ◇ **Optimal feedback synthesis** \rightsquigarrow Open for now

Natural question : General framework to study these problems ?
 \rightsquigarrow Yes! Reformulate controlled dynamics as **differential inclusions**

Introduction – Mean-field optimal control

Mean-field control \rightsquigarrow control problems on **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t, \mu(t), u(t), \cdot) \mu(t)) = 0,$$

where $u : [0, T] \rightarrow U$ is a **control law**.

- ◇ **Existence of optimal trajectories** \rightsquigarrow (B., Frankowska, Fornasier, Marigonda, Quincampoix, Pogodaev, Rossi, Savaré, Solombrino, etc...)
- ◇ **1st-order optimality conditions** \rightsquigarrow (B., Cavagnari, Frankowska, Gimenez, Marigonda, Quincampoix, Piccoli, Pogodaev, Rossi, etc...)
- ◇ **Properties of value functions** \rightsquigarrow (B., Frankowska, Cardaliaguet, Cavagnari, Gangbo, Gimenez, Marigonda, Quincampoix, etc...)
- ◇ **Optimal feedback synthesis** \rightsquigarrow Open for now

Natural question : General framework to study these problems ?
 \rightsquigarrow Yes! Reformulate controlled dynamics as **differential inclusions**

Introduction – Mean-field optimal control

Mean-field control \rightsquigarrow control problems on **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t, \mu(t), u(t), \cdot) \mu(t)) = 0,$$

where $u : [0, T] \rightarrow U$ is a **control law**.

- ◇ **Existence of optimal trajectories** \rightsquigarrow (B., Frankowska, Fornasier, Marigonda, Quincampoix, Pogodaev, Rossi, Savaré, Solombrino, etc...)
- ◇ **1st-order optimality conditions** \rightsquigarrow (B., Cavagnari, Frankowska, Gimenez, Marigonda, Quincampoix, Piccoli, Pogodaev, Rossi, etc...)
- ◇ **Properties of value functions** \rightsquigarrow (B., Frankowska, Cardaliaguet, Cavagnari, Gangbo, Gimenez, Marigonda, Quincampoix, etc...)
- ◇ **Optimal feedback synthesis** \rightsquigarrow Open for now

Natural question : General framework to study these problems ?

\hookrightarrow Yes! Reformulate controlled dynamics as **differential inclusions**

Introduction – Mean-field optimal control

Mean-field control \rightsquigarrow control problems on **continuity equations**

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t, \mu(t), u(t), \cdot) \mu(t)) = 0,$$

where $u : [0, T] \rightarrow U$ is a **control law**.

- ◇ **Existence of optimal trajectories** \rightsquigarrow (B., Frankowska, Fornasier, Marigonda, Quincampoix, Pogodaev, Rossi, Savaré, Solombrino, etc...)
- ◇ **1st-order optimality conditions** \rightsquigarrow (B., Cavagnari, Frankowska, Gimenez, Marigonda, Quincampoix, Piccoli, Pogodaev, Rossi, etc...)
- ◇ **Properties of value functions** \rightsquigarrow (B., Frankowska, Cardaliaguet, Cavagnari, Gangbo, Gimenez, Marigonda, Quincampoix, etc...)
- ◇ **Optimal feedback synthesis** \rightsquigarrow Open for now

Natural question : General framework to study these problems ?

\hookrightarrow Yes! Reformulate controlled dynamics as **differential inclusions**

Outline of the talk

Introduction to mean-field control

From differential inclusions to continuity inclusions

Main contributions

Differential inclusions – *The classical case*

Intuition: Differential inclusions \iff ODEs with **set-valued** r.h.s

Definition (differential inclusion)

Let \mathcal{M} be a smooth manifold and $F : \mathcal{M} \rightrightarrows T\mathcal{M}$. Then

$$\dot{x}(t) \in F(x(t)) \iff \begin{cases} x(0) = x_0, \\ x(\cdot) \in F(x(\cdot)), \quad \forall \cdot \in [0, T] \end{cases}$$



Differential inclusions – *The classical case*

Intuition: Differential inclusions \Longleftrightarrow ODEs with **set-valued** r.h.s

Definition (differential inclusion)

Let \mathcal{M} be a smooth manifold and $F : \mathcal{M} \rightrightarrows T\mathcal{M}$. Then

$$\dot{x}(t) \in F(x(t)) \Longleftrightarrow \begin{cases} \dot{x}(t) = f(t), \\ f(t) \in F(x(t)), \end{cases} \quad \mathcal{L}^1\text{-a.e. in } [0, T]$$




Differential inclusions – *The classical case*

Intuition: Differential inclusions \Longleftrightarrow ODEs with **set-valued** r.h.s

Definition (differential inclusion)

Let \mathcal{M} be a smooth manifold and $F : \mathcal{M} \rightrightarrows T\mathcal{M}$. Then

$$\dot{x}(t) \in F(x(t)) \Longleftrightarrow \begin{cases} \dot{x}(t) = f(t), \\ f(t) \in F(x(t)), \end{cases} \quad \mathcal{L}^1\text{-a.e. in } [0, T]$$




Differential inclusions – *The classical case*

Intuition: Differential inclusions \Longleftrightarrow ODEs with **set-valued** r.h.s

Definition (differential inclusion)

Let \mathcal{M} be a smooth manifold and $F : \mathcal{M} \rightrightarrows T\mathcal{M}$. Then

$$\dot{x}(t) \in F(x(t)) \Longleftrightarrow \begin{cases} \dot{x}(t) = f(t), \\ f(t) \in F(x(t)), \end{cases} \quad \mathcal{L}^1\text{-a.e. in } [0, T]$$



Differential inclusions – *Link with control system*

Key point: Control systems are **equivalent** to diff. inclusions

Equivalence (heuristic statement)

If $f : [0, T] \times \mathcal{M} \times U \rightarrow T\mathcal{M}$ is **Carathéodory**, then

$$\dot{x}(t) = f(t, x(t), u(t)) \iff \dot{x}(t) \in F(t, x(t)),$$

where

$$F(t, x) := \left\{ f(t, x, u) \text{ s.t. } u \in U \right\}.$$

Recap': Building differential inclusions

- 1) Check that the ambient space has a manifold-like structure
- 2) Identify an ODE structure \rightarrow make the vehicles not-washed

Differential inclusions – *Link with control system*

Key point: Control systems are **equivalent** to diff. inclusions

Equivalence (heuristic statement)

If $f : [0, T] \times \mathcal{M} \times U \rightarrow T\mathcal{M}$ is **Carathéodory**, then

$$\dot{x}(t) = f(t, x(t), u(t)) \iff \dot{x}(t) \in F(t, x(t)),$$

where

$$F(t, x) := \left\{ f(t, x, u) \text{ s.t. } u \in U \right\}.$$

Recap': Building differential inclusions

- Check that the ambient space has a manifold-like structure
- Identify an ODE structure \rightarrow make the vector field set-valued

Differential inclusions – *Link with control system*

Key point: Control systems are **equivalent** to diff. inclusions

Equivalence (heuristic statement)

If $f : [0, T] \times \mathcal{M} \times U \rightarrow T\mathcal{M}$ is **Carathéodory**, then

$$\dot{x}(t) = f(t, x(t), u(t)) \iff \dot{x}(t) \in F(t, x(t)),$$

where

$$F(t, x) := \left\{ f(t, x, u) \text{ s.t. } u \in U \right\}.$$

Recap': Building differential inclusions

- 1) Check that the ambient space has a manifold-like structure
- 2) Identify an ODE structure \rightsquigarrow make the velocities set-valued

Differential inclusions – *Link with control system*

Key point: Control systems are **equivalent** to diff. inclusions

Equivalence (heuristic statement)

If $f : [0, T] \times \mathcal{M} \times U \rightarrow T\mathcal{M}$ is **Carathéodory**, then

$$\dot{x}(t) = f(t, x(t), u(t)) \iff \dot{x}(t) \in F(t, x(t)),$$

where

$$F(t, x) := \left\{ f(t, x, u) \text{ s.t. } u \in U \right\}.$$

Recap': Building differential inclusions

- 1) Check that the ambient space has a **manifold-like** structure
- 2) Identify an ODE structure \rightsquigarrow make the velocities **set-valued**

Differential inclusions – *Link with control system*

Key point: Control systems are **equivalent** to diff. inclusions

Equivalence (heuristic statement)

If $f : [0, T] \times \mathcal{M} \times U \rightarrow T\mathcal{M}$ is **Carathéodory**, then

$$\dot{x}(t) = f(t, x(t), u(t)) \iff \dot{x}(t) \in F(t, x(t)),$$

where

$$F(t, x) := \left\{ f(t, x, u) \text{ s.t. } u \in U \right\}.$$

Recap': Building differential inclusions

- 1) Check that the ambient space has a **manifold-like** structure
- 2) Identify an ODE structure \rightsquigarrow make the velocities **set-valued**

Differential inclusions – *Link with control system*

Key point: Control systems are **equivalent** to diff. inclusions

Equivalence (heuristic statement)

If $f : [0, T] \times \mathcal{M} \times U \rightarrow T\mathcal{M}$ is **Carathéodory**, then

$$\dot{x}(t) = f(t, x(t), u(t)) \iff \dot{x}(t) \in F(t, x(t)),$$

where

$$F(t, x) := \left\{ f(t, x, u) \text{ s.t. } u \in U \right\}.$$

Recap': Building differential inclusions

- 1) Check that the ambient space has a **manifold-like** structure
- 2) Identify an ODE structure \rightsquigarrow make the velocities **set-valued**

Differential inclusions – *What about continuity equations ?*

Idea: Endow $\mathcal{P}_2(\mathbb{R}^d)$ with the **optimal transport** metric W_2

$$W_2(\mu, \nu) := \inf_{\gamma \in \Gamma(\mu, \nu)} \left(\int_{\mathbb{R}^{2d}} |x-y|^2 d\gamma(x, y) \right)^{1/2}$$



Wasserstein geometry & ODEs [Ambrosio, Gigli, McCann, Otto, Savaré]

- ◇ $(\mathcal{P}_2(\mathbb{R}^d), W_2)$ is a “**manifold**” with $T_\mu \mathcal{P}_2(\mathbb{R}^d) \subset L^2(\mu)$
- ◇ Continuity equations

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t, \mu(t), \cdot) \mu(t)) = 0,$$


are **ODEs** over $\mathcal{P}_2(\mathbb{R}^d)$, with velocities $\mathbf{v}(t, \mu(t), \cdot) \in L^2(\mu(t))$

Conclusion: Make $(t, \mu) \mapsto \mathbf{v}(t, \mu, \cdot) \in L^2(\mu)$ **set-valued** !

Differential inclusions – *What about continuity equations ?*

Idea: Endow $\mathcal{P}_2(\mathbb{R}^d)$ with the **optimal transport** metric W_2

$$W_2(\mu, \nu) := \inf_{\gamma \in \Gamma(\mu, \nu)} \left(\int_{\mathbb{R}^{2d}} |x-y|^2 d\gamma(x, y) \right)^{1/2}$$



Wasserstein geometry & ODEs [Ambrosio, Gigli, McCann, Otto, Savaré]

- ◇ $(\mathcal{P}_2(\mathbb{R}^d), W_2)$ is a “**manifold**” with $T_\mu \mathcal{P}_2(\mathbb{R}^d) \subset L^2(\mu)$
- ◇ Continuity equations

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t, \mu(t), \cdot) \mu(t)) = 0,$$


are **ODEs** over $\mathcal{P}_2(\mathbb{R}^d)$, with velocities $\mathbf{v}(t, \mu(t), \cdot) \in L^2(\mu(t))$

Conclusion: Make $(t, \mu) \mapsto \mathbf{v}(t, \mu, \cdot) \in L^2(\mu)$ **set-valued** !

Differential inclusions – *What about continuity equations ?*

Idea: Endow $\mathcal{P}_2(\mathbb{R}^d)$ with the **optimal transport** metric W_2

$$W_2(\mu, \nu) := \inf_{\gamma \in \Gamma(\mu, \nu)} \left(\int_{\mathbb{R}^{2d}} |x-y|^2 d\gamma(x, y) \right)^{1/2}$$



Wasserstein geometry & ODEs [Ambrosio, Gigli, McCann, Otto, Savaré]

- ◇ $(\mathcal{P}_2(\mathbb{R}^d), W_2)$ is a “**manifold**” with $T_\mu \mathcal{P}_2(\mathbb{R}^d) \subset L^2(\mu)$
- ◇ Continuity equations

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t, \mu(t), \cdot) \mu(t)) = 0,$$


are **ODEs** over $\mathcal{P}_2(\mathbb{R}^d)$, with velocities $\mathbf{v}(t, \mu(t), \cdot) \in L^2(\mu(t))$

Conclusion: Make $(t, \mu) \mapsto \mathbf{v}(t, \mu, \cdot) \in L^2(\mu)$ **set-valued** !

Differential inclusions – *What about continuity equations ?*

Idea: Endow $\mathcal{P}_2(\mathbb{R}^d)$ with the **optimal transport** metric W_2

$$W_2(\mu, \nu) := \inf_{\gamma \in \Gamma(\mu, \nu)} \left(\int_{\mathbb{R}^{2d}} |x-y|^2 d\gamma(x, y) \right)^{1/2}$$



Wasserstein geometry & ODEs [Ambrosio, Gigli, McCann, Otto, Savaré]

- ◇ $(\mathcal{P}_2(\mathbb{R}^d), W_2)$ is a “**manifold**” with $T_\mu \mathcal{P}_2(\mathbb{R}^d) \subset L^2(\mu)$
- ◇ Continuity equations

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t, \mu(t), \cdot) \mu(t)) = 0,$$


are **ODEs** over $\mathcal{P}_2(\mathbb{R}^d)$, with velocities $\mathbf{v}(t, \mu(t), \cdot) \in L^2(\mu(t))$

Conclusion: Make $(t, \mu) \mapsto \mathbf{v}(t, \mu, \cdot) \in L^2(\mu)$ **set-valued** !

Differential inclusions – *What about continuity equations ?*

Idea: Endow $\mathcal{P}_2(\mathbb{R}^d)$ with the **optimal transport** metric W_2

$$W_2(\mu, \nu) := \inf_{\gamma \in \Gamma(\mu, \nu)} \left(\int_{\mathbb{R}^{2d}} |x-y|^2 d\gamma(x, y) \right)^{1/2}$$



Wasserstein geometry & ODEs [Ambrosio, Gigli, McCann, Otto, Savaré]

- ◇ $(\mathcal{P}_2(\mathbb{R}^d), W_2)$ is a “**manifold**” with $T_\mu \mathcal{P}_2(\mathbb{R}^d) \subset L^2(\mu)$
- ◇ Continuity equations

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t, \mu(t), \cdot) \mu(t)) = 0,$$

are **ODEs** over $\mathcal{P}_2(\mathbb{R}^d)$, with velocities $\mathbf{v}(t, \mu(t), \cdot) \in L^2(\mu(t))$

Conclusion: Make $(t, \mu) \mapsto \mathbf{v}(t, \mu, \cdot) \in L^2(\mu)$ **set-valued** !

Outline of the talk

Introduction to mean-field control

From differential inclusions to continuity inclusions

Main contributions

Continuity inclusions – *Definition & controlled systems*

Definition (Continuity inclusions) [B., Frankowska]

Let $(t, \mu) \in [0, T] \times \mathcal{P}_2(\mathbb{R}^d) \Rightarrow V(t, \mu) \subset L^2(\mu)$. Then, $\mu(\cdot)$ solves

$$\partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \quad (I)$$

if there exists $t \in [0, T] \mapsto \mathbf{v}(t) \in V(t, \mu(t))$ s.t.

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t)\mu(t)) = 0. \quad (E)$$

Certificate: Is this notion suited for control systems ? \rightsquigarrow Yes!

Theorem (Correspondence with control systems) [B., Frankowska]

Take $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times U \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is **Cauchy-Lip** and set

$$V(t, \mu) := v(t, \mu, U) \subset C^0(\mathbb{R}^d, \mathbb{R}^d).$$

Then $\mu(\cdot)$ solves (I) $\iff \mu(\cdot)$ solves (E) with $\mathbf{v}(t) := v(t, \mu(t), v(t))$

Continuity inclusions – *Definition & controlled systems*

Definition (Continuity inclusions) [B., Frankowska]

Let $(t, \mu) \in [0, T] \times \mathcal{P}_2(\mathbb{R}^d) \Rightarrow V(t, \mu) \subset L^2(\mu)$. Then, $\mu(\cdot)$ solves

$$\partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \quad (\text{I})$$

if there exists $t \in [0, T] \mapsto \mathbf{v}(t) \in V(t, \mu(t))$ s.t.

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t)\mu(t)) = 0. \quad (\text{E})$$

Certificate: Is this notion suited for control systems ? \rightsquigarrow Yes!

Theorem (Correspondence with control systems) [B., Frankowska]

Take $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times U \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is **Cauchy-Lip** and set

$$V(t, \mu) := v(t, \mu, U) \subset C^0(\mathbb{R}^d, \mathbb{R}^d).$$

Then $\mu(\cdot)$ solves (I) $\iff \mu(\cdot)$ solves (E) with $\mathbf{v}(t) := v(t, \mu(t), v(t))$

Continuity inclusions – Definition & controlled systems

Definition (Continuity inclusions) [B., Frankowska]

Let $(t, \mu) \in [0, T] \times \mathcal{P}_2(\mathbb{R}^d) \Rightarrow V(t, \mu) \subset L^2(\mu)$. Then, $\mu(\cdot)$ solves

$$\partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \quad (\text{I})$$

if there exists $t \in [0, T] \mapsto \mathbf{v}(t) \in V(t, \mu(t))$ s.t.

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t)\mu(t)) = 0. \quad (\text{E})$$

Certificate: Is this notion suited for control systems ? \rightsquigarrow Yes!

Theorem (Correspondence with control systems) [B., Frankowska]

Take $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times U \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is **Cauchy-Lip** and set

$$V(t, \mu) := v(t, \mu, U) \subset C^0(\mathbb{R}^d, \mathbb{R}^d).$$

Then $\mu(\cdot)$ solves (I) $\iff \mu(\cdot)$ solves (E) with $\mathbf{v}(t) := v(t, \mu(t), u(t))$

Continuity inclusions – Definition & controlled systems

Definition (Continuity inclusions) [B., Frankowska]

Let $(t, \mu) \in [0, T] \times \mathcal{P}_2(\mathbb{R}^d) \Rightarrow V(t, \mu) \subset L^2(\mu)$. Then, $\mu(\cdot)$ solves

$$\partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \quad (\text{I})$$

if there exists $t \in [0, T] \mapsto \mathbf{v}(t) \in V(t, \mu(t))$ s.t.

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t)\mu(t)) = 0. \quad (\text{E})$$

Certificate: Is this notion suited for control systems ? \rightsquigarrow Yes!

Theorem (Correspondence with control systems) [B., Frankowska]

Take $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times U \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is **Cauchy-Lip** and set

$$V(t, \mu) := v(t, \mu, U) \subset C^0(\mathbb{R}^d, \mathbb{R}^d).$$

Then $\mu(\cdot)$ solves (I) $\iff \mu(\cdot)$ solves (E) with $\mathbf{v}(t) := v(t, \mu(t), u(t))$

Continuity inclusions – Definition & controlled systems

Definition (Continuity inclusions) [B., Frankowska]

Let $(t, \mu) \in [0, T] \times \mathcal{P}_2(\mathbb{R}^d) \Rightarrow V(t, \mu) \subset L^2(\mu)$. Then, $\mu(\cdot)$ solves

$$\partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \quad (\text{I})$$

if there exists $t \in [0, T] \mapsto \mathbf{v}(t) \in V(t, \mu(t))$ s.t.

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t)\mu(t)) = 0. \quad (\text{E})$$

Certificate: Is this notion suited for control systems ? \rightsquigarrow **Yes!**

Theorem (Correspondence with control systems) [B., Frankowska]

Take $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times U \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is **Cauchy-Lip** and set

$$V(t, \mu) := v(t, \mu, U) \subset C^0(\mathbb{R}^d, \mathbb{R}^d).$$

Then $\mu(\cdot)$ solves (I) $\iff \mu(\cdot)$ solves (E) with $\mathbf{v}(t) := v(t, \mu(t), u(t))$

Continuity inclusions – Definition & controlled systems

Definition (Continuity inclusions) [B., Frankowska]

Let $(t, \mu) \in [0, T] \times \mathcal{P}_2(\mathbb{R}^d) \Rightarrow V(t, \mu) \subset L^2(\mu)$. Then, $\mu(\cdot)$ solves

$$\partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \quad (\text{I})$$

if there exists $t \in [0, T] \mapsto \mathbf{v}(t) \in V(t, \mu(t))$ s.t.

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t)\mu(t)) = 0. \quad (\text{E})$$

Certificate: Is this notion suited for control systems ? \rightsquigarrow **Yes!**

Theorem (Correspondence with control systems) [B., Frankowska]

Take $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times U \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is **Cauchy-Lip** and set

$$V(t, \mu) := v(t, \mu, U) \subset C^0(\mathbb{R}^d, \mathbb{R}^d).$$

Then $\mu(\cdot)$ solves (I) $\iff \mu(\cdot)$ solves (E) with $\mathbf{v}(t) := v(t, \mu(t), u(t))$

Continuity inclusions – *Definition & controlled systems*

Definition (Continuity inclusions) [B., Frankowska]

Let $(t, \mu) \in [0, T] \times \mathcal{P}_2(\mathbb{R}^d) \Rightarrow V(t, \mu) \subset L^2(\mu)$. Then, $\mu(\cdot)$ solves

$$\partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \quad (\text{I})$$

if there exists $t \in [0, T] \mapsto \mathbf{v}(t) \in V(t, \mu(t))$ s.t.

$$\partial_t \mu(t) + \operatorname{div}(\mathbf{v}(t)\mu(t)) = 0. \quad (\text{E})$$

Certificate: Is this notion suited for control systems ? \rightsquigarrow **Yes!**

Theorem (Correspondence with control systems) [B., Frankowska]

Take $v : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \times U \times \mathbb{R}^d \rightarrow \mathbb{R}^d$ is **Cauchy-Lip** and set

$$V(t, \mu) := v(t, \mu, U) \subset C^0(\mathbb{R}^d, \mathbb{R}^d).$$

Then $\mu(\cdot)$ solves (I) $\iff \mu(\cdot)$ solves (E) with $\mathbf{v}(t) := v(t, \mu(t), u(t))$

Continuity inclusions – *Compactness & Relaxation*

Theorem (Compactness of solution sets)[B., Frankowska]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **closed convex** images. Then for any μ^0 , the solution set

$$\mathcal{S}(\mu^0) := \left\{ \mu(\cdot) \text{ s.t. } \partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \mu(0) = \mu^0 \right\},$$

is **compact** in the $C^0 W_2$ -topology \rightsquigarrow **Existence of minimisers!**

Theorem (Relaxation)[B., Frankowska]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **closed** images. Then, $\mathcal{S}(\mu^0)$ is dense in the solution set of

$$\partial_t \mu(t) \in -\operatorname{div}(\overline{\operatorname{co}} V(t, \mu(t))\mu(t)), \quad \mu(0) = \mu^0,$$

where $\overline{\operatorname{co}} V(t, \mu)$ denotes the **closed convex hull** of $V(t, \mu)$.

Continuity inclusions – *Compactness & Relaxation*

Theorem (Compactness of solution sets)[B., Frankowska]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **closed convex** images. Then for any μ^0 , the solution set

$$\mathcal{S}(\mu^0) := \left\{ \mu(\cdot) \text{ s.t. } \partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \mu(0) = \mu^0 \right\},$$

is **compact** in the $C^0 W_2$ -topology \rightsquigarrow **Existence of minimisers!**

Theorem (Relaxation)[B., Frankowska]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **closed** images. Then, $\mathcal{S}(\mu^0)$ is dense in the solution set of

$$\partial_t \mu(t) \in -\operatorname{div}(\overline{\operatorname{co}} V(t, \mu(t))\mu(t)), \quad \mu(0) = \mu^0,$$

where $\overline{\operatorname{co}} V(t, \mu)$ denotes the closed convex hull of $V(t, \mu)$.

Continuity inclusions – *Compactness & Relaxation*

Theorem (Compactness of solution sets)[B., Frankowska]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **closed convex** images. Then for any μ^0 , the solution set

$$\mathcal{S}(\mu^0) := \left\{ \mu(\cdot) \text{ s.t. } \partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \mu(0) = \mu^0 \right\},$$

is **compact** in the $C^0 W_2$ -topology \rightsquigarrow **Existence of minimisers!**

Theorem (Relaxation)[B., Frankowska]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **closed** images. Then, $\mathcal{S}(\mu^0)$ is dense in the solution set of

$$\partial_t \mu(t) \in -\operatorname{div}(\overline{\operatorname{co}} V(t, \mu(t))\mu(t)), \quad \mu(0) = \mu^0,$$

where $\overline{\operatorname{co}} V(t, \mu)$ denotes the closed convex hull of $V(t, \mu)$.

Continuity inclusions – *Compactness & Relaxation*

Theorem (Compactness of solution sets)[B., Frankowska]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **closed convex** images. Then for any μ^0 , the solution set

$$\mathcal{S}(\mu^0) := \left\{ \mu(\cdot) \text{ s.t. } \partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \mu(0) = \mu^0 \right\},$$

is **compact** in the $C^0 W_2$ -topology \rightsquigarrow **Existence of minimisers!**

Theorem (Relaxation)[B., Frankowska]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **closed** images. Then, $\mathcal{S}(\mu^0)$ is dense in the solution set of

$$\partial_t \mu(t) \in -\operatorname{div}(\overline{\operatorname{co}} V(t, \mu(t))\mu(t)), \quad \mu(0) = \mu^0,$$

where $\overline{\operatorname{co}} V(t, \mu)$ denotes the closed convex hull of $V(t, \mu)$.

Continuity inclusions – *Compactness & Relaxation*

Theorem (Compactness of solution sets)[B., Frankowska]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **closed convex** images. Then for any μ^0 , the solution set

$$\mathcal{S}(\mu^0) := \left\{ \mu(\cdot) \text{ s.t. } \partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \mu(0) = \mu^0 \right\},$$

is **compact** in the $C^0 W_2$ -topology \rightsquigarrow **Existence of minimisers!**

Theorem (Relaxation)[B., Frankowska]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **closed** images. Then, $\mathcal{S}(\mu^0)$ is dense in the solution set of

$$\partial_t \mu(t) \in -\operatorname{div}(\overline{\operatorname{co}} V(t, \mu(t))\mu(t)), \quad \mu(0) = \mu^0,$$

where $\overline{\operatorname{co}} V(t, \mu)$ denotes the **closed convex hull** of $V(t, \mu)$.

Continuity inclusions – *Compactness & Relaxation*

Theorem (Compactness of solution sets)[B., Frankowska]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **closed convex** images. Then for any μ^0 , the solution set

$$\mathcal{S}(\mu^0) := \left\{ \mu(\cdot) \text{ s.t. } \partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \mu(0) = \mu^0 \right\},$$

is **compact** in the $C^0 W_2$ -topology \rightsquigarrow **Existence of minimisers!**

Theorem (Relaxation)[B., Frankowska]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **closed** images. Then, $\mathcal{S}(\mu^0)$ is dense in the solution set of

$$\partial_t \mu(t) \in -\operatorname{div}(\overline{\operatorname{co}} V(t, \mu(t))\mu(t)), \quad \mu(0) = \mu^0,$$

where $\overline{\operatorname{co}} V(t, \mu)$ denotes the **closed convex hull** of $V(t, \mu)$.

Continuity inclusions – Compactness & Relaxation

Theorem (Compactness of solution sets)[B., Frankowska]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **closed convex** images. Then for any μ^0 , the solution set

$$\mathcal{S}(\mu^0) := \left\{ \mu(\cdot) \text{ s.t. } \partial_t \mu(t) \in -\operatorname{div}(V(t, \mu(t))\mu(t)), \mu(0) = \mu^0 \right\},$$

is **compact** in the $C^0 W_2$ -topology \rightsquigarrow **Existence of minimisers!**

Theorem (Relaxation)[B., Frankowska]

Suppose that $V : [0, T] \times \mathcal{P}_c(\mathbb{R}^d) \rightrightarrows C^0(\mathbb{R}^d, \mathbb{R}^d)$ is **Cauchy-Lip**, with **closed** images. Then, $\mathcal{S}(\mu^0)$ is dense in the solution set of

$$\partial_t \mu(t) \in -\operatorname{div}(\overline{\operatorname{co}} V(t, \mu(t))\mu(t)), \quad \mu(0) = \mu^0,$$

where $\overline{\operatorname{co}} V(t, \mu)$ denotes the **closed convex hull** of $V(t, \mu)$.

Conclusion – *That's all folks!*

Thank you for your attention !

