H–E–S–M Comparative Research Map

Plant regeneration, nutrient biology, and human malnutrition: shared questions without unsupported clinical translation.

Prepared by

Kyle Harrison | 2639389 Alberta Ltd.
October 23, 2026 | Technical discussion paper | Version 1.0

Purpose

Connect the supplied plant-regeneration proposal to a carefully bounded discussion of human nutrient biology and malnutrition, using Orbital Architecture and HAGD-Sci to separate observations, analogies, hypotheses, and reviewed findings.

Research boundary

No pediatric doses, supplement ratios, schedules, treatment instructions, or claims of clinical benefit are proposed. Plant findings do not establish human interventions. A small dose is not evidence of safety or effectiveness.

Source status

Plant descriptions below summarize the author’s supplied proposal; its full bibliography has not been independently verified here. Human nutrient statements are linked to NIH Office of Dietary Supplements sources. Malnutrition scope is linked to WHO guidance. [P1; H1–H7]

1 | A comparative model, not an equivalence

Shared categories can organize questions; they do not establish identical mechanisms.

H | Hormonal or developmental signals

Plant profile: proposed growth-regulator and elicitor inputs. Human research profile: observed endogenous signaling and relevant physiological context. This mapping does not authorize hormone manipulation in children.

E | Environmental conditions

Plant profile: light, temperature, photoperiod, culture conditions, or spaceflight context. Human profile: contextual variables such as feeding environment, infection burden, and access to care, as specified by domain professionals. These are not interchangeable exposures.

S | State and transition

Plant profile: defined callus, embryo, shoot, propagule, or storage-organ endpoints. Human profile: explicitly defined nutritional and developmental observations. Recovery from malnutrition is not equivalent to plant organogenesis.

M | Nutrient availability

Plant profile: measured medium composition and nutrient availability. Human profile: dietary intake, absorption, health status, and clinically interpreted evidence. Do not transfer medium concentrations, nutrient ratios, or stage timing.

Proposed model

S(next) = F[S(current), H, E, M; domain-specific parameters]. This is a proposed modeling structure, not a proven equation. Nutrients and environmental conditions may act concurrently rather than in a fixed H → E → S → M sequence.

2 | Plant-domain evidence map

Preserve the seven channels, but keep their endpoints distinct.

Fruit channels | Apple, citrus, banana

Record species, genotype, explant, growth-regulator regime, environment, basal medium, timing, and outcome. The supplied proposal compares wound-induced organogenesis, bud regeneration, and somatic embryogenesis; each requires protocol-specific evidence. [P1]

Marine and medicinal channels

Marine macrophyte establishment and Catharanthus roseus growth or metabolite production are separate outcomes. Restoration, biomass, organogenesis, and alkaloid yield must not be combined as one regeneration-success measure. [P1]

Roots and space/BLSS channels

Storage-organ development and space-related growth studies require distinct endpoints and controls. Actual spaceflight, simulated microgravity, ground analogues, and complete life-support closure must be classified separately. [P1]

Status of the original scoring matrix

The 0–5 controllability, efficiency, sensitivity, robustness, and feasibility values should be labeled preliminary author-assigned scores until scoring rubrics, source selection, datasets, and reviewer agreement are supplied.

Minimum experimental record

Intervention and control; dose/concentration and units; timing; experimental unit; biological replication; batch; confounders; endpoint definition; analysis plan; adverse outcomes; negative and inconclusive findings.

3 | Nutrient comparisons: iron and zinc

Human functions are evidence anchors; cross-domain benefits remain hypotheses.

Iron | Plant-side proposal

The paper associates iron availability with plant metabolic processes, photosynthetic function, and culture outcomes. These plant claims require source-specific confirmation rather than being treated as one universal bottleneck. [P1]

Iron | Established human relevance

Iron contributes to hemoglobin and myoglobin and is required for growth and development. Iron deficiency and iron-deficiency anemia are clinically important, but nutritional requirements and assessment differ by age and context. [H1]

Iron | Valid comparison and boundary

Both profiles can examine how inadequate nutrient availability relates to measured biological function. Plant browning, tissue conversion, and human anemia are not equivalent endpoints. No pediatric iron dose or recovery schedule follows from the analogy.

Zinc | Established human relevance

Zinc has roles in enzyme activity, protein and DNA synthesis, immune function, wound healing, and growth. Deficiency can delay growth and cause diarrhea in children. [H2]

Zinc | Valid comparison and boundary

The paper’s plant-development questions may motivate comparative growth research. They do not show that extra zinc improves growth in an already sufficient child or justify growth-phase pulses. High zinc intake can interfere with copper status. [P1; H2; H3]

4 | Nutrient comparisons: copper, manganese, boron

Similar biological themes do not make all trace elements therapeutic targets.

Copper | Human evidence

Copper is an enzyme cofactor involved in functions including connective-tissue synthesis and iron metabolism. Copper deficiency is uncommon but can produce anemia and connective-tissue or bone abnormalities. Excess zinc can impair copper absorption. [H3]

Copper | Comparative boundary

The plant lignification versus human connective-tissue comparison is a mechanistic research analogy, not a validated fixed zinc:copper supplementation ratio. Record each mechanism and species-specific evidence separately.

Manganese | Human evidence

Manganese supports several enzyme systems, including manganese superoxide dismutase. Human deficiency is very rare. Essentiality does not establish a need for routine supplementation or prove a benefit in malnutrition recovery. [H4]

Manganese | Comparative boundary

Photosystem II oxygen production and mitochondrial antioxidant activity are different processes. Their shared use of manganese does not establish matching nutrient schedules or developmental checkpoints.

Boron | Critical exception

Boron is essential in plant biology as discussed in the supplied proposal, but NIH does not classify it as an essential human nutrient; human deficiency signs are not firmly established. Do not describe boron as a verified pediatric growth gate or treatment for malnutrition. [P1; H5]

5 | Malnutrition: keep the clinical scope intact

A trace-mineral analogy is not a complete model of nutritional recovery.

What WHO guidance actually covers

The WHO guideline addresses prevention and management of wasting and nutritional oedema in children under five. Its implementation guidance covers identification, assessment, referral, and nutritional, medical, and psychosocial management in different care settings. [H6; H7]

Why this matters to the framework

Micronutrient deficiency and acute malnutrition are not interchangeable categories. A shared mineral list cannot replace the broader clinical assessment, food and feeding support, medical care, and follow-up described in applicable guidance. [H6; H7]

Shared trait worth investigating

The defensible comparison is that biological outcomes can depend on nutrient availability and context. The research question is which relationships are supported within each organism and clinical population—not whether a plant protocol can become a child supplement plan.

Clinical data profile | For professional definition

Record age, clinical classification, dietary and feeding context, measurements selected by the care team, illness context, current treatment, source guideline, and follow-up. This document sets no diagnostic thresholds and supplies no treatment algorithm.

Safety boundary

Any real child with suspected malnutrition needs assessment by a qualified healthcare team. This research map must not delay referral or substitute for a recognized care protocol. [H6; H7]

6 | Put the model into Orbital Architecture

A proposed evidence workflow, not an established deployment.

Step 1 | DB1 governs research scope

Define plant versus human research boundaries, authorized uses, consent/ethics requirements where applicable, protocol versions, reviewer roles, and prohibited clinical extrapolations.

Step 2 | DB2 preserves evidence

Store separate plant and human evidence records. Include exact source, population/species, experiment or study type, conditions, measured outcomes, and limitations. Preserve provenance and contradictory results.

Step 3 | DB3 generates candidates

Produce comparative questions or candidate relationships. Label each as established evidence, interpretation, analogy, hypothesis, prediction, or unknown. Never present inferred human dosing as a verified result.

Step 4 | HAGD-Sci challenges the comparison

Qualified researchers check mechanistic fit, confounders, source support, testability, feasibility, and clinical boundaries. Accept, modify, reject, or escalate, preserving reasons and revisions.

Step 5 | DBX checks release prerequisites

Verify explicit evidence fields, authorization, domain labels, versions, and required approvals. A technical PASS is not proof of biological correctness or permission to prescribe.

Step 6 | DB4 stores scoped findings

Promote only reviewed claims with their limitations and permitted uses. Red-tag unsupported treatment implications without deleting independently supported nutrient functions. Record restoration, revision, and retirement decisions.

7 | A bounded validation roadmap

Start with evidence quality; do not start by treating children.

Phase 1 | Literature and claim audit

Verify the plant bibliography and human references. Break every paragraph into atomic claims. Identify direct support, indirect support, analogy, contradiction, and missing evidence. Remove unvalidated cross-domain scores and intervention claims from authoritative outputs.

Phase 2 | Plant-domain experimental validation

Select one tractable channel with qualified plant researchers. Define input windows, concurrent nutrient conditions, controls, endpoint definitions, replication, randomization, and analysis. Test a narrow falsifiable prediction before generalizing across channels.

Phase 3 | Human-domain question assessment

Pediatric nutrition and malnutrition specialists decide whether a specific comparative question adds value beyond existing human literature. A human study requires an independently justified design, appropriate review, safeguards, and approvals; plant success is not sufficient justification.

Phase 4 | Results-based integration

If evidence supports a finding, update its scope and review status. Preserve negative results and unresolved uncertainty. Do not auto-convert a research association into an intervention or autonomous clinical action.

Suggested metrics

Source-verification rate; claim–evidence alignment; mislabeled-analogy rate; reviewer agreement; revision traceability; protocol adherence; endpoint reproducibility. Clinical efficacy is not measured by how many claims a software workflow approves.

8 | Source ledger and final position

Selected evidence anchors; not an exhaustive systematic review.

P1 | Author-supplied research proposal

Kyle Harrison. Comparative Analysis of Regeneration Mechanisms: Sea → Jungle → Fruits → Roots → Space. Technical proposal supplied in this conversation. Plant mechanisms and channel scores require the independent audit described above.

H1–H3 | NIH nutrient fact sheets

H1 Iron: https://ods.od.nih.gov/factsheets/Iron-HealthProfessional/
H2 Zinc: https://ods.od.nih.gov/factsheets/Zinc-HealthProfessional/
H3 Copper: https://ods.od.nih.gov/factsheets/Copper-HealthProfessional/

H4–H5 | NIH nutrient fact sheets

H4 Manganese: https://ods.od.nih.gov/factsheets/Manganese-HealthProfessional/
H5 Boron: https://ods.od.nih.gov/factsheets/Boron-HealthProfessional/

H6–H7 | WHO malnutrition guidance

H6 WHO guideline on prevention and management of wasting and nutritional oedema in children under five (2023), publication identifier 9789240082830.
H7 WHO implementation guidance for inpatient and outpatient management of wasting and nutritional oedema (2026), publication identifier 9789240122840.
Available through WHO Publications. Sources reviewed October 23, 2026.

Final position

H–E–S–M can organize comparative questions. Human nutrition evidence defines human nutrient claims. Qualified professionals govern clinical interpretation. Orbital Architecture preserves the boundary between an interesting analogy and an evidence-supported, appropriately authorized use.