

MULTI-DIMENSIONAL FEMTOSECOND-LASER INDUCED DYNAMICS OF CO ON METALS:

ACCOUNTING FOR ELECTRONIC FRICTION AND SURFACE MOTION WITH COMBINED MODELS

Robert Scholz¹, Peter Saalfrank¹, Ivor Lončarić², Jean Cristophe Tremblay³, Gernot Fächsel³, and Gereon Floß¹

¹Institut für Chemie, Universität Potsdam, Karl-Liebknecht-Str. 24-25, 14476 Potsdam, Germany


²Ruder Bosković Institute, Div. of Theor. Physics, Bijenička cesta 54, 10000 Zagreb, Croatia

³Freie Universität, Inst. für Chemie und Biochemie, Takustr. 3, 14195 Berlin, Germany

Introduction


Motivation

- **Aim:** gain **precise understanding** of **adsorbate bonding** on metals
⇒ Important for **Catalysis**
- **Why femtosecond(fs)-lasers?**
 - **produce** non-equilibrium **2-T-states**
⇒ **different than** normal **heating**
 - further **tool** besides STM and scattering
 - **direct** future **applications** possible
⇒ “**femtochemistry**”[1]
- **Why CO/Ru(001) and CO/Cu(100)?**
 - both are well studied **model systems**
 - recently, interesting **fs-laser experiments**[2][3]
 - also, **ab-initio** based **6-dim. potentials** available [4][5]




How do fs-Lasers affect Adsorbate-Metal Systems?

- **fs-Laser** (purple wavy line) interacts with the **Ru (0001)** surface (blue line).
- **Desorption** (blue arrow) and **Diffusion** (yellow arrow) are shown.
- **Electron-phonon coupling** (1), **Electronic friction** (2), and **Phonon-adsorbate interaction** (3) are indicated.
- **Fermi-Dirac-Distribution** plot shows $f(E)$ vs E for T_{el} (low) and T_{ph} (high).
- **only electrons** of metal absorb laser
- **electron-hole pairs** thermalize fast
⇒ “**hot**” Fermi-Dirac-distribution
- electrons transfer energy to ion lattice, via **1) electron-phonon coupling**
- **equilibration within ps-timescale**
⇒ Thus, for few ps **two temperatures**:
 - T_{el} - electron temperature
 - T_{ph} - phonon temperature
- both can **couple** to adsorbed **molecule**
- low electron heat **capacity** ⇒ T_{el} higher




Results

Desorption (Data for Ru)



- desorption mainly during first 50 ps
- fluence dependence of desorption yield close to experiment
- no barrier in PMF


Vibrations (Data for Cu)



- Frequency-shift from time-resolved SFG
- Preliminary results from our dynamics


Models and Methods

Basis of the Dynamics: the Six-dimensional Potential Energy Surfaces (PES)[4][5]




- **precomputed** with DFT (GGA)
- all **six dimensions** of the adsorbate
- **analytical** ⇒ **very fast**
⇒ **many trajectories** possible
- but: surface frozen ⇒ **no phonons**

Two-Temperature Model (2TM)[6]




Electronic Friction: Langevin Dynamics[7] and Local Density Friction Approximation (LDFA)[8]



- the **Langevin** equation of motion, a **stochastic** differential equation:
- describes **movement** of CO and **interaction** with **electron-hole pairs** (friction and excitation)
- **Local Density Friction Approx.** (LDFA): *ab-initio* model that gives **friction coefficients** $\eta_{el,k}$
 - Atom k embedded in **free electron gas** with density of bare surface at current position \underline{r}_k
- **Random forces** $\underline{R}_{el,k}$: white noise, **dependent** on both $\eta_{el,k}$ (from LDFA) and T_{el} (from 2TM)

Inclusion of Phonons: Generalized Langevin Oscillator(GLO)-model[9]



Conclusions

- **system:** CO on Ru(001) or Cu(100)
- **full-dimensional (6D)** simulation of dynamics driven by **fs-laser irradiation**
- **first principles**, no “free” parameters
- accounting for **1) electronic friction** and excitation by hot electron-hole pairs (via LDFA and the Langevin approach) and **2) substrate motion** (via GLO)
- detailed time- and space-resolved insights

Outlook

- better description of electronic friction: η_{el} as $f(T_{el})$ and going **beyond LDFA**
⇒ Long term goal: **friction tensors**[10]
- further **2TM improvements**, consider **non-equilibrium lattice (NLM)**[11]
- simulate **other coverages** and interaction between adsorbate molecules
⇒ bigger super cells needed
- go to **larger** or more complex **systems** hydrocarbons; coadsorbate CO + H or O

References

- [1] M. Bonn, S. Funk, Ch. Hess, D.N. Denzler *et al.*, *Science* **285**, 1042 (1999).
- [2] M. Dell’Angela, T. Anniyev, M. Beye, R. Coffee *et al.*, *Science* **339**, 1302 (2013).
- [3] K. Inoue, K. Watanabe, T. Sugimoto *et al.*, *Phys.Rev.Lett.* **117**, 186101 (2016).
- [4] G. Fächsel, J.C. Tremblay, and P. Saalfrank, *J.Chem.Phys.* **141**, 094704 (2014).
- [5] R. Marquardt, F. Cuvelier, R.O. Olsen *et al.*, *J.Chem.Phys.* **132**, 074108 (2010).
- [6] S.I. Anisimov, B.L. Kapeliovich and T.L. Perelman, *Sov.Phys.-JETP* **39**, 375 (1974).
- [7] M. Head-Gordon and J.C. Tully, *J.Chem.Phys.* **103**, 10137 (1995).
- [8] J.I. Juaristi, M. Alducin, R. Díez Muiño *et al.*, *Phys.Rev.Lett.* **100**, 116102 (2008).
- [9] H.F. Busnengo, M.A. Di Césare, W. Dong *et al.*, *Phys.Rev.B* **72**, 125411 (2005).
- [10] R. J. Maurer, M. Askerka, V. S. Batista, J. C. Tully *Phys.Rev.B* **94**, 115432 (2016).
- [11] L. Waldecker, R. Bertoni, R. Ernstorfer, J. Vorberger *Phys.Rev.X* **6**, 021003 (2016).