

FEMTOSECOND-LASER INDUCED DYNAMICS OF CO ON Ru(0001):

NEW INSIGHTS FROM A HOT-ELECTRON, ELECTRONIC FRICTION MODEL INCLUDING SURFACE MOTION

Robert Scholz^{1,2}, Gereon Floß¹, Peter Saalfrank¹, Gernot Fuchsels³, Ivor Lončarić⁴, and J. I. Juaristi^{4,5,6}

¹Institut für Chemie, Universität Potsdam, Karl-Liebknecht-Str. 24-25, D-14476 Potsdam, Germany

²Fritz-Haber-Institut der Max-Planck-Gesellschaft, Faradayweg 4-6, D-14195 Berlin, Germany

³Universiteit Leiden, Gorlaeus Laboratories, Einsteinweg 55, 2333 Leiden, The Netherlands

⁴Centro de Física de Materiales CFM/MPC (CSIC-UPV/EHU), Paseo Manuel de Lardizabal 5, 20018 Donostia-San Sebastián, Spain


⁵Departamento de Física de Materiales, Facultad de Químicas, Universidad del País Vasco (UPV/EHU), Apartado 1072, 20080 San Sebastián, Spain

⁶Donostia International Physics Center DIPC, P. Manuel de Lardizabal 4, 20018 San Sebastián, Spain

Introduction




Motivation

- research on small molecules adsorbed to metals is important for:
 - catalytic applications
 - fundamental understanding of bonding
- femtosecond(fs)-lasers \Rightarrow very valuable tool
 - allow for investigations on small timescales
 - open up new processes compared to heating or “normal” lasers \Rightarrow femtochemistry, e.g. [1]:
 - may enable specific control over catalytic reactions \Rightarrow photocatalysis
- specific motivation for system CO/Ru(0001)
 - experimentally well studied regarding fs-laser irradiation, e.g. [2, 3]
 - fulldimensional *ab-initio* potential recently developed in our group[4]
 - details of this indicate interpretation of experiment [3] may be wrong



How does fs-laser-irradiation affect metal surfaces?


- fs-Laser** \rightarrow **Desorption** **Diffusion (and possibly Reactions)**
- ① **Electron-phonon coupling**
② **Electronic friction**
③ **Phonon-adsorbate interaction**
- metals: ion lattice plus quasi-free electron gas
 - visible light is absorbed only by the electrons
 - produced electron-hole pairs thermalize quickly \Rightarrow “hot” Fermi-Dirac-distribution (after ~ 10 fs)
 - electrons transfer part of energy to ion lattice, via ① **electron-phonon coupling** (phonons = lattice vibrations; quasi-particles)
 - electrons couple to phonons as their fast movement causes “shockwaves” in ion lattice
 - equilibration process completes after ~ 1 ps
- \Rightarrow Thus, with fs-lasers, two different temperatures:
- T_{el} - electron temperature
 - T_{ph} - phonon temperature
- electron-hole pairs and phonons both couple to adsorbed molecule



Models and Methods

Six-dimensional Potential Energy Surface (6D PES)[4]

- Basis for dynamics: precomputed PES from DFT (rPBE + D2)
 - all 6 dimensions of the adsorbate
 - analytical PES and gradients \Rightarrow very fast
- \Rightarrow number and length of trajectories can be large
- downsides:
 - surface atoms frozen \Rightarrow no phonons
 - had to be constructed first



Two-Temperature Model (2TM)[5]


- describes interaction of metal with laser, using two differential equations:

$$C_{el} \frac{\partial T_{el}}{\partial t} = \frac{\partial}{\partial z} \kappa \frac{\partial T_{el}}{\partial z} - g(T_{el} - T_{ph}) + S(z, t),$$

$$C_{ph} \frac{\partial T_{ph}}{\partial t} = g(T_{el} - T_{ph}).$$

\Rightarrow get T_{el} and T_{ph} as $f(z, t)$ from laser parameters and material properties:


- laser wavelength λ (affects penetration depth into material)
- electron and phonon heat capacities C_{el} and C_{ph}
- (effective) absorbed fluence F (energy/area)
- electron heat conductivity κ
- pulse duration τ (all three appear in the “source term” $S(z, t)$)
- electron-phonon coupling constant g



Electronic Friction: Langevin Dynamics[6] and Local Density Friction Approximation (LDFA)[7]


- Langevin equation of motion, a stochastic differential equation:

$$m_k \frac{d^2 \underline{r}_k}{dt^2} = \underbrace{-\nabla_k V(\underline{r}_1, \underline{r}_2)}_{\text{Force on Atom } k} - \underbrace{\eta_{el,k}(\underline{r}_k) \frac{d \underline{r}_k}{dt}}_{\text{Friction force slows movement}} + \underbrace{\underline{R}_{el,k}(t)}_{\text{Random force from e-h pairs}}.$$
- describes movement of CO on the PES and interaction with electron-hole pairs (friction and excitation)
- Local Density Friction Approx. (LDFA): most simple model to calculate friction coefficients $\eta_{el,k}$
 - Atom k embedded in free electron gas with density of bare surface at current position \underline{r}_k
- Random forces $\underline{R}_{el,k}$: gaussian white noise, dependent on both $\eta_{el,k}$ (from LDFA) and T_{el} (from 2TM)
 - justified by the 2. fluctuation dissipation theorem [11], which relates friction and thermal movement




Inclusion of Phonons: Generalized Langevin Oscillator(GLO)-model[8, 9, 10]

- influence of phonons modeled in an effective way (augments frozen surface)
- entire surface understood as 3D oscillator (coordinates \underline{r}_s , mass $m_s = m_{Ru}$)
- coupling to molecule via shifting: $V_{GLO}(\underline{r}_C, \underline{r}_O; \underline{r}_s) = V(\underline{r}_C - \underline{r}_s, \underline{r}_O - \underline{r}_s)$
- additionally coupled to ghost oscillator \underline{r}_g to model influence of the bulk
 - ghost oscillator is subject to friction η_{ph} and random forces $\underline{R}_{ph}(T_{ph})$




Results

Desorption



Diffusion



Conclusions

- 6D Langevin dynamics of CO on Ru(0001)
- based on first principles, no “free” parameters
- accounting for (via LDFA) electronic friction, hot electron excitation and (via GLO) substrate motion
- allows for detailed time- and space-resolved insights
- no physisorbed state, molecules desorb directly

Outlook

- calculation of RIXS spectra currently performed
- employ better electronic friction (beyond LDFA)
- enhance 2TM with electron-electron-scattering
- simulate other coverages (here only 0.25ML)
- include interaction between adsorbate molecules

References

- [1] M. Bonn, S. Funk, Ch. Hess, D.N. Denzler et al., *Science* **285**, 1042 (1999).
 [2] S. Funk, M. Bonn, D.N. Denzler, C. Hess et al., *J.Chem.Phys.* **112**, 9888 (2000).
 [3] M. Dell’Angela, T. Anniyev, M. Beye, R. Coffee et al., *Science* **339**, 1302 (2013).
 [4] G. Fuchsels, J.C. Tremblay, and P. Saalfrank, *J.Chem.Phys.* **141**, 094704 (2014).
 [5] S.I. Anisimov, B.L. Kapeliovich and T.L. Perelman, *Sov.Phys.-JETP* **39**, 375 (1974).
 [6] M. Head-Gordon and J.C. Tully, *J.Chem.Phys.* **103**, 10137 (1995).
 [7] J.I. Juaristi, M. Alducin, R. Diez Muiño, et al., *Phys.Rev.Lett.* **100**, 116102 (2008).
 [8] S.A. Adelman and J.D. Doll, *J.Chem.Phys.* **64**, 2375 (1976).
 [9] J.C. Tully, *J.Chem.Phys.* **73**, 1975 (1980).
 [10] H.F. Busnengo, M.A. Di Césare, W. Dong et al., *Phys.Rev.B* **72**, 125411 (2005).
 [11] R. Kubo, *Rep.Prog.Phys.* **29**, 255 (1966).