

FEMTOSECOND-LASER INDUCED DYNAMICS OF CO ON Ru(0001): NEW INSIGHTS FROM A HOT-ELECTRON, ELECTRONIC FRICTION MODEL INCLUDING SURFACE MOTION

Robert Scholz^{1,2}, Gereon Floß¹, Peter Saalfrank¹, Gernot Fuchsels³, Ivor Lončarić⁴, and J. I. Juaristi^{4,5,6}

¹Institut für Chemie, Universität Potsdam, Karl-Liebknecht-Str. 24-25, D-14476 Potsdam, Germany

²Fritz-Haber-Institut der Max-Planck-Gesellschaft, Faradayweg 4-6, D-14195 Berlin, Germany

³Universiteit Leiden, Gorlaeus Laboratories, Einsteinweg 55, 2333 Leiden, The Netherlands

⁴Centro de Física de Materiales CFM/MPC (CSIC-UPV/EHU), Paseo Manuel de Lardizabal 5, 20018 Donostia-San Sebastián, Spain

⁵Departamento de Física de Materiales, Facultad de Químicas, Universidad del País Vasco (UPV/EHU), Apartado 1072, 20080 San Sebastián, Spain


⁶Donostia International Physics Center DIPC, P. Manuel de Lardizabal 4, 20018 San Sebastián, Spain

Introduction


Motivation

- research on small molecules adsorbed to metals is important for:
 - catalytic applications
 - fundamental understanding of bonding
- femtosecond(fs)-lasers are a valuable tool for such research as they
 - allow for investigations on small timescales
 - open up new processes compared to heating (femtochemistry)
 - may enable specific control over catalytic reactions (photocatalysis)
- specific motivation for system CO/Ru(0001)
 - experimentally well studied regarding fs-laser irradiation, e.g. [1, 2]
 - fulldimensional *ab-initio* potential recently developed in our group[3]
 - details of this indicate interpretation of experiment [2] may be wrong

How does fs-laser-irradiation affect metal surfaces?



- metals: ion lattice plus quasi-free electron gas
- visible light is absorbed only by the electrons
- produced electron hole pairs thermalize quickly \Rightarrow “hot” Fermi-Dirac-distribution (after ~ 10 fs)
- electrons transfer part of energy to ion lattice, via ① **electron-phonon coupling** (phonons = lattice vibrations; quasi-particles)
 - electrons couple to phonons as their fast movement causes “shockwaves” in ion lattice
 - equilibration process completes after ~ 1 ps



\Rightarrow Thus, with fs-lasers, two different temperatures:


- T_{el} - electron temperature
- T_{ph} - phonon temperature

- can be simulated using a Two-Temperature Model (2TM) [4] (see right)

Models and Methods

Six-dimensional Potential Energy Surface (6D PES)

- Basis for dynamics: precomputed PES from DFT (rPBE + D2)
 - all 6 dimensions of the adsorbate
 - analytical PES and gradients \Rightarrow very fast \Rightarrow number and length of trajectories can be large
 - downside: surface atoms frozen \Rightarrow no phonons



Two-Temperature Model (2TM) [4]

- consists of two coupled differential equations: $C_{el} \frac{\partial T_{el}}{\partial t} = \frac{\partial}{\partial z} \kappa \frac{\partial T_{el}}{\partial z} - g(T_{el} - T_{ph}) + S(z, t)$, $C_{ph} \frac{\partial T_{ph}}{\partial t} = g(T_{el} - T_{ph})$
- describes interaction of the metal surface and laser
- calculates T_{el} and T_{ph} as $f(z, t)$ from laser parameters and material properties:
 - laser wavelength λ (affects penetration depth into material)
 - electron and phonon heat capacities C_{el} and C_{ph}
 - (effective) absorbed fluence F (energy/area)
 - electron heat conductivity κ
 - pulse duration τ (all three appear in the “source term” $S(z, t)$)
 - electron-phonon coupling constant g

Electronic Friction: LDFA and Langevin Dynamics

Inclusion of Phonons: GLO-model

References

- [1] S. Funk, M. Bonn, D. N. Denzler, C. Hess et al., *J. Chem. Phys.* **112**, 9888 (2000).
- [2] M. Dell’Angela, T. Anniyev, M. Beye et al., *Science* **339**, 1302 (2013).
- [3] G. Fuchsels, J. C. Tremblay, and P. Saalfrank, *J. Chem. Phys.* **141**, 094704 (2014).

- [4] S. I. Anisimov, B. L. Kapeliovich, and T. L. Perel’man, *Sov. Phys.-JETP* **39**, 375 (1974).
- [5] J. I. Juaristi, M. Alducin, R. Díez Muiño, et al., *Phys. Rev. Lett.* **100**, 116102 (2008).
- [6] S. A. Adelman and J. D. Doll, *J. Chem. Phys.* **64**, 2375 (1976).
- [7] J. C. Tully, *J. Chem. Phys.* **73**, 1975 (1980).
- [8] H. F. Busnengo, M. A. Di Césare, W. Dong, and A. Salin, *Phys. Rev. B* **72**, 125411 (2005).