

FEMTOSECOND-LASER INDUCED DYNAMICS OF CO ON Ru(0001): NEW INSIGHTS FROM A HOT-ELECTRON, ELECTRONIC FRICTION MODEL INCLUDING SURFACE MOTION

Robert Scholz^{1,2}, Gereon Floß¹, Peter Saalfrank¹, Gernot Fuchs³, Ivor Lončarić⁴, and J. I. Juaristi^{4,5,6}

¹Institut für Chemie, Universität Potsdam, Karl-Liebknecht-Str. 24-25, D-14476 Potsdam, Germany

²Fritz-Haber-Institut der Max-Planck-Gesellschaft, Faradayweg 4-6, D-14195 Berlin, Germany

³Universiteit Leiden, Gorlaeus Laboratories, Einsteinweg 55, 2333 Leiden, The Netherlands

⁴Centro de Física de Materiales CFM/MPC (CSIC-UPV/EHU), Paseo Manuel de Lardizabal 5, 20018 Donostia-San Sebastián, Spain

⁵Departamento de Física de Materiales, Facultad de Químicas, Universidad del País Vasco (UPV/EHU), Apartado 1072, 20080 San Sebastián, Spain





⁶Donostia International Physics Center DIPC, P. Manuel de Lardizabal 4, 20018 San Sebastián, Spain

Introduction

Motivation

- research on small molecules adsorbed to metals is important for:
 - catalytic applications
 - fundamental understanding of bonding
- femtosecond(fs)-lasers are a valuable tool for such research as they
 - allow for investigations on small timescales
 - open up new processes compared to heating (femtophotochemistry)
 - may enable specific control over catalytic reactions (photocatalysis)


How does fs-laser-irradiation affect metal surfaces?

- fs-Laser** **Desorption**
Diffusion (and possibly Reactions)
- ① Electron-phonon coupling**
② Electronic friction
③ Phonon-adsorbate interaction
- metals: ion lattice plus quasi-free electron gas
visible light is absorbed only by the electrons
produced electron hole pairs thermalize quickly
⇒ “hot” Fermi-Dirac-distribution (after ~10 fs)
electrons transfer part of energy to ion lattice, via **① electron-phonon coupling** (phonons = lattice vibrations; quasi-particles)
– electrons couple to phonons as their fast movement causes “shockwaves” in ion lattice
– equilibration process completes after ~1 ps
⇒ Thus, with fs-lasers, two different temperatures:
– T_{el} - electron temperature
– T_{ph} - phonon temperature
can be simulated using a Two-Temperature Model (2TM) [1] (see right)
- 
- 
- 
- 

Models and Methods

Six-dimensional Potential Energy Surface (6D PES)

- Basis for dynamics: precomputed Potential Energy Surface (PES)
 - all 6 dimensions of the adsorbate
 - analytical PES and gradients ⇒ very fast
⇒ number and length of trajectories can be large
 - downside: surface atoms frozen ⇒ no phonons



Two-Temperature Model (2TM) [1]

- consists of two coupled differential equations:
$$C_{\text{el}} \frac{\partial T_{\text{el}}}{\partial t} = \frac{\partial}{\partial z} \kappa \frac{\partial}{\partial z} T_{\text{el}} - g(T_{\text{el}} - T_{\text{ph}}) + S(z, t),$$
$$C_{\text{ph}} \frac{\partial T_{\text{ph}}}{\partial t} = g(T_{\text{el}} - T_{\text{ph}}).$$
- describes interaction of the metal surface and laser
- calculates T_{el} and T_{ph} from laser parameters and material properties e.g.:
 - laser wavelength λ (affects penetration depth into material)
 - electron and phonon heat capacities C_{el} and C_{ph}
 - (effective) absorbed fluence F (energy/area)
 - electron heat conductivity κ
 - pulse duration τ (all three appear in the “source term” $S(z, t)$)
 - electron-phonon coupling constant g

Electronic Friction: LDFA and Langevin Dynamics

Inclusion of Phonons: GLO-model

References

[1] S. I. Anisimov, B. L. Kapeliovich, and T. L. Perel'man, *Sov. Phys.-JETP* **39**, 375 (1974).

[2] M. Dell'Angela, T. Anniyev, M. Beye et al., *Science* **339**, 1302 (2013).