

FEMTOSECOND-LASER INDUCED DYNAMICS OF CO ON Ru(0001): NEW INSIGHTS FROM A HOT-ELECTRON, ELECTRONIC FRICTION MODEL INCLUDING SURFACE MOTION

Robert Scholz^{1,2}, Gereon Floß¹, Peter Saalfrank¹, Gernot Fuchsels³, Ivor Lončarić⁴, and J. I. Juaristi^{4,5,6}

¹Institut für Chemie, Universität Potsdam, Karl-Liebknecht-Str. 24-25, D-14476 Potsdam, Germany

²Fritz-Haber-Institut der Max-Planck-Gesellschaft, Faradayweg 4-6, D-14195 Berlin, Germany

³Universiteit Leiden, Gorlaeus Laboratories, Einsteinweg 55, 2333 Leiden, The Netherlands

⁴Centro de Física de Materiales CFM/MPC (CSIC-UPV/EHU), Paseo Manuel de Lardizabal 5, 20018 Donostia-San Sebastián, Spain

⁵Departamento de Física de Materiales, Facultad de Químicas, Universidad del País Vasco (UPV/EHU), Apartado 1072, 20080 San Sebastián, Spain




⁶Donostia International Physics Center DIPC, P. Manuel de Lardizabal 4, 20018 San Sebastián, Spain

Introduction

Motivation



- research on small molecules adsorbed to metals is important for:
 - catalytic applications
 - fundamental understanding of bonding
- femtosecond(fs)-lasers are a valuable tool for such research as they
 - allow for investigations on small timescales
 - open up new processes compared to heating (femtochemistry)
 - may enable specific control over catalytic reactions (photocatalysis)
- specific motivation for system CO/Ru(0001)
 - experimentally well studied regarding fs-laser irradiation, e.g. [1, 2]
 - fulldimensional *ab-initio* potential recently developed in our group[3]
 - details of this indicate interpretation of experiment [2] may be wrong

How does fs-laser-irradiation affect metal surfaces?

- 
- fs-Laser
- Desorption
- Diffusion (and possibly Reactions)
- ① Electron-phonon coupling
- ② Electronic friction
- ③ Phonon-adsorbate interaction
- Ru (0001)
- T_{el} T_{ph}
- metals: ion lattice plus quasi-free electron gas
 - visible light is absorbed only by the electrons
 - produced electron-hole pairs thermalize quickly \Rightarrow “hot” Fermi-Dirac-distribution (after ~ 10 fs)
 - electrons transfer part of energy to ion lattice, via ① **electron-phonon coupling** (phonons = lattice vibrations; quasi-particles)
 - electrons couple to phonons as their fast movement causes “shockwaves” in ion lattice
 - equilibration process completes after ~ 1 ps
- \Rightarrow Thus, with fs-lasers, two different temperatures:
- T_{el} - electron temperature
 - T_{ph} - phonon temperature
- can be simulated using a Two-Temperature Model (2TM)[4] (see right)
- 
- 

Models and Methods

Six-dimensional Potential Energy Surface (6D PES)[3]

- Basis for dynamics: precomputed PES from DFT (rPBE + D2)
 - all 6 dimensions of the adsorbate
 - analytical PES and gradients \Rightarrow very fast
- \Rightarrow number and length of trajectories can be large
- downsides:
 - surface atoms frozen \Rightarrow no phonons
 - had to be constructed first
- 
- 


Two-Temperature Model (2TM)[4]

- describes interaction of metal with laser, using two differential equations:


$$C_{el} \frac{\partial T_{el}}{\partial t} = \frac{\partial}{\partial z} \kappa \frac{\partial T_{el}}{\partial z} - g(T_{el} - T_{ph}) + S(z, t),$$

$$C_{ph} \frac{\partial T_{ph}}{\partial t} = g(T_{el} - T_{ph}).$$

\Rightarrow get T_{el} and T_{ph} as $f(z, t)$ from laser parameters and material properties:


- laser wavelength λ (affects penetration depth into material)
 - (effective) absorbed fluence F (energy/area)
 - pulse duration τ (all three appear in the “source term” $S(z, t)$)
 - electron and phonon heat capacities C_{el} and C_{ph}
 - electron heat conductivity κ
 - electron-phonon coupling constant g
- 

Electronic Friction: Langevin Dynamics[5] and Local Density Friction Approximation (LDFA)[6]

- 
- Langevin equation of motion, a stochastic differential equation:

$$m_k \frac{d^2 \underline{r}_k}{dt^2} = \underbrace{-\nabla_k V(\underline{r}_1, \underline{r}_2)}_{\text{Force on Atom } k} - \underbrace{\eta_{el,k}(\underline{r}_k) \frac{d \underline{r}_k}{dt}}_{\text{Friction force slows movement}} + \underbrace{\underline{R}_{el,k}(t)}_{\text{Random force from e-h pairs}}.$$
 - describes movement of CO on the PES and interaction with electron-hole pairs (friction and excitation)
 - Local Density Friction Approx. (LDFA): most simple model to calculate friction coefficients $\eta_{el,k}$
 - Atom k embedded in free electron gas with density of bare surface at current position \underline{r}_k
 - Random forces $\underline{R}_{el,k}$: gaussian white noise, dependent on both $\eta_{el,k}$ and T_{el}

Inclusion of Phonons: Generalized Langevin Oscillator(GLO)-model[7, 8, 9]



References

- [1] S. Funk, M. Bonn, D. N. Denzler, C. Hess, M. Wolf and G. Ertl, *J. Chem. Phys.* **112**, 9888 (2000).
- [2] M. Dell’Angela, T. Anniyev, M. Beye, R. Coffee, A. Föhlisch et al., *Science* **339**, 1302 (2013).
- [3] G. Fuchsels, J. C. Tremblay, and P. Saalfrank, *J. Chem. Phys.* **141**, 094704 (2014).
- [4] S. I. Anisimov, B. L. Kapeliovich, and T. L. Perel’man, *Sov. Phys.-JETP* **39**, 375 (1974).

- [5] M. Head-Gordon and J. C. Tully, *J. Chem. Phys.* **103**, 10137 (1995).

- [6] J. I. Juaristi, M. Alducin, R. Díez Muiño, H. F. Busnengo and A. Salin, *Phys. Rev. Lett.* **100**, 116102 (2008).

- [7] S. A. Adelman and J. D. Doll, *J. Chem. Phys.* **64**, 2375 (1976).

- [8] J. C. Tully, *J. Chem. Phys.* **73**, 1975 (1980).

- [9] H. F. Busnengo, M. A. Di Césare, W. Dong, and A. Salin, *Phys. Rev. B* **72**, 125411 (2005).