

Laser-driven dynamics of CO on Ru(0001)

a computational study using electronic friction (MDEF) and
the generalized Langevin oscillator (GLO)

Robert Scholz

Institut für Chemie
Universität Potsdam

30. November 2016

1 Introduction

- Motivation - in general and system specific
- First impressions of fs-laser-driven dynamics

2 Models and methods

- Foundations: 6D potential and two-temperature model
- Electronic friction: non-adiabatic coupling approximated
- The generalized Langevin oscillator (GLO)
- “Half-time”: short summary (and maybe time for a few questions)

3 Results and discussion

- Laser-driven diffusion
- Laser-driven desorption
- Physisorbed precursor states?

4 Summary and Outlook

1 Introduction

- Motivation - in general and system specific
- First impressions of fs-laser-driven dynamics

2 Models and methods

- Foundations: 6D potential and two-temperature model
- Electronic friction: non-adiabatic coupling approximated
- The generalized Langevin oscillator (GLO)
- “Half-time”: short summary (and maybe time for a few questions)

3 Results and discussion


- Laser-driven diffusion
- Laser-driven desorption
- Physisorbed precursor states?

4 Summary and Outlook

General motivation

Why investigate fs-laser-driven surface dynamics?

- gain fundamental understanding of adsorbate bonding
⇒ additional tool besides scattering experiments and STM
- possible direct application in catalysis: “femtochemistry”
⇒ new reaction pathways opened up by fs-lasers



CO/O-coadsorbate @ Ru(0001)

M. Bonn *et al.*, Science 1999

Specific motivation for investigating CO/Ru(0001)

CO/Ru(0001) system important for catalysis, e.g.:



- Fischer-Tropsch synthesis
- exhaust gas converters (systems with similar metals)

Experimentally well studied system

- especially regarding fs-laser irradiation:
 - e.g. Bonn *et al.*, *Science* 1999; Funk *et al.*, *JChemPhys* 2000
 - (both Ertl group ⇒ chemistry Nobel prize 2007).
- desorption kinetics: large prefactors ⇒ still not understood
- recently, time resolved x-ray spectra (XAS and XES)
⇒ “movie” of changes in orbital density of states
 - Dell'Angela *et al.*, *Science* 2013

Further specific motivation for investigating CO/Ru

Füchsel *et al.*, *JChemPhys* 2014



Important prior theory work was done at our group

Füchsel *et al.*, *JChemPhys* 2014

- Development of a potential energy surface (PES)
 - from over 90 000 DFT points!
 - all 6 dimensions of the adsorbate
 - very fast because preconstructed
- ⇒ **enables large-scale dynamics!**

Outline

1 Introduction

- Motivation - in general and system specific
- First impressions of fs-laser-driven dynamics

2 Models and methods


- Foundations: 6D potential and two-temperature model
- Electronic friction: non-adiabatic coupling approximated
- The generalized Langevin oscillator (GLO)
- “Half-time”: short summary (and maybe time for a few questions)

3 Results and discussion

- Laser-driven diffusion
- Laser-driven desorption
- Physisorbed precursor states?


4 Summary and Outlook

What happens after fs-laser excitation of the metal?



Coupling between 3 kinds of degrees of freedom:

- electron gas (T_{el})
 - initially absorbs laser energy
 - low heat capacity \Rightarrow high T_{el} ($\approx 5\text{-}10 \text{ kK}$)
- lattice vibrations (T_{ph})
 - thermalization with electrons: ps time scale \Rightarrow fs-laser causes two distinct temperatures!
- adsorbate movement (T_{ads})



Details of the time-resolved x-ray experiment

Dell'Angela *et al.*, *Science* 2013 (experimental part by Nilsson group, SLAC/LCLS, Stanford)


What was done?

- pump: *vis-fs-laser*
- probe: x-ray free e^- laser
 - K-edge of O-atom

What is observed?



- orbital density of states at O
- energies shift towards gas-phase values of CO
- intensities change
 - $2\tilde{\pi}^*$ \Rightarrow increase by $\sim 30\%$
 - \tilde{d}_π \Rightarrow decrease by $\sim 30\%$
 - participant peak appears

⇒ physisorbed precursor(?)



Details of the accompanying theory

still Dell'Angela *et al.*, *Science* 2013 (theory part by Nørskov group, SUNCAT, Stanford)



Dell'Angela *et al.*, *Science* 2013

Scholz *et al.*, *PhysRevB* 2016

Outline

1 Introduction

- Motivation - in general and system specific
- First impressions of fs-laser-driven dynamics

2 Models and methods


- Foundations: 6D potential and two-temperature model
- Electronic friction: non-adiabatic coupling approximated
- The generalized Langevin oscillator (GLO)
- “Half-time”: short summary (and maybe time for a few questions)

3 Results and discussion

- Laser-driven diffusion
- Laser-driven desorption
- Physisorbed precursor states?


4 Summary and Outlook

Ab-initio-based, full-dimensional (6D) potential energy surface (PES)



Two-temperature model (TTM)

$$C_{\text{el}} \frac{\partial T_{\text{el}}}{\partial t} = \frac{\partial}{\partial z} \kappa \frac{\partial}{\partial z} T_{\text{el}} - g(T_{\text{el}} - T_{\text{ph}}) + S(z, t),$$
$$C_{\text{ph}} \frac{\partial T_{\text{ph}}}{\partial t} = g(T_{\text{el}} - T_{\text{ph}}).$$



Outline

1 Introduction

- Motivation - in general and system specific
- First impressions of fs-laser-driven dynamics

2 Models and methods


- Foundations: 6D potential and two-temperature model
- **Electronic friction: non-adiabatic coupling approximated**
- The generalized Langevin oscillator (GLO)
- “Half-time”: short summary (and maybe time for a few questions)

3 Results and discussion

- Laser-driven diffusion
- Laser-driven desorption
- Physisorbed precursor states?

4 Summary and Outlook

Different approaches to describe non-adiabatic coupling



Langevin Dynamics

$$\underbrace{m_k \frac{d^2 \underline{r}_k}{dt^2}}_{\text{Force on Atom } k} = \underbrace{-\nabla_k V(\underline{r}_1, \underline{r}_2)}_{\text{Force due to PES}} - \underbrace{\eta_{\text{el},k}(\underline{r}_k) \frac{d\underline{r}_k}{dt}}_{\text{Friction force slows movement}} + \underbrace{R_{\text{el},k}(t)}_{\text{Random force from e-h pairs}}.$$

- $R_{\text{el},k}(t)$ = Gaussian white noise
 - describes excitation by hot electron-hole pairs
 - dependent on: $\eta_{\text{el},k}(\underline{r}_k)$ and T_{el}

Local density friction approx. plus independent atoms

1 Introduction

- Motivation - in general and system specific
- First impressions of fs-laser-driven dynamics

2 Models and methods

- Foundations: 6D potential and two-temperature model
- Electronic friction: non-adiabatic coupling approximated
- **The generalized Langevin oscillator (GLO)**
- “Half-time”: short summary (and maybe time for a few questions)

3 Results and discussion

- Laser-driven diffusion
- Laser-driven desorption
- Physisorbed precursor states?

4 Summary and Outlook

Generalized Langevin Oscillator


$$m_s \frac{d^2 \underline{r}_s}{dt^2} = -\underbrace{\nabla_s V(\underline{r}_1 - \underline{r}_s, \underline{r}_2 - \underline{r}_s)}_{\text{Force due to PES}} - m_s \underline{\Omega}^2 \underline{r}_s + m_s \underline{\Lambda}_{gs} \underline{r}_s$$

Surface oscillator

$$m_s \frac{d^2 \underline{r}_g}{dt^2} = -\underbrace{m_s \underline{\Omega}^2 \underline{r}_g}_{\text{Harmonic oscillator}} + \underbrace{m_s \underline{\Lambda}_{gs} \underline{r}_s}_{\text{Coupling to surface force}} - \eta_{ph} \frac{d \underline{r}_g}{dt} + R_{ph}(T_{ph})$$

ghost oscillator

both oscillate in the 3 dimensions x, y, z



Outline

1 Introduction

- Motivation - in general and system specific
- First impressions of fs-laser-driven dynamics

2 Models and methods

- Foundations: 6D potential and two-temperature model
- Electronic friction: non-adiabatic coupling approximated
- The generalized Langevin oscillator (GLO)
- **“Half-time”:** short summary (and maybe time for a few questions)

3 Results and discussion

- Laser-driven diffusion
- Laser-driven desorption
- Physisorbed precursor states?

4 Summary and Outlook

Outline

1 Introduction

- Motivation - in general and system specific
- First impressions of fs-laser-driven dynamics

2 Models and methods

- Foundations: 6D potential and two-temperature model
- Electronic friction: non-adiabatic coupling approximated
- The generalized Langevin oscillator (GLO)
- “Half-time”: short summary (and maybe time for a few questions)


3 Results and discussion

- **Laser-driven diffusion**
- Laser-driven desorption
- Physisorbed precursor states?

4 Summary and Outlook

Example trajectory

(in this case without GLO-model)



Outline

1 Introduction

- Motivation - in general and system specific
- First impressions of fs-laser-driven dynamics

2 Models and methods


- Foundations: 6D potential and two-temperature model
- Electronic friction: non-adiabatic coupling approximated
- The generalized Langevin oscillator (GLO)
- “Half-time”: short summary (and maybe time for a few questions)

3 Results and discussion


- Laser-driven diffusion
- **Laser-driven desorption**
- Physisorbed precursor states?

4 Summary and Outlook


Heating of DOFs Z and θ



Fluence-dependence of desorption yield P_{des}



Two-pulse correlation of desorption



Outline

1 Introduction

- Motivation - in general and system specific
- First impressions of fs-laser-driven dynamics

2 Models and methods


- Foundations: 6D potential and two-temperature model
- Electronic friction: non-adiabatic coupling approximated
- The generalized Langevin oscillator (GLO)
- “Half-time”: short summary (and maybe time for a few questions)

3 Results and discussion

- Laser-driven diffusion
- Laser-driven desorption
- **Physisorbed precursor states?**


4 Summary and Outlook

Anisotropic diffusion patterns + dynamical trapping



Surprising patterns in XY-distribution

- preference of **hcp-site** after 5-7 ps, despite it being a local maximum!
⇒ **dynamical trapping** (cf. 30 ps)
- effect dependent on fluence
⇒ consistent with experiment
(weaker “precursor”-signal for lower fluence)



Outline

1 Introduction

- Motivation - in general and system specific
- First impressions of fs-laser-driven dynamics

2 Models and methods

- Foundations: 6D potential and two-temperature model
- Electronic friction: non-adiabatic coupling approximated
- The generalized Langevin oscillator (GLO)
- “Half-time”: short summary (and maybe time for a few questions)

3 Results and discussion

- Laser-driven diffusion
- Laser-driven desorption
- Physisorbed precursor states?

4 Summary and Outlook

Introduction
oooooooo

Models and methods
oooooooooooo

Results and discussion
oooooooo

Summary and Outlook

Summary

Introduction
oooooooo

Models and methods
oooooooooooo

Results and discussion
oooooooo

Summary and Outlook

Outlook