

**Springer Series in  
Operations Research  
and Financial Engineering**

**Sidney I. Resnick**

**Extreme Values,  
Regular Variation,  
and Point Processes**

 Springer

# Contents

	<i>Preface to the Soft Cover Edition</i> .....	v
	<i>Preface to the Hard Cover Edition</i> .....	vii
0	<i>Preliminaries</i> .....	1
0.1	Uniform Convergence .....	1
0.2	Inverses of Monotone Functions .....	3
0.3	Convergence to Types Theorem and Limit Distributions of Maxima .....	7
0.4	Regularly Varying Functions of a Real Variable .....	12
0.4.1	Basics .....	13
0.4.2	Deeper Results; Karamata's Theorem .....	16
0.4.3	Extensions of Regular Variation: $\Pi$ -Variation, $\Gamma$ -Variation ...	26
1	<i>Domains of Attraction and Norming Constants</i> .....	38
1.1	Domain of Attraction of $\Lambda(x) = \exp\{-e^{-x}\}$ .....	38
1.2	Domain of Attraction of $\Phi_\alpha(x) = \exp\{-x^{-\alpha}\}$ , $x > 0$ .....	54
1.3	Domain of Attraction of $\Psi_\alpha(x) = \exp\{-(-x)^\alpha\}$ , $x < 0$ .....	59
1.4	Von Mises Conditions .....	62
1.5	Equivalence Classes and Computation of Normalizing Constants .....	67
2	<i>Quality of Convergence</i> .....	76
2.1	Moment Convergence .....	76
2.2	Density Convergence .....	85
2.3	Large Deviations .....	94
2.4	Uniform Rates of Convergence to Extreme Value Laws .....	107
2.4.1	Uniform Rates of Convergence to $\Phi_\alpha(x)$ .....	107
2.4.2	Uniform Rates of Convergence to $\Lambda(x)$ .....	114
3	<i>Point Processes</i> .....	123
3.1	Fundamentals .....	123
3.2	Laplace Functionals .....	128
3.3	Poisson Processes .....	130

3.3.1	Definition and Construction . . . . .	130
3.3.2	Transformations of Poisson Processes . . . . .	134
3.4	Vague Convergence . . . . .	139
3.5	Weak Convergence of Point Processes and Random Measures . . . . .	150
4	<i>Records and Extremal Processes</i> . . . . .	162
4.1	Structure of Records . . . . .	164
4.2	Limit Laws for Records . . . . .	174
4.3	Extremal Processes . . . . .	179
4.4	Weak Convergence to Extremal Processes . . . . .	196
4.4.1	Skorohod Spaces . . . . .	196
4.4.2	Weak Convergence of Maximal Processes to Extremal Processes via Weak Convergence of Induced Point Processes . . . . .	209
4.5	Extreme Value Theory for Moving Averages . . . . .	224
4.6	Independence of $k$ -Record Processes . . . . .	242
5	<i>Multivariate Extremes</i> . . . . .	250
5.1	Max-Infinite Divisibility . . . . .	251
5.2	An Example: The Bivariate Normal . . . . .	255
5.3	Characterizing Max-id Distributions . . . . .	257
5.4	Limit Distributions for Multivariate Extremes . . . . .	263
5.4.1	Characterizing Max-Stable Distributions . . . . .	266
5.4.2	Domains of Attraction; Multivariate Regular Variation . . . . .	276
5.5	Independence and Dependence . . . . .	290
5.6	Association . . . . .	298
	<i>References</i> . . . . .	307
	<i>Index</i> . . . . .	315

criteria are only a control on the right tail. For instance, if  $F \in D(\Phi_\alpha)$  then  $1 - F(x) \sim x^{-\alpha} L(x)$  as  $x \rightarrow \infty$ . This implies

$$\int_0^\infty x^k F(dx) < \infty \quad \text{if } k < \alpha$$

(Exercise 1.2.2), but no control is provided over the left tail and it is possible for  $\int_{-\infty}^0 |x|^k F(dx) = \infty$  for any  $k > 0$ . Similarly  $F \in D(\Lambda)$  implies when  $x_0 = \infty$  that

$$\int_0^\infty x^k F(dx) < \infty \quad \text{for all } k > 0$$

(Exercise 1.1.1) but implies nothing about behavior of the left tail.

Thus in investigating (2.1) it is necessary to impose some condition on the left tail of  $F$ .

**Proposition 2.1.** *For an extreme value distribution  $G$ , suppose  $F \in D(G)$ .*

(i) *If  $G = \Phi_\alpha$ , set  $a_n = (1/(1 - F))^{+}(n)$ ,  $b_n = 0$ . If for some integer  $0 < k < \alpha$*

$$\int_{-\infty}^0 |z|^k F(dx) < \infty \quad (2.3)$$

*then*

$$\lim_{n \rightarrow \infty} E(M_n/a_n)^k = \int_{-\infty}^\infty x^k \Phi_\alpha(dx) = \Gamma(1 - \alpha^{-1}k).$$

(ii) *If  $G = \Psi_\alpha$  and  $F$  has right end  $x_0$  set*

$$a_n = x_0 - (1/(1 - F))^{+}(n), \quad b_n = x_0.$$

*If for some integer  $k > 0$*

$$\int_{-\infty}^{x_0} |x|^k F(dx) < \infty \quad (2.4)$$

*then*

$$\lim_{n \rightarrow \infty} E((M_n - x_0)/a_n)^k = \int_{-\infty}^0 x^k \Psi_\alpha(dx) = (-1)^k \Gamma(1 + \alpha^{-1}k).$$

(iii) *If  $G = \Lambda$  and  $F$  has right end  $x_0$  with representation (1.5) set  $b_n = (1/(1 - F))^{+}(n)$ ,  $a_n = f(b_n)$ . If for some integer  $k > 0$*

$$\int_{-\infty}^0 |x|^k F(dx) < \infty \quad (2.5)$$

*then*

$$\lim_{n \rightarrow \infty} E((M_n - b_n)/a_n)^k = \int_{-\infty}^\infty x^k \Lambda(dx) = (-1)^k \Gamma^{(k)}(1)$$

where  $\Gamma^{(k)}(1)$  is the  $k$ th derivative of the gamma function at  $x = 1$ .

Question 2

*Remarks.* (i) Conditions (2.3), (2.4), and (2.5) can be weakened slightly. See Exercise 2.1.1.

(ii) For any norming constants  $a_n, b_n$  satisfying (2.1) (not just the ones specified in the statement of the proposition), we also have (2.2) satisfied provided the appropriate condition (2.3), (2.4), or (2.5) holds. See Exercise 2.1.2.

We only prove (i) and (iii) in Proposition 2.1. Part (iii) requires that our tool box be equipped with the following inequalities.

**Lemma 2.2.** *Suppose  $F \in D(\Lambda)$  with representation (1.5) and that  $a_n$  and  $b_n$  are as specified in part (iii) of Proposition 2.1.*

(a) *Given  $\varepsilon > 0$ , we have for  $s > 0$  and all sufficiently large  $n$*

$$f(b_n)/f(a_ns + b_n) \geq (1 + \varepsilon s)^{-1} \quad (2.6)$$

*and consequently if  $y > 0$  and  $n$  is large*

$$1 - F^n(a_n y + b_n) \leq (1 + \varepsilon)^3 (1 + \varepsilon y)^{-\varepsilon^{-1}}. \quad (2.7)$$

(b) *Recall the meaning of  $z_0$  in the representation (1.5). Given  $\varepsilon$ , pick  $z_1 \in (z_0, x_0)$  such that  $|f'(x)| \leq \varepsilon$  if  $x > z_1$ . Then for large  $n$  and  $u \in (a_n^{-1}(z_1 - b_n), 0)$  we have*

$$f(b_n)/f(a_n u + b_n) \geq (1 + \varepsilon|u|)^{-1} \quad (2.8)$$

*and consequently for large  $n$  and  $s \in (a_n^{-1}(z_1 - b_n), 0)$*

$$F^n(a_n s + b_n) \leq e^{-(1-\varepsilon)^2(1+\varepsilon|s|)^{\varepsilon^{-1}}}. \quad (2.9)$$

**PROOF OF LEMMA 2.2.** (a) For  $n$  such that  $|f'(t)| \leq \varepsilon$  if  $t \geq b_n$  we have for  $s > 0$

$$(f(a_n s + b_n)/f(b_n)) - 1 = \int_{b_n}^{a_n s + b_n} (f'(u)/f(b_n)) du$$

and recalling  $a_n = f(b_n)$  this is

$$\int_0^s f'(a_n u + b_n) du \leq \varepsilon s.$$

Consequently  $f(b_n)/f(a_n s + b_n) \geq (1 + \varepsilon s)^{-1}$  as asserted.

To check (2.7) note that

$$1 - F(b_n) \sim n^{-1}$$

so that for large  $n$  and  $y > 0$

$$n(1 - F(a_n y + b_n)) \leq (1 + \varepsilon)(1 - F(a_n y + b_n))/(1 - F(b_n))$$

and from (1.5) this is

$$(1 + \varepsilon)c(a_n y + b_n)c^{-1}(b_n)e^{-\int_{b_n}^{a_n y + b_n} (1/f(s)) ds}.$$

Since  $c(x) \rightarrow c > 0$  as  $x \uparrow x_0$  we have the preceding

$$\leq (1 + \varepsilon)^2 e^{-\int_0^s (f(b_n)/f(a_n s + b_n)) ds}$$