

A model for neural activity in the absence of external stimuli

Aline Duarte and Guilherme Ost *

Universidade de São Paulo and GSSI - L'Aquila †

November 7, 2014

Abstract

We study a stochastic process describing the continuous time evolution of the membrane potentials of finite system of neurons in the absence of external stimuli. The values of the membrane potentials evolve under the effect of *chemical synapses*, *electrical synapses* and a *leak current*. The evolution of the process can be informally described as follows. Each neuron spikes randomly following a point process with rate depending on its membrane potential. When a neuron spikes, its membrane potential is immediately reset to a resting value. Simultaneously, the membrane potential of the neurons which are influenced by it receive an additional positive value. Furthermore, between consecutive spikes, the system follows a deterministic motion due both to electrical synapses and the leak current. Electrical synapses push the system towards its average potential, while the leak current attracts the membrane potential of each neuron to the resting value.

We show that in the absence leakage the process converges exponentially fast to an unique invariant measure, whenever the initial configuration is non null. More interesting, when leakage is present, we proved the system stops spiking after a finite amount of time almost surely. This implies that the unique invariant measure is supported only by the null configuration.

Key words : piecewise deterministic Markov process, limiting distribution, neuronal systems, chemical synapses, electrical synapses, leak current

AMS Classification: 60K35, 60F99, 60J25

1 Introduction

We study the behavior of a finite number of interacting neurons in the absence of external stimuli our goal being to determine the long-run behavior of the process. Our system is composed of N neurons whose state at time $t \geq 0$ is specified by $U(t) = (U_1(t), \dots, U_N(t))$, with $U_i(t) \in \mathbb{R}_+^N$. For each neuron $i = 1, \dots, N$ and each time $t \geq 0$, $U_i(t)$ represents the

*e-mail addresses: aline.duarte@gmail and guilhermeost@gmail.com

†The authors are PhD students from the Universidade de São Paulo and this work was developed during their partial completion doctoral program at GSSI both being fully supported by CNPq.

membrane potential of neuron i at time t . We consider two kinds of interactions among neurons and also a constant interplay between neurons and the environment.

More precisely the neurons interact via *electrical* and *chemical synapses*. Electrical synapses are due to so-called *gap-junction channels* between neurons which induce a constant sharing of potential, pushing the system towards its average value. By contrast, chemical synapses are point events which can be described as follows. Each neuron spikes randomly at rate $\varphi(U) \geq 0$ which depends on its membrane potential U , we suppose φ a non decreasing function, positive at $U > 0$ and both integrable and vanishing at 0 (in agreement with the assumption of non external stimuli). When neuron i spikes, its membrane potential is immediately reset to a resting potential 0. Simultaneously, the neurons which are influenced by neuron i receive an additional positive value to their membrane potential. This value may vary for each pair of neurons. Moreover, in the whole time, the neurons loose potential to the environment, due to *leakage channels* which pushes down the membrane potential of each neuron toward zero. This outgoing constant flow of potential is defined as the *leak current*. For technical details we refer the reader to Gersnter and Kistler (2002).

Our system is inspired by the one introduced in Galves and Löcherbach (2013). This model is an example of piecewise-deterministic Markov processes (PDPs) introduced by Davis (1984). Such processes combine random jump events, the chemical synapses, with deterministic continuous evolutions, in our case due both to electrical synapses and the leak current. The PDPs have been used also to model neuronal systems by other authors, see for instance the papers by Riedler, Thieullen and Wainrib (2012), De Masi et al. (2014), Fournier and Löcherbach (2014) and Robert and Touboul (2014).

Chemical synapses and leakage make the system non-conservative. Moreover, there is an evident competition between the incoming energy induced by the spikes and the outgoing energy induced by the leak current. Therefore it is natural to ask about the limiting behavior of the system as time $t \rightarrow \infty$. The main results of the paper, presented in Theorem 2.3 and Theorem 2.4, provide a complete description of the asymptotic distribution of the process. The theorem 2.3 states that under the presence of the leakage then almost surely there are only a finite number of spikes and the system converges to an “inactive global state” interpreted as “brain sleep”.

When the leakage is absent we prove a Harris-type condition which shows exponential convergence to an unique non trivial invariant measure, whenever the initial configuration is non null. This is the content of the Theorem 2.4.

Our paper is organized as follows. In section 2, we introduce the process, prove its existence and we state the main results, Theorem 2.3 and Theorem 2.4. In Section 3, we proof the theorem 2.3, while in section 4, we prove Theorem 2.4. In section 5, we briefly compare similar results recently obtained by Roberto and Touboul (2014).

2 Model definition and main results

Let $\mathcal{N} = \{1, \dots, N\}$ be a finite set of neurons, for some fixed integer $N \geq 1$ and consider the family of non-negative synaptic weights $(W_{i \rightarrow j})_{i,j \in \mathcal{N}} \in \mathbb{R}_+^{\binom{N}{2}}$ such that $W_{i \rightarrow i} = 0$ for all $i \in \mathcal{N}$. The value $W_{i \rightarrow j}$ corresponds to the value added to the membrane potential of neuron j when the neuron i spikes.

We consider a continuous time Markov process

$$U(t) = (U_1(t), \dots, U_N(t)), \quad t \geq 0,$$

taking values in \mathbb{R}_+^N , whose infinitesimal generator is given for any smooth test function $f : \mathbb{R}_+^N \rightarrow \mathbb{R}$, by

$$\mathcal{L}f(u) = \sum_{i \in \mathcal{N}} \varphi(u_i) [f(\Delta_i(u)) - f(u)] - \lambda \sum_{i \in \mathcal{N}} \left(\frac{\partial f}{\partial u_i}(u) [u_i - \bar{u}] \right) - \alpha \sum_{i \in \mathcal{N}} \left(\frac{\partial f}{\partial u_i}(u) [u_i] \right), \quad (1)$$

where for all $i \in \mathcal{N}$, $\Delta_i : \mathbb{R}_+^N \rightarrow \mathbb{R}_+^N$ is defined by

$$(\Delta_i(u))_j = \begin{cases} u_j + W_{i \rightarrow j}, & \text{if } j \neq i \\ 0, & \text{if } j = i \end{cases},$$

$\lambda, \alpha \geq 0$ are positive parameters modelling, respectively, the strength of electrical synapses and the leakage effect, $\bar{u} = \frac{1}{N} \sum_{i=1}^N u_i$ and

Assumption 1. $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ is a non-decreasing function such that $\varphi(0) = 0$ and there exists a constant $r > 0$ satisfying

$$\int_0^{2r} \frac{\varphi(u)}{u} du < +\infty.$$

We are assuming that there is no external stimuli. This assumption appears in the condition $\varphi(0) = 0$. In addition, from the neurobiological point of view, it is reasonable to assume that φ is a non-decreasing function since an addition in the membrane potential increases the probability of a spiking occurs.

The first term in (1) depicts how the chemical synapses are incorporated in our model. Neurons whose potential is u spike at rate $\varphi(u)$. Intuitively this means that any initial configuration $u \in \mathbb{R}_+^N$ of the membrane potentials

$$\mathbb{P}(U(t) = \Delta_i(u) | U(0) = u) = \varphi(u_i)t + o(t), \quad \text{as } t \rightarrow 0$$

Thus, the function φ is called firing or spiking rate of the system.

The second and third terms in (1) represent the electrical synapses and the leak current respectively. They describe the deterministic time evolution of the system between two consecutive spikes. More specifically, in an interval of time $[a, b]$, without occurrence of spikes in the whole system, the membrane potential of neuron $i \in \mathcal{N}$ obeys the following ordinary differential equation

$$\frac{d}{dt} U_i(t) = -\alpha U_i(t) - \lambda(U_i(t) - \bar{U}(t)). \quad (2)$$

Notice that the first term of the right-hand side of (2) pushes simultaneously all neurons to the resting state, while the second term tends to attract the neurons to the average potential.

Our first theorem proves the existence of the process.

Theorem 2.1. *Let $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ be any function satisfying the Assumption 1. For any $N \geq 1$ and any $u \in \mathbb{R}_+^N$ there exists a unique strong Markov process $U(t)$ taking values in \mathbb{R}_+^N starting from u whose generator is given by (1).*

Proof. Let $N_i(t)$, $t \geq 0$, be the simple point process on \mathbb{R}_+ which counts the jump events of neuron $i \in \mathcal{N}$ up to time t . Define $E_i = \sum_{j \neq i} W_{i \rightarrow j}$ and $E = \max_{i \in \mathcal{N}} E_i$ and, following De Masi et al. (2014), consider the following random variable, for all $t > 0$,

$$k(t) = \sum_{i \in \mathcal{N}} \int_0^t \mathbf{1}\{U_i(s^-) \leq 2E\} dN_i(s).$$

The random variable $k(t)$ counts the number of spikes of neurons whose the potential is at most $2E$.

Suppose U_i fires at time t , in this case

$$\bar{U}(t) = \frac{1}{N} \sum_{j \neq i} (U_j(t^-) + W_{i \rightarrow j}) = \bar{U}(t^-) + \frac{1}{N} (E_i - U_i(t^-)).$$

Now, using the expression of $\bar{U}(t)$ above and adapting the proof of Theorem 1 of De Masi et al (2014), we have the following inequalities for all $t > 0$,

$$\bar{U}(t) \leq \bar{U}(0) + \frac{E}{N} k(t), \quad EN(t) \leq N\bar{U}(0) + 2Ek(t) \quad \text{and} \quad U^{max}(t) \leq (N+1)U^{max}(0) + 2Ek(t),$$

where $U^{max}(t) = \max_{i \in \mathcal{N}} U_i(t)$.

Since we can bound $Ek(t)$ by a Poisson process of intensity $N\varphi(2E)$, the second inequality above shows that number of jumps of the process is finite almost surely on any finite time interval. To conclude the proof just note that the construction of the process can be achieved by gluing together trajectories given by the deterministic flow between successive jump times. This procedure is feasible since the number of jumps of the process is finite on any finite interval. □

Now, we shall present an elementary argument which shows that for all leakage rate α large enough and if the firing rate φ is globally Lipschitz with $\varphi(0) = 0$, the system goes extinct. This result was the starting point of this paper. The idea of this proof was taken from discussions with Galves and Löcherbach. The result is the following.

Theorem 2.2. *For any $N \geq 1$, $\alpha \geq 0$, $\lambda \geq 0$ and c -Lipschitz function $\varphi : \mathbb{R}_+ \rightarrow \mathbb{R}_+$ such that $c > 0$ and $\varphi(0) = 0$, the following inequality holds, for all $t \geq 0$ and $U(0) = u \in \mathbb{R}_+^N$,*

$$\mathbb{E}[\bar{U}(t)] \leq \bar{u} e^{t(\alpha^* c - \alpha)},$$

where $\alpha^* = \max_{k \in \mathcal{N}} \sum_{j \in \mathcal{N}} W_{j \rightarrow k}$. In particular, if $\alpha > \alpha^* c$, then the process goes extinct.

Proof. For each $i \in \mathcal{N}$, plugging $f = \pi_i$ in (1), where π_i is the projection onto the i -th coordinate, we have

$$\frac{d}{dt} \mathbb{E}[U_i(t)] = \sum_{j \in \mathcal{N}} W_{j \rightarrow i} \mathbb{E}[\varphi(U_j(t))] - \mathbb{E}[U_i(t) \varphi(U_i(t))] - \alpha \mathbb{E}[U_i(t)] + \lambda \mathbb{E}[U_i(t) - \bar{U}(t)].$$

Summing over all $i \in \mathcal{N}$ and then using that φ is a non-negative c -Lipschitz function such that $\varphi(0) = 0$, it follows that

$$\frac{d}{dt}\mathbb{E}[\bar{U}(t)] \leq (\alpha^* c - \alpha)\mathbb{E}[\varphi(\bar{U}(t))].$$

Therefore applying the Grownwall's lemma to the inequality above we finish the proof. \square

Even assuming that the firing rate φ satisfies only the Assumption 1, we claim that for any fixed numbers of neurons, the presence of the leak current is a necessary and sufficient condition for the extinction of the process. In fact, we shall prove a stronger result. It states that, if there is the leakage, there will be only a finite numbers of spikes eventually almost surely. On the other hand, it is shown that, excluding the trivial initial configuration, the system is ergodic when there is no leakage. In particular, this results generalize the Theorem 2.2 above.

In order to state our main result, we need to introduce the some extra notation.

For each neuron $i \in \mathcal{N}$, let $T_1^i = \inf\{s > 0 : U_i(s) = 0\}$ be the first spiking time of neuron i and for each $k \geq 2$, let $T_k^i = \inf\{s > T_{k-1}^i : U_i(s) = 0\}$ be the k -th spiking time of neuron i . Then, the first and the k -th spiking time of the system are defined respectively by

$$T_1 = \inf_{i \in \mathcal{N}} T_1^i \quad \text{and} \quad T_k = \inf_{i \in \mathcal{N}, m \geq 1} \{T_k^i > T_{m-1}\}, \quad k \geq 2. \quad (3)$$

Our main theorem is given below.

Theorem 2.3. *Let $(U(t))_{t \geq 0}$ be the Markov process whose the infinitesimal generator is given by (1) and T_k be as defined in (3). Assume that φ satisfies the Assumption 1 with $r > \max_{i \in \mathcal{N}} \sum_{j \in \mathcal{N}} W_{j \rightarrow i}$. Then for any $\alpha > 0$ and $\lambda \geq 0$,*

$$\mathbb{P} \left(\sum_{k \geq 1} \mathbf{1}\{T_k < \infty\} < \infty \right) = 1.$$

Corollary 1. *Under the same hypothesis of Theorem 2.3, for all $i \in \mathcal{N}$, it holds*

$$\lim_{t \rightarrow +\infty} U_i(t) = 0 \quad a.s.$$

In particular, the delta of Dirac at point 0^N , δ_{0^N} , is the unique invariant measure for the process in the presence of the leak current.

It remains to analyse what happens in the long-run behavior of the system in the absence of the leakage. This is the content of the next result.

Theorem 2.4. *Let $(U(t))_{t \geq 0}$ be the Markov process whose the infinitesimal generator is given by (1) with $\alpha = 0$. Then the process admits only two different invariant probability measures either δ_{0^N} or $\mu \neq \delta_{0^N}$. If $U(0) = (0, \dots, 0)$, the invariant measure is δ_{0^N} , while whatever is the initial configuration $U(0) = u \in \mathbb{R}_+^N \setminus \{0^N\}$, the invariant measure is μ .*

3 Proof of Theorem 2.3

First of all, observe that, from the equation (2), for any time $t \in [T_n, T_{n+1})$,

$$U_i(t) = (U_i(T_n) - \bar{U}(T_n))e^{-(\alpha+\lambda)(t-T_n)} + \bar{U}(T_n)e^{-\alpha(t-T_n)}. \quad (4)$$

We shall explore this equation many times.

We start giving a lower bound to the probability of there is no spikes when the system starts with a initial condition small enough.

Proposition 3.1. *Let $(U(t))_{t \in \mathbb{R}}$ be the Markov process whose the generator is given by (1), $U^{max}(t) = \max\{U_i(t), i \in \mathcal{N}\}$ be the maximum potential of the system at time t and T_1 as defined in (3). Suppose that φ satisfies the Assumption 1. If $\alpha > 0$, then*

$$\mathbb{P}(T_1 = \infty \mid U^{max}(0) < r) \geq e^{-N\alpha \int_0^{2r} \frac{\varphi(u)}{u} du} > 0.$$

Proof. By the equation (4), we have, for all $0 \leq t < T_1$, the following inequalities

$$U^{max}(t) \leq \bar{U}(0)e^{-\alpha t} + U^{max}(0)e^{-\alpha t} \leq 2re^{-\alpha t}.$$

Using the inequality above and the non-decreasing assumption on φ , we have

$$\begin{aligned} \mathbb{P}(T_1 > t \mid U^{max}(0) < r) &\geq \exp \left\{ - \int_0^t N\varphi(U^{max}(s))ds \right\} \\ &\geq \exp \left\{ -N\alpha \int_{2re^{-\alpha t}}^{2r} \frac{\varphi(u)}{u} du \right\}. \end{aligned} \quad (5)$$

Therefore, when t goes to infinite, it follows the result. \square

Proposition 3.2. *Under the same hypothesis of Proposition 3.1. For $\theta < (\alpha + \lambda)^{-1}$, it holds that*

$$\mathbb{P}(T_1 > \theta \mid U^{max}(0) > r) \leq e^{-\theta\varphi(\beta)} < 1,$$

where $\beta = \beta(\theta) = r(1 - (\alpha + \lambda)\theta) > 0$.

Proof. Indeed,

$$\mathbb{P}(T_1 > \theta \mid U^{max}(0) > r) \leq \exp \left\{ - \int_0^\theta \varphi(U^{max}(s))ds \right\}. \quad (6)$$

Moreover, by (4) for all $0 < s \leq \theta$ is true that

$$U^{max}(s) \geq U^{max}(0)e^{-(\alpha+\lambda)s} \geq re^{-(\alpha+\lambda)s} \geq r(1 - (\alpha + \lambda)\theta). \quad (7)$$

From (6) and (7) we deduce the desired inequality

$$\mathbb{P}(T_1 > \theta \mid U^{max}(0) \geq r) \leq e^{-\theta\varphi(\beta)} < 1.$$

\square

The next result claims that even when all membrane potentials are large, there is a positive probability of all potentials become small after a fix time $T > 0$.

Proposition 3.3. *Consider $(U(t))_{t \in \mathbb{R}}$ the Markov process whose the generator is given by (1), $U^{max}(t)$ and T_1 as in proposition 3.1. Suppose that φ satisfies the Assumption 1 and $\alpha > 0$. If $r > \max_{i \in \mathcal{N}} \sum_{j \in \mathcal{N}} W_{j \rightarrow i}$, then there exists $T > 0$ such that*

$$\mathbb{P}(U^{max}(T) < r \mid U^{max}(0) > r) > 0.$$

Proof. Define $\tau_1 = \inf\{t > 0 : U^{max}(t) = 0\}$ and take $\theta < (\alpha + \lambda)^{-1}$ (this is possible since $\alpha > 0$). By the proposition 3.2, it holds that

$$\mathbb{P}(T_1 < \theta, T_1 = \tau_1 \mid U^{max}(0) > r) \geq \left(1 - e^{-\theta\varphi(\beta)}\right) \frac{1}{N}.$$

Define $T_0 = 0$, $T = N\theta$ and consider the following sequence of events

$$A_1 = \{T_1 < \theta, T_1 = \tau_1, U^{max}(T_1) < r, T_2 > T\},$$

and for $l = 2 \dots, N$,

$$A_l = \left\{ \cap_{k=1}^l \{T_k < T_{k-1} + \delta, T_k = \tau_k\}, \cap_{k=1}^{l-1} \{U^{max}(T_k) > r\}, U^{max}(T_l) < r, T_{l+1} > T \right\}.$$

The event A_k corresponds to the following situation. At the firsts $k - 1$ consecutive spikes of the maximum, there always exists at least one neuron whose potential is larger than r . But at k -th spike of the maximum all membrane potentials get less than r and no more spikes happen up to time T .

From the definition of the events A_k we have

$$\{U^{max}(T) < r\} \supseteq \bigcup_{k=1}^N A_k.$$

Wherefore, it suffices to compute the probability of each event A_k in the right side of the inclusion, conditioned to the event $\{U^{max}(0) > r\}$. Now, defining $\beta_0 = (1 - e^{-\delta\varphi(\beta)}) \frac{1}{N}$, and $\beta_1 = \exp\left\{\frac{-2Ncr}{\alpha}(1 - e^{-\alpha N\delta})\right\}$, by the proposition 3.2 and the inequality (5), it follows that

$$\mathbb{P}(A_1 \mid U^{max}(0) > r) \geq \beta_0 \beta_1 \mathbb{P}(U^{max}(T_1) < r \mid U^{max}(0) > r, T_1 < \delta, T_1 = \tau_1).$$

Similarly, we have

$$\begin{aligned} \mathbb{P}(A_2 \mid U^{max}(0) > r) &\geq \beta_0^2 \beta_1 \mathbb{P}(U^{max}(T_1) > r \mid U^{max}(0) > r, T_1 < \delta, T_1 = \tau_1) \\ &\quad \times \mathbb{P}(U^{max}(T_2) < r \mid U^{max}(T_1) > r, T_2 < T_1 + \delta, T_2 = \tau_2). \end{aligned}$$

Thus, summing these two inequalities, we get the following lower bound for $A_1 \cup A_2$,

$$\mathbb{P}(A_1 \cup A_2 \mid U^{max}(0) > r) \geq \beta_0^2 \beta_1 \mathbb{P}(U^{max}(T_2) < r \mid U^{max}(T_1) > r, T_2 < T_1 + \delta, T_2 = \tau_2).$$

Proceeding in this way for the other terms, we obtain that

$$\begin{aligned} \mathbb{P}(U^{max}(T) < r \mid U^{max}(0) > r) \\ \geq \beta_0^N \beta_1 \mathbb{P}(U^{max}(T_N) < r \mid U^{max}(T_{N-1}) > r, \cap_{k=1}^N \{T_k < T_{k-1} + \delta, T_k = \tau_k\}). \end{aligned}$$

Since $r > \max_{i \in \mathcal{N}} \sum_{j \in \mathcal{N}} W_{j \rightarrow i}$, we easily see that

$$\mathbb{P} \left(U^{max}(T_N) < r \mid U^{max}(T_{N-1}) > r, \cap_{k=1}^N \{T_k < T_{k-1} + \delta, T_k = \tau_k\} \right) = 1.$$

□

As a consequence of proposition 3.3 we have

Corollary 2. *Let $T > 0$ be the positive constant given by the proposition 3.3. Define the stopping times $R_1 = \inf\{n \geq 1; U^{max}(nT) \leq r\}$ and $R_k = \inf\{n \geq R_{k-1}; U^{max}(nT) \leq r\}$ for all $k \geq 2$. Then, under the same hypothesis of the proposition 3.3,*

$$\sum_{k=1}^{\infty} \mathbf{1}\{R_k < \infty\} = \infty \quad \mathbb{P} - a.s.$$

Proof of Theorem 2.3. For this proof we must define the following stopping times: $K_1 = \inf\{n \geq 1 : T_n > R_1 T\}$, $J_1 = \inf\{n \geq 1 : R_n T > T_{S_1}\}$, and for $k \geq 2$, $K_k = \inf\{n > J_{k-1} : T_n > R_{J_{k-1}} T\}$ and $J_k = \inf\{n \geq 1 : R_n T > T_{S_{k-1}}\}$. From corollary 2 all these stopping times are finite almost surely and, in particular, well defined. Now, from Theorem 2.1 and the definition of K_k

$$1 = \mathbb{P} \left(\bigcap_{k=1}^{\infty} \bigcup_{i=1}^{\infty} \{T_i > R_k T\} \mid \bigcap_{i=1}^{\infty} \{T_i < \infty\} \right) = \mathbb{P} \left(\bigcap_{k=1}^{\infty} \{K_k < \infty\} \mid \bigcap_{i=1}^{\infty} \{T_i < \infty\} \right).$$

Thus, $\cap_{k=1}^{\infty} \{T_k < \infty\}$ implies $\cap_{k=1}^{\infty} \{K_k < \infty\}$.

On the other hand, by proposition 3.1, one knows that

$$\mathbb{P} \left(\bigcap_{k=1}^n \{K_k < \infty\} \right) \leq \left(1 - e^{-N\alpha \int_0^{2r} \frac{\varphi(u)}{u} du} \right)^n,$$

which converges to 0 when n diverges.

□

4 Proof of Theorem 2.4

To simplify the proof of the theorem 4 we shall split it into several steps. The main part of the argument is to find a *positive recurrent regeneration set* B in sense that

- (i) B is positive recurrent and,
- (ii) there exist some $t_* > 0$, $\epsilon > 0$ and a probability measure ν on \mathbb{R}_+^N such that

$$P_{t_*}(u, A) \geq \epsilon \nu(A), \quad u \in B,$$

for all measurable set $A \in \mathcal{B}(\mathbb{R}_+^N)$.

Usually, a Markov processes with a regeneration set are called *Harris recurrent Markov processes*. For such Processes an invariant measure always exists.

In what follows, for any positive real number $a > 0$ we use the notation $R_a(x)$, meaning that there exists a constant $l > 0$ such that $|R_a(x)| \leq la$ for all x . When a function satisfies such condition it is called a function of order a . Note that we are not specifying the domain in which the function R is defined on.

For each $\varepsilon > 0$ define the following event

$$A_\varepsilon = \{(i-1)\varepsilon < T_i < i\varepsilon, \ i = 1, \dots, N\},$$

and let $(S_m)_{m \geq 1}$ be the spiking mark sequence of the system. The event $\{S_m = i\}$ means that the $U_i(T_m) = 0$. Finally, consider the event

$$S = \{S_i = i, i = 1, \dots, N\}.$$

The first lemma below says that, conditioning to the event A_ε , when ε is sufficiently small the process evolves, modulo an error of small order, as in the case without gap junction ($\lambda = 0$). Before stating this lemma we need to introduce a finite sequence of potential configurations.

Consider the sequence $(v(k))_{k=0, \dots, N}$ with $v(k) \in R_+^N$ given by

$$v(0) = \left(\sum_{j=2}^N W_{j \rightarrow 1}, \sum_{j=3}^N W_{j \rightarrow 1}, \dots, W_{N \rightarrow N-1}, 0 \right),$$

and for $1 \leq k \leq N$, $v_k(k) = 0$ and for $i \neq k$

$$v_i(k) = \mathbf{1}_{\{i > k\}} \left(v_i(0) + \sum_{j=1}^k W_{j \rightarrow i} \right) + \mathbf{1}_{\{i < k\}} \left(\sum_{j=i+1}^k W_{j \rightarrow i} \right).$$

Lemma 4.1. Fix $\delta > 0$. If $U(0) = u \in B(v(0), \delta)$, then conditioning to the event $A_\varepsilon \cap S$, for each $k = 1, \dots, N$ the following equalities hold:

- (i) $U_i(T_k) = v_i(k) + \sum_{r=i+1}^k \lambda(T_r - T_{r-1})d_i(r-1) + R_{\delta\varepsilon}(T_1^k, u) + R_{\varepsilon^2}(T_1^k, u)$, if $i < k$;
- (ii) $U_i(T_k) = v_i(k) + \sum_{r=1}^k \lambda(T_r - T_{r-1})d_i(r-1) + R_\delta(u) + R_{\delta\varepsilon}(T_1^k, u) + R_{\varepsilon^2}(T_1^k, u)$, if $i > k$;
- (iii) $\bar{U}(T_k) = \bar{v}(k) + R_\delta(u) + R_\varepsilon(T_1^k, u)$, if $k < N$ and $\bar{U}(T_N) = \bar{v}(0) + R_\varepsilon(T_1^N, u)$,

where $d_i(m) = \bar{v}(m) - v_i(m)$, $T_0 = 0$ and $T_1^k = (T_1, \dots, T_k)$. Furthermore, all the partial derivatives of the remainder functions $R_{\delta\varepsilon}(T_1^k, u)$, $R_{\varepsilon^2}(T_1^k, u)$ above are either of order δ or ε .

Proof. The proof is given by induction on k . On the event $A_\varepsilon \cap S$, we have that $U_1(T_1) = 0$ and for each $i \neq 1$ and $U(0) = u \in B(v(0), \delta)$,

$$\begin{aligned} U_i(T_1) &= W_{1 \rightarrow i} + \bar{U}(0) + (1 - \lambda T_1 + R_{\varepsilon^2}(T_1))(v_i(0) + R_\delta(u) - \bar{U}(0)) \\ &= v_i(1) + \lambda T_1 d_i(0) + R_\delta(u) + R_{\delta\varepsilon}(T_1, u) + R_{\varepsilon^2}(T_1, u), \end{aligned}$$

where in the first equality we have used the expansion series of the exponential function.

Thus, from the expression of $U_i(T_1)$ above, we conclude that

$$\begin{aligned}\bar{U}(T_1) &= \bar{v}(1) + \lambda T_1 \frac{1}{N} \sum_{i=2}^N d_i(0) + R_\delta(u) + R_{\varepsilon\delta}(T_1, u) + R_{\varepsilon^2}(T_1, u) \\ &= \bar{v}(1) + R_\delta(u) + R_{\varepsilon}(T_1, u).\end{aligned}$$

In addition, it is easy to check that all remainders functions above have partial derivatives with respect to T_1 and all of them are either R_δ or R_ϵ functions. Therefore (i), (ii) and (iii) it is verified for $k = 1$.

Now, suppose that (i), (ii) and (iii) hold for some fixed $1 < k < N$. As before, on the event $A_\epsilon \cap S$, we have $U_{k+1}(T_{k+1}) = 0$ and, for $i < k + 1$, by the inductive hypothesis,

$$\begin{aligned}U_i(T_{k+1}) &= W_{k+1 \rightarrow i} + \bar{U}(T_k) + (1 - \lambda(T_{k+1} - T_k) + R_{\varepsilon^2}(T_{k-1}, T_k))(U_i(T_k) - \bar{U}(T_k)) \\ &= v_i(k+1) + \sum_{r=i+1}^{k+1} \lambda(T_r - T_{r-1})d_i(r-1) + R_{\delta\epsilon}(T_1^{k+1}, u) + R_{\varepsilon^2}(T_1^{k+1}, u)\end{aligned}$$

Using the same argument for the case when $i > k + 1$, we get the first equation. Using again the inductive hypothesis and looking at the expression written in the first equality of the membrane potential, it is readily seen that the remainder functions possesses partial derivatives with to T_l , for $l = 1, \dots, k + 1$ and they are either of order δ or ϵ .

Finally, summing $U_i(T_{k+1})$ over all neurons $i = 1, \dots, N$

$$\bar{U}(T_{k+1}) = \bar{v}(k+1) + R_{\varepsilon}(T_1^{k+1}, u) + R_\delta(u).$$

□

Note that $v(T_N) = v(0)$, thus, from the previous lemma it follows the

Corollary 3. *Under the same assumptions of Lemma 4.1, if $T = N\varepsilon$, then for each $i = 1, \dots, N$, the following equality is verified*

$$U_i(T) = v_i(0) + \lambda(T - T_N)d_i(0) + \sum_{r=i+1}^N \lambda(T_r - T_{r-1})d_i(r-1) + R_{\delta\epsilon}(T_1^N, u) + R_{\epsilon^2}(T_1^N, u).$$

Remark 4.1. In order to simplify the notation, we shall denote the map $\gamma^0 : A_\epsilon \rightarrow \mathbb{R}_+^N$ by $\gamma^0(t_1^N) = (\gamma_1^0(t_1^N), \dots, \gamma_N^0(t_1^N))$ where for each $i = 1, \dots, N$,

$$\gamma_i^0(t_1^N) = v_i(0) + \lambda(T - t_N)d_i(0) + \sum_{r=i+1}^N \lambda(t_r - t_{r-1})d_i(r-1).$$

By corollary 3, conditioning to the event $A_\epsilon \cap S$, we have the following representation for all $U(0) = u \in B(v(0), \delta)$

$$U(T) = \gamma^0(T_1^N) + R_{\delta\epsilon}(T_1^N, u) + R_{\epsilon^2}(T_1^N, u),$$

where both $R_{\delta\epsilon}(T_1^N, u)$ and $R_{\epsilon^2}(T_1^N, u)$ are multivalued functions whose the L_1 -norms are remainders functions of order $\delta\epsilon$ and ϵ^2 respectively. Moreover, we have the following corollary

Corollary 4. For each $u \in B(v(0), \delta)$, the determinant of the Jacobian of map $A_\varepsilon \ni t_1^N \mapsto \gamma_u(t_1^N) = \gamma^0(t_1^N) + R_{\delta\varepsilon}(t_1^N, u) + R_{\varepsilon^2}(t_1^N, u)$, is given by

$$|J\gamma_u(t_1^N)| = \lambda^N \prod_{i=1}^N \bar{v}(i) + R_\varepsilon(t_1^N, u) + R_\delta(t_1^N, u),$$

which is different from zero for δ and ε small enough for all $(t_1^N) \in A_\varepsilon$ and $u \in B(v(0), \delta)$.

We shall use this representation to show that our process, at time T , satisfies a Harris-type condition (see proposition 4.1). Before proving this proposition, we need an extra lemma. Here again the non-decreasing assumption on φ is important.

Lemma 4.2. (i) Let $f_u(t_1, \dots, t_N) = f_{u, (S_1=1, \dots, S_N=N)}(t_1, \dots, t_N)$ denote the joint density of (T_1, \dots, T_N) conditioning to the event S , when the starting configuration is u . Then, for any $0 < \delta < \min_{i=1, \dots, N} \{v_i(0)\}$, there exist a constant $\beta > 0$ such that for all $u \in B(v(0), \delta)$,

$$f_u(t_1, \dots, t_N) \geq C_1, \text{ for } (t_1, \dots, t_N) \in A_\varepsilon.$$

(ii) For any given $\min_{i=1, \dots, N} \{v_i(0)\} > \delta > 0$ there exist $\varepsilon > 0$ and a constant $C_2 > 0$ such that for $u \in B(v(0), \delta)$, we have

$$P_T(u, B(v(0), \delta)) \geq C_2 \varepsilon^N > 0.$$

In particular, defining $T_{B(v(0), \delta)} = \inf\{t > 0 : U(t) \in B(v(0), \delta)\}$, we have

$$\sup_{u \in B(v(0), \delta)} \mathbb{E}_u[T_{B(u^*, \delta)}] \leq \frac{T}{C_2 \varepsilon^N}.$$

Proof. (i) Since $P_u(T_1 > t) = \exp\left[-\int_0^t \sum_{j=1}^N \varphi(U_j^u(s)) ds\right]$, we immediately see that the density function of T_1 given that $U(0) = u$ is

$$f_{u, T_1}(t_1) = \sum_{j=1}^N \varphi(U_j^u(t_1)) \exp\left[-\int_0^{t_1} \sum_{j=1}^N \varphi(U_j^u(s)) ds\right], \text{ for } t_1 \geq 0. \quad (8)$$

Since $u \in B(v(0), \delta)$ and $\delta < \min\{M_i\}$, we know that there exist positive constants c_1 and c_2 such that $c_1 < u_j < c_2$ for all $j = 1, \dots, N$. But then for all $u \in B(v(0), \delta)$ and j , we have that $c_1 < U_j(t_1) = \bar{u}(1 - e^{-\lambda t_1}) + u_j e^{-\lambda t_1} < c_2$. Thus from the previous inequality and the identity (8) that it follows that $f_{u, T_1}(t_1) \geq N \varphi(c_1) e^{-t_1 N \varphi(c_2)}$. Now, from the definition of the process one easily sees that the density function of the increment T_2 given $T_1 = t_1, S_1 = 1$ and $U(0) = u$, for $t_2 > t_1$, is

$$f_{T_2|T_1=t_1, S_1=1, u}(t_2) = \sum_{j=1}^N \varphi(U_j^{\Delta_1(U^u(t_1))}(t_2)) \exp\left[-\int_{t_1}^{t_2} \sum_{j=1}^N \varphi(U_j^{\Delta_1(U^u(t_1))}(s)) ds\right]. \quad (9)$$

Note that for $j \neq 1$, $W_{1 \rightarrow j} + c_1 < \Delta_1(U^u(t_1))_j = W_{1 \rightarrow j} + U^u(t_1^-) < W_{1 \rightarrow j} + c_2$, so that $\bar{U}^u(t_1) > c_1 \frac{N-1}{N} + \frac{1}{N} \sum_{j=2}^N W_{1 \rightarrow j}$. From these two inequalities and using that the average potential is constant between successive jumps it follows that there exist positive constants

c_1^1 and c_2^1 such that $f_{T_2|T_1=t_1, S_1=1, u}(t_2) \geq \varphi(c_1^1)e^{-(t_2-t_1)N(c_2^1)}$, for $t_2 > t_1$. Proceeding in this manner we obtain sequences $(c_1^n)_{n=1, \dots, N-1}$ and $(c_2^n)_{n=1, \dots, N-1}$ satisfying for $k = 2, \dots, N$,

$$f_{T_k|T_{k-1}=t_{k-1}, S_{k-1}=k-1, \dots, T_1=t_1, S_1=1, u}(t_k) \geq \varphi(c_1^{k-1})e^{-(t_k-t_{k-1})N(c_2^{k-1})},$$

where $t_k > t_{k-1} > \dots > t_1 \geq 0$. Thus, we have that, over A_ϵ , the product of these conditional densities is positive. This concludes the prove of item (i).

(ii) The first part follows immediately from the item (i) above and corollary 3. To prove the second part, we write $\mathbb{E}[T_B(v(0), \delta)] = \int_{t=0}^{\infty} \mathbb{P}(T_B(v(0), \delta) > t)dt$ and we note that $\{T_B(v(0), \delta) > t\} \subset \{U(lT) \notin B(v(0), \delta), l = 1, \dots, \lceil \frac{t}{T} \rceil\}$ for $t > T$. A simple application of the Markov property together with the lower bound given by the first part finishes the proof. \square

Proposition 4.1. (*Harris-type Condition*) *For any $u \in B(v(0), \delta)$, there exists a non negative function h_u such that for all measurable sets $A \in \beta(\mathbb{R}_N^+ \setminus \{0^N\})$,*

$$P_T(u, A) \geq \int_A h_u(v)dv.$$

Moreover, there exist a measurable set I , with positive Lebesgue measure, and a constant $C_3 > 0$ such that $h_u(v) \geq C_3 1_I(v)$ for all $u \in B(v(0), \delta)$.

Proof. For each $u \in B(v(0), \delta)$, as in Corollary 4 let us call $\gamma_u : A_\epsilon \rightarrow I_u$, the map

$$\gamma_u(t_1^N) = \gamma^0(t_1^N) + R_{\delta_\epsilon}(t_1^N) + R_{\epsilon^2}(t_1^N),$$

where $I_u = \gamma_u(A_\epsilon)$. From the corollaries 3, 4 and the remark 4.1, it follows that for each $u \in B(v(0), \delta)$, conditioning to $A_\epsilon \cap S$, the random vector $U^u(T)$ has a density h_u , where

$$h_u(v) = \begin{cases} f_u(g_u(v))|Jg_u(v)| & , \text{ if } v \in I_u \\ 0 & \text{ otherwise,} \end{cases}$$

with $g_u : I_u \rightarrow A_\epsilon$ being the inverse of γ_u . This concludes the first part of the proposition.

The proof of second part is more delicate and requires some work. From Corollary 4 and Lemma 4.2 part (i) it suffices to prove that there exists a set I such that $I \subset \cap_{u \in B(v(0), \delta)} I_u$.

Now, consider the event B_ϵ defined by

$$B_\epsilon = \left\{ (i-1)\epsilon + \frac{\epsilon}{4} < T_i < i\epsilon - \frac{\epsilon}{4}, i = 1, \dots, N \right\} \cap S,$$

define $I = \gamma_0(B_\epsilon)$ and fix $v \in I$. We want to show that for all $u \in B(v(0), \delta)$ there is an vector $t_1^N = t_1^N(u) = (t_1(u), \dots, t_N(u)) \in A_\epsilon$ such that $\gamma_u(t_1^N) = v$. To this end, we introduce the function $F(s, t_1^N) = v - \gamma_0(t_1^N) - s[R_{\delta_\epsilon}(t_1^N, u) + R_{\epsilon^2}(t_1^N, u)]$, for $s \in [0, 1]$ and $t_1^N \in A_\epsilon$. Note that we need to show the existence of vector $t_1^N \in A_\epsilon$ such that $F(1, t_1^N) = 0$. This means that we need to study the equation $F(s, t_1^N) = 0$ with t_1^N as function of s . To ease the notation, from now on we will write t instead of t_1^N .

Note that by the definition of I , there exists $t_0 \in B_\epsilon$ such that $F(0, t_0) = 0$. Besides, $t = t(s)$ is solution of the equation $F(s, t) = 0$ if and only if it satisfies

$$0 = -D\gamma_0(t(s)) \cdot \frac{dt(s)}{ds} - [R_{\delta_\epsilon}(t(s), u) + R_{\epsilon^2}(t(s), u)] - s[DR_{\delta_\epsilon}(t(s), u) + DR_{\epsilon^2}(t(s), u)] \cdot \frac{dt(s)}{ds}$$

or equivalently,

$$\left[-D\gamma_0(t(s)) - s(DR_{\delta\epsilon}(t(s), u) + DR_{\epsilon^2}(t(s), u)) \right] \cdot \frac{dt(s)}{ds} = R_{\delta\epsilon}(t(s), u) + R_{\epsilon^2}(t(s), u), \quad (10)$$

where $Df(\cdot)$ stands for the differential operator of f . By corollary 4, the linear operator inside the brackets is invertible and the function on the right hand side is of order $\delta\epsilon + \epsilon^2$ whose the derivative is of order $\delta + \epsilon$. Therefore, it follows that $t = t(s)$ is a solution of (10) if and only if it is the solution of the ODE of the form

$$\frac{d}{ds}t(s) = H(s, t(s)), \quad t(0) = t_0, \quad (11)$$

where the derivative of H with respect to t exists and it is of order $\delta + \epsilon$. In particular, it is limited and therefore the ODE has unique solution $t = t(s)$ for all s such that $t(s) \in A_\epsilon$. Since $t(s)$ moves as H which is of order $\delta\epsilon + \epsilon^2$ and the initial condition t_0 is at a distance of order ϵ of A_ϵ , we have that $t(1) \in A_\epsilon$, decreasing both δ and ϵ if necessary. Hence, we have that $I \subset \cap_{u \in B} I_u$. \square

Proof of Theorem 4. Item (ii) of lemma 4.2 implies that the time of $B(v(0), \delta)$ is a positive recurrent set. Now by proposition 4.1 the Harris-type condition holds once the process reach $B(v(0), \delta)$, so that in fact $B(v(0), \delta)$ is a positive recurrent regenerative set. \square

5 Final discussion

The present paper presented a new class of stochastic processes and studied its asymptotic distribution in absence and presence of the leak current, without external stimuli in either cases. Similar results have been recently obtained by Robert and Touboul in [7]. Their model considers a constant leakage rate without the gap junction term ($\alpha = 1$ and $\lambda = 0$ in our context) where the synaptic weights are positive i.i.d random variables with finite mean. They showed that if $\int_0^1 \frac{\varphi(x)}{x} dx < +\infty$, then no spike occurs after some finite time. This condition imposed on the spiking rate φ is the same type as ours. They also proved that when $\varphi(0) > 0$ and the synaptic weights are bounded almost surely, the process admits a unique non trivial invariant measure. In other words, when external stimuli are considered the system remains active almost surely, contrasting with non external stimuli case. We did not study this case, nonetheless it could be treated similarly as we did in Theorem 4.

6 Acknowledges

We are indebted to R. Cofré for the discussions which originate this work. We thank E. Presutti for helpful suggestions and stimulating discussions; A. Galves and E. Löcherbach for the discussions which led us to the proof of Theorem 2.2. This article was produced as part of the activities of FAPESP Center for Neuromathematics (FAPESP grant 2013/07699-0). A. Duarte is fully supported by a CNPq fellowship (grant 141270/2013-6) and G. Ost is fully supported by a CNPq fellowship (grant 141482/2013-3).

References

- [1] M. H. A. Davis. Piecewise-deterministic Markov processes: a general class of nondiffusion stochastic models. *J. Roy. Statist. Soc. Ser. B*, 46(3):353–388, 1984.
- [2] A. De Masi, A. Galves, E. Löcherbach, and E. Presutti. Hydrodynamical limit for a system of interacting neurons. *ArXiv*, 2014.
- [3] N. Fournier and E. Löcherbach. On a toy model of interacting neurons. *ArXiv*, 2014.
- [4] A. Galves and E. Löcherbach. Infinite systems of interacting chains with memory of variable length: a stochastic model for biological neural nets. *Journal of Statistical Physics*, 151(5):896–921, 2013.
- [5] Wulfram Gerstner and Werner Kistler. *Spiking Neuron Models: An Introduction*. Cambridge University Press, New York, NY, USA, 2002.
- [6] M. Thiell, M. Riedler and G. Wainrib. Limit theorems for infinite-dimensional piecewise deterministic markov processes. applications to stochastic excitable membrane models. *Electron. J. Probab.*, 17:no. 55, 1–48, 2012.
- [7] P. Robert and J. Touboul. On the dynamics of random neuronal networks. *ArXiv*, 2014.