

Shock of three-state model for intracellular transport of kinesin KIF1A

Yunxin Zhang*

Shanghai Key Laboratory for Contemporary Applied Mathematics,

Centre for Computational System Biology,

School of Mathematical Sciences, Fudan University, Shanghai 200433, China.

Recently, a three-state model is presented to describe the intracellular traffic of unconventional (single-headed) kinesin KIF1A [Phys. Rev. Lett. **95**, 118101 (2005)], in which each motor can bind strongly or weakly to its microtubule track, and each binding site of the track might be empty or occupied by one motor. As the usual two-state model, i.e. the totally asymmetric simple exclusion process (TASEP) with motor detachment and attachment, in steady state of the system, this three-state model also exhibits shock (or domain wall separating the high-density and low density phases) and boundary layers. In this study, using mean-field analysis, the conditions of existence of shock and boundary layers are obtained theoretically. Combined with numerical calculations, the properties of shock are also studied. This study will be helpful to understand the biophysical properties of the collective transport of kinesin KIF1A.

PACS numbers: 87.16.Nn, 87.16.A-, 05.60.-k, 05.70.Ln

Keywords: TASEP; molecular motor; shock; domain wall; KIF1A

*Email: xyz@fudan.edu.cn

I. INTRODUCTION

Molecular motors are biogenic force generators acting in the nanometer range. They are responsible for intracellular transport of wide varieties of cargo from one location to another in eukaryotic cells [1–6]. Linear motors produce sliding movements along filamentous structures called protein tracks; for example, myosin slides along actin filament [7–10], kinesin [11–15] and dynein [16–20] along microtubule. Microtubule and filamentary protein actin are protein filaments which form part of a dual-purpose scaffolding called cytoskeleton, act like struts or girders for the cellular architecture and, at the same time, also serve as tracks for the intracellular transportation networks.

However, experiments found, one single filamentary track is usually traveled along by multiple motors. Fundamental understanding of these collective physical phenomena may expose the causes of motor-related diseases (e.g., Alzheimer’s disease). In literatures, these phenomena are usually described by the totally asymmetric simple exclusion process (TASEP), which is originally proposed in [21], consists of particles hopping unidirectionally with hard-core exclusion along a 1D lattice. TASEP is one of many examples for driven systems with stationary nonequilibrium states, which cannot be described in terms of Boltzmann weights. To model the attachment and detachment of motors to and from tracks, Parmeggiani *et al* [22, 23] discussed a class of driven lattice gas obtained by coupling 1D TASEP to Langmuir kinetics, in which the attachment and detachment of motors is modeled as particle creation and annihilation respectively. Furthermore, Lipowsky *et al* [24, 25] suggested a more general model, in which the diffusion of motors in the cell is considered explicitly. However, in reality, a motor protein is not a mere particle, but an enzyme whose mechanical movement is coupled with its biochemical cycle. Therefore, recently, a three-state model is presented by Nishinari *et al* to describe the intracellular transport of single-headed kinesin KIF1A [26, 27]. In which, the microtubule (MT) binding motor might be in two states: strongly MT binding state and weakly MT binding state, denoted by S, W respectively. Biochemically, the strongly binding state corresponds to bare

motor or ATP binding motor state, and the weakly binding state corresponds to ADP binding state.

One of the important feature of the collective motion of motors along one single track is the possible appearance of shock or domain wall, which is defined as the interface between the low-density and high density regions. For the usual two-state TASEP, using mean field method, the existence and properties of the shock have been discussed recently [28]. In this study, similar analysis to the Nishinari's three-state model will be presented, the efficient and necessary conditions of the existence of shock will be given theoretically, and with the aid of numerical calculations the properties of the shock will also be discussed. The method used in this study can be regarded as a generalization of the one presented in [28].

In the next section, the three-state model and its mean field approximation will be briefly introduced, and then in Sec. III the conditions of the existence of shock will be presented. The existence of boundary layers and the properties of shock will be discussed in Sec. IV and V. Finally, this study will be shortly summarized in Sec. VI.

II. TASEP WITH THREE INTERNAL STATES

The three-state TASEP given in [27] can be mathematically described as follows. Let S_i and W_i denote the probabilities of finding a molecular motor in the states 1 and 2 at the lattice site i at time t respectively (states 1 and 2 correspond to strongly bound and weakly bound states of molecular motors). Then S_i, W_i are governed by the following master equations

$$\begin{aligned} \frac{dS_i}{dt} = & \omega_a(1 - S_i - W_i) - \omega_h S_i - \omega_d S_i + \omega_s W_i \\ & + \omega_f W_{i-1}(1 - S_i - W_i) + (1 - c)\omega_f W_i(S_{i+1} + W_{i+1}), \end{aligned} \quad (1)$$

$$\begin{aligned}
\frac{dW_i}{dt} = & -(\omega_s + \omega_f)W_i(1 - S_{i+1} - W_{i+1}) + \omega_h S_i \\
& - [\omega_s + (1 - c)\omega_f]W_i(S_{i+1} + W_{i+1}) \\
& - \omega_b W_i(2 - S_{i+1} - W_{i+1} - S_{i-1} - W_{i-1}) \\
& + \omega_b(W_{i-1} + W_{i+1})(1 - S_i - W_i),
\end{aligned} \tag{2}$$

where ω_a is the rate of a molecular motor binding to the empty lattice site i , i.e. the transition rate of state 0 to state 1, ω_h is the transition rate of state 1 to state 2, i.e. the rate of ATP hydrolysis, ω_d is the transition rate of state 1 to state 0, i.e. the rate of detachment, ω_b is the rate of random Brownian motion. After the release of ADP, the motor steps forward to the next binding site in front with rate ω_f , stays at the current location with rate ω_s . c is an interpolating parameter ($0 \leq c \leq 1$). The corresponding equations for the left boundary ($i = 1$) are given by

$$\frac{dS_1}{dt} = \alpha(1 - S_1 - W_1) - \omega_h S_1 - \gamma_1 S_1 + \omega_s W_1 + (1 - c)\omega_f W_1(S_2 + W_2), \tag{3}$$

$$\begin{aligned}
\frac{dW_1}{dt} = & -(\omega_s + \omega_f)W_1 + \omega_h S_1 + c\omega_f W_1(S_2 + W_2) \\
& + \omega_b W_2(1 - S_1 - W_1) - \omega_b W_1(1 - S_2 - W_2) - \gamma_2 W_1,
\end{aligned} \tag{4}$$

where α is the rate of attachment of the motors at the left boundary (i.e. the lattice site $i = 1$), γ_1 , γ_2 are the rates of detachment of motors in state 1 and state 2 at the left boundary respectively. The equations for the right boundary ($i = N$) are give by

$$\begin{aligned}
\frac{dS_N}{dt} = & \delta(1 - S_N - W_N) + \omega_f W_{N-1}(1 - S_N - W_N) \\
& + \omega_s W_N - \omega_h S_N - \beta_1 S_N,
\end{aligned} \tag{5}$$

$$\begin{aligned}
\frac{dW_N}{dt} = & \omega_h S_N - \omega_s W_N - \beta_2 W_N + \omega_b W_{N-1}(1 - S_N - W_L) \\
& - \omega_b W_N(1 - S_{N-1} - W_{N-1}),
\end{aligned} \tag{6}$$

where δ is the rate of attachment of the motors at the right boundary (i.e. the lattice site $i = N$), β_1 β_2 are the rates of detachment of motors in state 1 and state 2 at the right boundary respectively.

It should be pointed out that the exclusion process described above is different from the one discussed in [29], where multiple occupancy of sites is allowed if particles are

in different internal states. Here, the multiple occupancy is unallowed. However, the particles bounding to the lattice site might be in two different states 1 and 2, corresponding to the strongly bound and weakly bound states. As in [22], attachment and detachment of a motor are modeled as, effectively, creation and annihilation of motors on the lattice. Moreover, the transition between states 1 and 2 is described by the rates ω_h , ω_s , ω_f , the Brownian ratched mechanism is described by rate ω_b .

Mean Field Approximation: In the large N limit, we can make the continuum mean field approximation to Eqs. (1) and (2). Let $\Delta x = \frac{1}{N-1}$ and $x = (i-1)\Delta x$. Obviously, $0 \leq x \leq 1$, since $1 \leq i \leq N$. Using the Taylor expansion

$$S(x \pm \Delta x) = S(x) + \sum_{k=1}^{+\infty} \frac{(\pm \Delta x)^k}{k!} \frac{\partial^k S(x)}{\partial x^k}, \quad W(x \pm \Delta x) = W(x) + \sum_{k=1}^{+\infty} \frac{(\pm \Delta x)^k}{k!} \frac{\partial^k W(x)}{\partial x^k}. \quad (7)$$

The continuum limits of Eqs. (1) and (2) are then

$$\begin{aligned} \frac{\partial S(x, t)}{\partial t} = & \omega_a(1 - S - W) + \omega_s W - (\omega_h + \omega_d)S \\ & + \omega_f \left(W - \Delta x \frac{\partial W(x, t)}{\partial x} \right) (1 - S - W) \\ & + (1 - c)\omega_f W \left(S + W + \Delta x \frac{\partial S(x, t)}{\partial x} + \Delta x \frac{\partial W(x, t)}{\partial x} \right) \\ & + O(\Delta x^2), \end{aligned} \quad (8)$$

$$\begin{aligned} \frac{\partial W(x, t)}{\partial t} = & c\omega_f W \left(S + W + \Delta x \frac{\partial S(x, t)}{\partial x} + \Delta x \frac{\partial W(x, t)}{\partial x} \right) \\ & - (\omega_s + \omega_f)W + \omega_h S + O(\Delta x^2). \end{aligned} \quad (9)$$

Thus, the probability density $\rho(x, t) = S(x, t) + W(x, t)$ of finding a molecular motor at lattice site x at time t satisfies [summing Eqs. (8) and (9)]

$$\frac{\partial \rho(x, t)}{\partial t} = \omega_a(1 - \rho) - \omega_d S + \omega_f \frac{\partial W(\rho - 1)}{\partial x} \Delta x + O(\Delta x^2). \quad (10)$$

As the discussion in [30], in the thermodynamic limit $N \rightarrow \infty$, there are three regimes to be distinguished. If ω_a and ω_d are of order $[1/(N-1)]^\alpha$ with $\alpha < 1$, then at the steady state, the system, Eqs. (9) and (10), reduces to

$$\begin{cases} \omega_a(1 - W - S) - \omega_d S = 0, \\ c\omega_f W (S + W) - (\omega_s + \omega_f)W + \omega_h S = 0. \end{cases} \quad (11)$$

So the probability S satisfies

$$ck(k+1)S^2 + [(k+1)(l+1) + n - c(2k+1)]S + (c-l-1) = 0, \quad (12)$$

and $W = 1 - (k+1)S$, $\rho = S + W$, where $l = \omega_s/\omega_f$, $n = \omega_h/\omega_f$, $k = \omega_d/\omega_a$ (because of the particle-hole symmetry, we restrict the discussion to the case $\omega_a \geq \omega_d$, i.e. $0 \leq k \leq 1$).

For the cases that ω_a and ω_d are of order $[1/(N-1)]^\alpha$ but with $\alpha > 1$, the local kinetics is negligible and the system will be

$$\begin{cases} W(W+S-1) = C, \\ c\omega_f W(S+W) - (\omega_s + \omega_f)W + \omega_h S = 0 \end{cases}, \quad (13)$$

where the constant C is determined by the left or right boundary conditions [31, 32]. The case of the local rates ω_a and ω_d being of the order $1/(N-1)$ is the most interesting one, and will be investigated further in this study. In the following, we always assume that the local rates ω_a and ω_d are of the order $1/(N-1)$.

III. THE EXISTENCE OF SHOCK

Let

$$\Omega_a = \frac{\omega_a}{\Delta x}, \quad \Omega_d = \frac{\omega_d}{\Delta x}, \quad (14)$$

then at steady state, the leading terms of Δx of Eqs. (9) (10) are

$$\begin{cases} \omega_f \frac{\partial W(\rho-1)}{\partial x} + \Omega_a(1-\rho) - \Omega_d S = 0 \\ c\omega_f W(S+W) - (\omega_s + \omega_f)W + \omega_h S = 0 \end{cases} \quad (15)$$

or

$$\begin{cases} \omega_f \frac{\partial W(\rho-1)}{\partial x} + \Omega_a(1-\rho) - \Omega_d(\rho-W) = 0, \\ c\omega_f W\rho - (\omega_s + \omega_f)W + \omega_h(\rho-W) = 0. \end{cases} \quad (16)$$

The second equation implies

$$W = \frac{n\rho}{n+l+1-c\rho}. \quad (17)$$

The steady state flux is then proportional to

$$J = W(1 - \rho) = \frac{n\rho(1 - \rho)}{n + l + 1 - c\rho}. \quad (18)$$

The same as in [27], at the steady state, we can obtain the left boundary conditions

$$S(0) = \frac{\alpha - [c\alpha(\alpha - \omega_s)/\omega_f]}{c\alpha + \omega_h}, \quad W(0) = \frac{\alpha}{\omega_f}, \quad (19)$$

and the right boundary conditions

$$S(1) = \frac{\omega_s + \beta}{\omega_h} \left[\frac{\omega_h}{\omega_h + \omega_s + \beta} - \frac{\beta}{\omega_f} \right], \quad W(1) = \frac{\omega_h}{\omega_h + \omega_s + \beta} - \frac{\beta}{\omega_f}. \quad (20)$$

One can find that Eqs. (16) involves only the first-order derivatives of ρ and W with respect to x whereas there are two sets of boundary conditions (19) and (20). Therefore, if we integrate the equations (16) with the left boundary conditions (19), the solution (denoted by ρ_l , W_l , S_l respectively) may not, in general, match smoothly with the solution (denoted by ρ_r , W_r , S_r respectively) obtained for the same equations but with the right boundary conditions (20). This discontinuity corresponds to a shock or domain wall. However, at any position x , the continuity condition of motor flux, or equivalently $J_l(x) = J_r(x)$, should be satisfied, where $J_l(x) = W(x-)[1 - \rho(x-)]$ and $J_r(x) = W(x+)[1 - \rho(x+)]$. At the shock position x_s ,

$$\begin{cases} W(x_s-) = W_l(x_s), & \rho(x_s-) = \rho_l(x_s), \\ W(x_s+) = W_r(x_s), & \rho(x_s+) = \rho_r(x_s). \end{cases} \quad (21)$$

So the continuity condition $J_l(x_s) = J_r(x_s)$ implies

$$\frac{n\rho_l(x_s)(1 - \rho_l(x_s))}{n + l + 1 - c\rho_l(x_s)} = J(x_s) = \frac{n\rho_r(x_s)(1 - \rho_r(x_s))}{n + l + 1 - c\rho_r(x_s)}, \quad (22)$$

or

$$\rho_l(x_s) + \rho_r(x_s) = 1 + \frac{c}{n}J(x_s) = 1 + \gamma\rho_l(x_s)\rho_r(x_s), \quad (23)$$

where $\gamma = \frac{c}{n+l+1} < 1$.

From Eqs. (16), one can show the probability ρ satisfies

$$(\gamma\rho^2 - 2\rho + 1)\rho_x = \Omega_{ah}(1 - \gamma\rho) \left[(k+1)c\rho^2 - \left(\frac{c}{\gamma} + c + k(l+1) \right) \rho + \frac{c}{\gamma} \right], \quad (24)$$

with $\Omega_{ah} = \Omega_a/\omega_h$. It can be proved that, for $0 \leq c \leq 1$ and $l \geq 0$, the discriminant

$$\Delta := \left[\frac{c}{\gamma} + c + k(l+1) \right]^2 - \frac{4(k+1)c^2}{\gamma} \geq 0.$$

So the equation (24) can be reformulated as

$$(\rho - \rho_3)(\rho - \rho_4)\rho_x = -\Omega_{ah}(k+1)c(\rho - \rho_0)(\rho - \rho_1)(\rho - \rho_2), \quad (25)$$

where

$$\begin{aligned} \rho_0 &= \frac{1}{\gamma}, \quad \rho_{3,4} = \frac{1 \mp \sqrt{1-\gamma}}{\gamma}, \\ \rho_{1,2} &= \frac{\left[\frac{c}{\gamma} + c + k(l+1) \right] \mp \sqrt{\left[\frac{c}{\gamma} + c + k(l+1) \right]^2 - \frac{4(k+1)c^2}{\gamma}}}{2(k+1)c}. \end{aligned} \quad (26)$$

One can easily show that, for $0 \leq c \leq 1, l \geq 0$ and $k \geq 0$,

$$\rho_0 \geq 1, \quad \rho_4 \geq 1, \quad \rho_2 \geq 1, \quad \rho_1 \leq 1 \quad \frac{1}{2} \leq \rho_3 \leq 1. \quad (27)$$

Particularly,

$$\begin{aligned} \rho_1 &= \frac{\frac{2c}{\gamma}}{\left[\frac{c}{\gamma} + c + k(l+1) \right] + \sqrt{\left[\frac{c}{\gamma} + c + k(l+1) \right]^2 - \frac{4(k+1)c^2}{\gamma}}} \\ &\rightarrow \begin{cases} 1 & \text{as } k \rightarrow 0, \\ \frac{l+n+1}{l+n+1+k(l+1)} \leq 1 & \text{as } c \rightarrow 0. \end{cases} \end{aligned} \quad (28)$$

Moreover, one can easily show that the function $f(x) = x - \sqrt{x^2 - x}$ decreases with $x \geq 1$ monotonously. So, for $c \leq (1-k)(l+1) + n$,

$$\begin{aligned} \rho_1 &= \frac{\left[\frac{c}{\gamma} + c + k(l+1) \right] - \sqrt{\left[\frac{c}{\gamma} + c + k(l+1) \right]^2 - \frac{4(k+1)c^2}{\gamma}}}{2(k+1)c} \\ &= \frac{(k+1)(l+1) + n + c}{2(k+1)c} - \sqrt{\left(\frac{(k+1)(l+1) + n + c}{2(k+1)c} \right)^2 - \frac{l+n+1}{(k+1)c}} \\ &\geq \frac{(k+1)(l+1) + n + c}{2(k+1)c} - \sqrt{\left(\frac{(k+1)(l+1) + n + c}{2(k+1)c} \right)^2 - \frac{(k+1)(l+1) + n + c}{2(k+1)c}} \\ &\geq \frac{l+n+1}{c} - \sqrt{\left(\frac{l+n+1}{c} \right)^2 - \frac{l+n+1}{c}} \\ &= \rho_3. \end{aligned} \quad (29)$$

Therefore, in the following, we always assume that $\rho_0, \rho_2, \rho_4 \geq 1$, and $\frac{1}{2} \leq \rho_3 \leq \rho_1 \leq 1$.

The general solutions of Eq. (25) are

$$F(\rho) = x + C, \quad (30)$$

where C is an arbitrary constant and

$$F(\rho) = -\frac{1}{\Omega_{ah}(k+1)_c} [A \ln |\rho - \rho_0| + B \ln |\rho - \rho_1| + D \ln |\rho - \rho_2|], \quad (31)$$

with

$$A = \frac{\rho_0(\rho_0 - 1)}{(\rho_0 - \rho_1)(\rho_2 - \rho_0)}, \quad B = -\frac{\rho_1^2 - 2\rho_0\rho_1 + \rho_0}{(\rho_0 - \rho_1)(\rho_1 - \rho_2)}, \quad D = \frac{\rho_2^2 - 2\rho_0\rho_2 + \rho_0}{(\rho_0 - \rho_2)(\rho_1 - \rho_2)}. \quad (32)$$

So the solution of Eq. (25), which satisfies the left boundary condition $\rho(0) = S(0) + W(0)$, see Eq. (19), is

$$F(\rho_l) = x + F_0, \quad \text{or} \quad \rho_l(x) = F^{-1}(x + F_0), \quad (33)$$

where $F_0 = F[\rho(0)]$. Similarly, the solution of Eq. (25), which satisfies the right boundary condition $\rho(1) = S(1) + W(1)$, see Eq. (20), is

$$F(\rho_r) = x + F_1 - 1, \quad \text{or} \quad \rho_r(x) = F^{-1}(x + F_1 - 1), \quad (34)$$

where $F_1 = F(\rho(1))$. In the following, we assume $\rho_l \neq \rho_r$ (otherwise, there would be no shock and boundary layers).

Combining (21) (23) (33) (34), one sees that, at the shock position x_s

$$\begin{cases} F(\rho_l(x_s)) - F(\rho_r(x_s)) + F_1 - F_0 - 1 = 0, \\ \rho_l(x_s) + \rho_r(x_s) = 1 + \gamma \rho_l(x_s) \rho_r(x_s). \end{cases} \quad (35)$$

These are the efficient and necessary condition of the existence of shock at the position x_s . In other words, at the shock position x_s , $H(x_s, \Omega_a, k, \omega_f, l, n, c) :=$

$$F(\rho_l(x_s)) - F\left(\frac{1 - \rho_l(x_s)}{1 - \gamma \rho_l(x_s)}\right) + F_1 - F_0 - 1 = 0. \quad (36)$$

If there exists $0 < x_s < 1$, such that $H(x_s, \Omega_a, k, \omega_f, l, n, c) = 0$, the shock will appear at x_s , and the height of the shock is

$$\varepsilon_s = |\rho_r(x_s) - \rho_l(x_s)| = \left| \frac{1 - 2\rho_l(x_s) + \gamma \rho_l^2(x_s)}{1 - \gamma \rho_l(x_s)} \right|. \quad (37)$$

Let

$$H(\rho) := F(\rho) - F\left(\frac{1-\rho}{1-\gamma\rho}\right) + F_1 - F_0 - 1 = 0, \quad (38)$$

then from (31), one can obtain

$$\begin{aligned} \frac{dH(\rho)}{d\rho} &= \frac{dF(\rho)}{d\rho} - \frac{d}{d\rho} F\left(\frac{1-\rho}{1-\gamma\rho}\right) = \frac{\gamma\rho^2 - 2\rho + 1}{\Omega_{ah}(1-\gamma\rho)} \\ &\times \frac{[(1-\rho) - \rho_1(1-\gamma\rho)][(1-\rho) - \rho_2(1-\gamma\rho)] + (\gamma-1)(\rho - \rho_1)(\rho - \rho_2)}{(\rho - \rho_1)(\rho - \rho_2)[(1-\rho) - \rho_1(1-\gamma\rho)][(1-\rho) - \rho_2(1-\gamma\rho)]} \\ &= \frac{\gamma(1-\rho)^2(1-\gamma\rho_1)(\rho - \rho_3)(\rho - \rho_4) \left(\frac{1-\rho_1}{1-\gamma\rho_1} - \rho_2\right)}{\Omega_{ah}(1-\gamma\rho)^3(\rho - \rho_1)(\rho - \rho_2) \left(\frac{1-\rho}{1-\gamma\rho} - \rho_1\right) \left(\frac{1-\rho}{1-\gamma\rho} - \rho_2\right)} \\ &\begin{cases} > 0 & \text{if } 0 \leq \rho < \frac{1-\rho_1}{1-\gamma\rho_1}, \\ < 0 & \text{if } \frac{1-\rho_1}{1-\gamma\rho_1} < \rho < \rho_1, \\ > 0 & \text{if } \rho_1 < \rho \leq 1. \end{cases} \end{aligned} \quad (39)$$

Using this property of the function $H(\rho)$, we can obtain the following results:

(I) For $0 \leq \rho(0) < \rho_3$ and $\rho_3 < \rho(1) \leq \rho_1$, the conditions of existence of shock in interval $(0, 1)$ is

$$F\left(\frac{1-\rho(1)}{1-\gamma\rho(1)}\right) < F_0 + 1 \quad \text{and} \quad F\left(\frac{1-\rho(0)}{1-\gamma\rho(0)}\right) < F_1 - 1, \quad (40)$$

see Fig. 1 **(a)**. From (25) (31), one can find the function $F(\rho)$ increases with ρ for $0 \leq \rho < \rho_3$ and $\rho_1 < \rho \leq 1$, and decreases with ρ for $\rho_3 < \rho < \rho_1$. Thus

$$\begin{aligned} F\left(\frac{1-\rho(1)}{1-\gamma\rho(1)}\right) < F_0 + 1 &\iff \rho_l^{-1} \left(\frac{1-\rho(1)}{1-\gamma\rho(1)}\right) < 1, \\ F\left(\frac{1-\rho(0)}{1-\gamma\rho(0)}\right) < F_1 - 1 &\iff \rho_r(0) < \frac{1-\rho(0)}{1-\gamma\rho(0)}. \end{aligned} \quad (41)$$

Therefore, the conditions presented in (40) are generalizations of the ones obtained in [28] for the usual TASEP with motor detachment and attachment:

$$\rho_l^{-1}(1-\rho(1)) < 1, \quad \text{and} \quad \rho_r(0) < 1-\rho(0). \quad (42)$$

In fact, if the parameter $c = 0$ (i.e. $\gamma = 0$), the Eq. (24) reduces to

$$(1-2\rho)\rho_x = \Omega_{ah}[(l+n+1) - [(l+n+1) + k(l+1)]\rho], \quad (43)$$

which is similar as the model discussed in [22] for the usual TASEP. For such reduced cases, $\rho_3 = 0.5$, and the conditions (40) of existence of shock is reduced to (42).

(II) For $0 \leq \rho(0) < \rho_3$ and $\rho_1 \leq \rho(1) \leq 1$, the conditions of existence of shock in interval $(0, 1)$ is

$$F\left(\frac{1-\rho(1)}{1-\gamma\rho(1)}\right) < F_0 + 1, \quad \text{and} \quad F\left(\frac{1-\rho(0)}{1-\gamma\rho(0)}\right) > F_1 - 1, \quad (44)$$

see Fig. 1 **(b)**. Similar as in **(I)**,

$$\begin{aligned} F\left(\frac{1-\rho(1)}{1-\gamma\rho(1)}\right) < F_0 + 1 &\iff \rho_l^{-1}\left(\frac{1-\rho(1)}{1-\gamma\rho(1)}\right) < 1, \\ F\left(\frac{1-\rho(0)}{1-\gamma\rho(0)}\right) > F_1 - 1 &\iff \rho_r^{-1}\left(\frac{1-\rho(0)}{1-\gamma\rho(0)}\right) > 0. \end{aligned} \quad (45)$$

Therefore, conditions (44) are also generalizations of the ones for the usual TASEP [28]:

$$\rho_l^{-1}(1-\rho(1)) < 1, \quad \text{and} \quad \rho_r^{-1}(1-\rho(0)) > 0. \quad (46)$$

(III) For $0 \leq \rho(0), \rho(1) \leq \rho_3$, the condition of the existence of shock in $(0, 1)$ is

$$\rho_l^{-1}(\rho_3) < 1 \quad \text{and} \quad \tilde{\rho}_r(0) < \frac{1-\rho(0)}{1-\gamma\rho(0)} \quad (47)$$

where $\tilde{\rho}_r(x)$ is one of the solutions of differential equation (24), which satisfies $\tilde{\rho}_r(1) = \rho_3$ and $\tilde{\rho}_r(0) > \rho_3$. See Fig. 1 **(c)**.

(IV) For $\rho(0), \rho(1) > \rho_3$, there exists no shock in $(0, 1)$. It can be readily verified that the function $f(\rho_l, \rho_r) = \rho_l + \rho_r - \gamma\rho_l\rho_r - 1$ increases monotonously with $0 \leq \rho_l, \rho_r \leq 1$, and $f(\rho_3, \rho_3) = 0$. For $\rho(0), \rho(1) > \rho_3$, one knows that $\rho_3 \leq \min(\rho_l, \rho_r) < \max(\rho_l, \rho_r)$ (note: we always assume $\rho_l \neq \rho_r$). Thus $f(\rho_l, \rho_r) = \rho_l + \rho_r - \gamma\rho_l\rho_r - 1 > 0$. It is to say that there exists no shock [see (35)].

(V) For $0 \leq \rho(1) < \rho_3$ and $\rho_l(0) > \rho_3$, there exists no shock in $(0, 1)$.

In conclusion, $\rho(0) < \rho_3$ is one necessary condition of the existence of shock in $(0, 1)$.

IV. THE EXISTENCE OF BOUNDARY LAYER

Generally speaking, if $\rho_l \neq \rho_r$ and there is no shock in $(0, 1)$, boundary layer will appear at least at one of the boundaries 0 and 1. Similar as in [28], we have the

following results:

(I) For $0 \leq \rho(0) < \rho_3$ and $\rho_3 < \rho(1) \leq \rho_1$: if

$$F\left(\frac{1-\rho(1)}{1-\gamma\rho(1)}\right) > F_0 + 1 \quad \text{and} \quad F\left(\frac{1-\rho(0)}{1-\gamma\rho(0)}\right) < F_1 - 1, \quad (48)$$

there exists boundary layer at the right boundary $x = 1$, see Fig. 1 **(d)**; if

$$F\left(\frac{1-\rho(1)}{1-\gamma\rho(1)}\right) < F_0 + 1 \quad \text{and} \quad F\left(\frac{1-\rho(0)}{1-\gamma\rho(0)}\right) > F_1 - 1, \quad (49)$$

there exists boundary layer at the left boundary $x = 0$ (in these cases, the shock position $x_s < 0$), see Fig. 2 **(b)**.

In view of the property (39) of function $H(\rho)$, the conditions (48) can be simplified as

$$F\left(\frac{1-\rho(1)}{1-\gamma\rho(1)}\right) > F_0 + 1, \quad (50)$$

and the conditions (49) can be simplified as

$$F\left(\frac{1-\rho(0)}{1-\gamma\rho(0)}\right) > F_1 - 1. \quad (51)$$

(II) For $0 \leq \rho(0) < \rho_3$ and $\rho_1 \leq \rho(1) \leq 1$: if

$$F\left(\frac{1-\rho(0)}{1-\gamma\rho(0)}\right) < F_1 - 1, \quad (52)$$

there exists boundary layer at the left boundary $x = 0$ (i.e. the shock position $x_s < 0$), see Fig. 2 **(c)**; if

$$F\left(\frac{1-\rho(1)}{1-\gamma\rho(1)}\right) > F_0 + 1, \quad (53)$$

there exists boundary layer at the right boundary $x = 1$ (i.e. the shock position $x_s > 1$), see Fig. 2 **(a)**.

(III) For $0 \leq \rho(0), \rho(1) \leq \rho_3$ and $\rho_l(1) \neq \rho(1)$, there exists boundary layer at $x = 1$.

See Fig. 1 **(c)**, Fig. 2 **(d)** and Fig. 3 **(a)**. If

$$\rho_l^{-1}(\rho_3) < 1 \quad \text{and} \quad \tilde{\rho}_r(0) > \frac{1-\rho(0)}{1-\gamma\rho(0)}, \quad (54)$$

there is also the boundary layer at the left boundary $x = 0$.

(IV) For $\rho_3 \leq \rho(0) < 1$, $0 \leq \rho(1) \leq \rho_3$ and $\rho(0) \neq \tilde{\rho}_r(0)$, there exist boundary layers at both $x = 0$ and $x = 1$, see Fig. 3 (b) and Fig. 3 (c). For these cases, $\rho(x) = \tilde{\rho}_r(x)$ for $0 < x < 1$.

(V) For $\rho_3 \leq \rho(0)$, $\rho(1) \leq 1$ and $\rho(0) \neq \rho_r(0)$, there exists boundary layer at $x = 0$. For these cases, $\rho(x) = \rho_r(x)$ for $0 < x \leq 1$, see Fig. 3 (d) and Fig. 4.

V. THE PROPERTIES OF SHOCK

Finally, we discuss the properties of shock briefly. From the discussion in Sec. III, one knows that $\rho(0) < \rho_3$ is necessary for the existence of shock. So, at the shock position x_s , $\rho_l(x_s) < \rho_r(x_s)$, see Eq. (23), and the height of the shock is [see Eq. (37)]

$$\varepsilon_s = \rho_r(x_s) - \rho_l(x_s) = \frac{1 - 2\rho_l(x_s) + \gamma\rho_l^2(x_s)}{1 - \gamma\rho_l(x_s)}. \quad (55)$$

The derivative of the height ε_s with respect to $\rho_l(x_s)$ is

$$\frac{\partial \varepsilon_s}{\partial \rho_l(x_s)} = \frac{\gamma - 1}{(1 - \gamma\rho_l(x_s))^2} - 1 < 0. \quad (56)$$

At the same time, $\rho_l(0) = \rho(0) < \rho_3$ means $\frac{\partial \rho_l(x_s)}{\partial x_s} > 0$ [see (25)]. So

$$\frac{\partial \varepsilon_s}{\partial x_s} = \frac{\partial \varepsilon_s}{\partial \rho_l(x_s)} \frac{\partial \rho_l(x_s)}{\partial x_s} < 0, \quad (57)$$

which means, the shock height ε_s decreases with the shock position x_s . Therefore, we only need to give the relations between shock position x_s and the model parameters $\Omega_a, \Omega_d, \omega_f, \omega_s, \omega_h, \omega_b, c$.

Because of the complexity of the function F [see (31)], it is difficult to get theoretical results as in [28]. From numerical calculations, we find that, the shock position x_s decreases with parameters $\Omega_a, \alpha, \omega_b, \omega_s$, but increases with parameters $\Omega_d, \beta, \omega_f, \omega_h, c$, see Fig. 5. In the calculations, x_s is obtained by (36) and (33).

VI. CONCLUDING REMARKS

In this study, the three-state process, which is presented in [26, 27] to model the intracellular transport of single-headed kinesin KIF1A, is theoretically analyzed

using mean field approximation. By similar methods as for the usual TASEP [28], the conditions of the existence of shock or domain wall, which is defined as the interface of low-density and high-density phases, are obtained. With the aid of numerical calculations, the parameters dependent properties of the shock are also discussed. The results obtained in this study will be helpful to understand the real biophysical properties of motor traffic in eukaryotic cells.

Acknowledgments

This study is funded by the Natural Science Foundation of Shanghai (under Grant No. 11ZR1403700).

-
- [1] J. Howard. *Mechanics of Motor Proteins and the Cytoskeleton*. Sinauer Associates and Sunderland, MA, 2001.
 - [2] D. Bray. *Cell movements: from molecules to motility, 2nd Edn.* Garland, New York, 2001.
 - [3] G. M. Cooper. *The Cell: A Molecular Approach, 2nd Edn.* Sinauer Associates, Inc., Sunderland, Mass., 2000.
 - [4] M. Schliwa. *Molecular Motors*. Wiley-Vch, Weinheim, 2003.
 - [5] A. O. Sperry. *Molecular Motors: Methods and Protocols (Methods in Molecular Biology Vol 392)*. Humana Press Inc., Totowa, New Jersey, 2007.
 - [6] R. D. Vale. The molecular motor toolbox for intracellular transport. *Cell*, 112:467–480, 2003.
 - [7] A. D. Mehta, R. S. Rock, M. Rief, J. A. Spudich, M. S. Mooseker, and R. E. Cheney. Myosin-V is a processive actin-based motor. *Nature*, 400:590–593, 1999.
 - [8] A. M. Hooft, E. J. Maki, K. K. Cox, and J. E. Baker. An accelerated state of myosin-based actin motility. *Biochemistry*, 46:3513–3520, 2007.

- [9] J. C. M. Gebhardt, A. E.-M. Clemen, J. Jaud, and M. Rief. Myosin-V is a mechanical ratchet. *Proc. Natl. Acad. Sci. USA*, 103:8680–8685, 2006.
- [10] K. Shiroguchi and Jr. K. Kinoshita. Myosin V walks by lever brownian motion. *Science*, 316:1208–1212, 2007.
- [11] S. M. Block, L. S. B. Goldstein, and B. J. Schnapp. Bead movement by single kinesin molecules studied with optical tweezers. *Nature*, 348:348–352, 1990.
- [12] R. D. Vale, T. Funatsu, D. W. Pierce, L. Romberg, Y. Harada, and T. Yanagida. Direct observation of single kinesin molecules moving along microtubules. *Nature*, 380:451–453, 1996.
- [13] A. Yildiz, M. Tomishige, R. D. Vale, and P. R. Selvin. Kinesin walks hand-over-hand. *Science*, 303:676–678, 2004.
- [14] N. J. Carter and R. A. Cross. Mechanics of the kinesin step. *Nature*, 435:308–312, 2005.
- [15] N. R. Guydosh and S. M. Block. Direct observation of the binding state of the kinesin head to the microtubule. *Nature*, 08259, 2009.
- [16] R. Mallik, B. C. Carter, S. A. Lex, S. J. King, and S. P. Gross. Cytoplasmic dynein functions as a gear in response to load. *Nature*, 427:649–652, 2004.
- [17] S. L. Reck-Peterson, A. Yildiz, A. P. Carter, A. Gennerich, N. Zhang, and R. D. Vale. Single-molecule analysis of dynein processivity and stepping behavior. *Cell*, 126:335–348, 2006.
- [18] S. Toba, T. M. Watanabe, L. Yamaguchi-Okimoto, Y. Y. Toyoshima, and H. Higuchi. Overlapping hand-over-hand mechanism of single molecular motility of cytoplasmic dynein. *Proc. Natl. Acad. Sci. USA*, 103:5741–5745, 2006.
- [19] A. Gennerich, A. P. Carter, S. L. Reck-Peterson, and R. D. Vale. Force-induced bidirectional stepping of cytoplasmic dynein. *Cell*, 131:952–965, 2007.
- [20] A. Houdusse and A. P. Carter. Dynein swings into action. *Cell*, 136:395–396, 2009.

- [21] C. MacDonald, J. Gibbs, and A. Pipkin. Kinetics of biopolymerization on nucleic acid templates. *Biopolymers*, 6:1–25, 1968.
- [22] A. Parmeggiani, T. Franosch, and E. Frey. Phase coexistence in driven one-dimensional transport. *Physical Review Letters*, 90:086601, 2003.
- [23] A. Parmeggiani, T. Franosch, and E. Frey. Totally asymmetric simple exclusion process with langmuir kinetics. *Phys. Rev. E*, 70:046101, 2004.
- [24] R. Lipowsky, S. Klumpp, and T.M. Nieuwenhuizen. Random walks of cytoskeletal motors in open and closed compartments. *Physical Review Letters*, 87:108101, 2001.
- [25] R. Lipowsky, Y. Chai, S. Klumpp, S. Liepelt, and M. J. I. Muller. Molecular motor traffic: From biological nanomachines to macroscopic transport. *Physica A*, 372:34–51, 2006.
- [26] K. Nishinari, Y. Okada, A. Schadschneider, and D. Chowdhury. Intracellular transport of single-headed molecular motors kif1a. *Phys.Rev. Lett.*, 95:118101, 2005.
- [27] P. Greulich, A. Garai, K. Nishinari, A. Schadschneider, and D. Chowdhury. Intracellular transport by single-headed kinesin kif1a: Effects of single-motor mechanochemistry and steric interactions. *Phys. Rev. E*, 75:041905, 2007.
- [28] Y. Zhang. Domain wall of the totally asymmetric exclusion process without particle number conservation. *Chin. J. Phys.*, 48:607–618, 2010.
- [29] T. Reichenbach, T. Franosch, and E. Frey. Exclusion processes with internal states. *Phys. Rev. Lett.* 74,, 97:050603, 2006.
- [30] V. Popkov, A. Rakos, R. D. Willmann, A. B. Kolomeisky, and G. M. Schütz. Localization of shocks in driven diffusive systems without particle number conservation. *Phys. Rev. E*, 67:066117, 2003.
- [31] B. Derrida, M. R. Evans, V. Hakim, and V. Pasquier. Exact solution of a 1d asymmetric exclusion model using a matrix formulation. *J. Phys. A: Math. Gen.*, 26:1493–1517, 1993.

- [32] G. Schütz and E. Domany. Phase transitions in an exactly soluble one-dimensional exclusion process. *J. Stat. Phys.*, 72:277–304, 1993.




FIG. 1: There exists shock in $(0, 1)$ if $0 \leq \rho(0) < \rho_3$, $\rho_3 < \rho(1) \leq \rho_1$ and $F\left(\frac{1-\rho(1)}{1-\gamma\rho(1)}\right) < F_0 + 1$, $F\left(\frac{1-\rho(0)}{1-\gamma\rho(0)}\right) < F_1 - 1$ (a); Or $0 \leq \rho(0) < \rho_3$, $\rho_1 \leq \rho(1) \leq 1$ and $F\left(\frac{1-\rho(1)}{1-\gamma\rho(1)}\right) < F_0 + 1$, $F\left(\frac{1-\rho(0)}{1-\gamma\rho(0)}\right) > F_1 - 1$ (b). There exists shock in $(0, 1)$ and boundary layer at $x = 1$ if $0 \leq \rho(0), \rho(1) \leq \rho_3$ and $\rho_l^{-1}(\rho_3) < 1$, $\tilde{\rho}_r(0) < \frac{1-\rho(0)}{1-\gamma\rho(0)}$ (c). There exists boundary layer at $x = 1$ if $0 \leq \rho(0) \leq \rho_3, \rho_3 \leq \rho(0) \leq \rho_1$ and $F\left(\frac{1-\rho(1)}{1-\gamma\rho(1)}\right) > F_0 + 1$ (d).




FIG. 2: There exists boundary layers at $x = 1$ if $0 \leq \rho(0) \leq \rho_3, \rho_1 \leq \rho(1) \leq 1$ and $F\left(\frac{1-\rho(1)}{1-\gamma^*\rho(1)}\right) > F_0 + 1$ **(a)**; Or $0 \leq \rho(0), \rho(1) \leq \rho_3$ and $\rho_l(1) > \rho(1)$ **(d)**. There exists boundary layers at $x = 0$ if $0 \leq \rho(0) < \rho_3, \rho_3 < \rho(1) \leq \rho_1$ and $F\left(\frac{1-\rho(0)}{1-\gamma^*\rho(0)}\right) > F_1 - 1$ **(b)**; Or $0 \leq \rho(0) \leq \rho_3, \rho_1 \leq \rho(1) \leq 1$ and $F\left(\frac{1-\rho(0)}{1-\gamma^*\rho(0)}\right) < F_1 - 1$ **(c)**.




FIG. 3: There exists boundary layers at $x = 1$ if $0 \leq \rho(0), \rho(1) \leq \rho_3$ and $\rho_l(1) < \rho(1)$ **(a)**. There exist boundary layers at both $x = 0$ and $x = 1$ if $\rho_3 \leq \rho(0) < 1, 0 \leq \rho(1) \leq \rho_3$ and $\rho(0) < \tilde{\rho}_r(0)$ **(b)**; Or $\rho_3 \leq \rho(0) < \rho_1, 0 \leq \rho(1) \leq \rho_3$ and $\rho(0) > \tilde{\rho}_r(0)$ **(c)**. There are boundary layers at $x = 0$ if $\rho_3 \leq \rho(0), \rho(1) \leq \rho_1$ and $\rho(0) < \rho_r(0)$ **(d)**.




FIG. 4: There are boundary layers at $x = 0$ if $\rho_3 \leq \rho(1) \leq \rho_1, \rho_1 \leq \rho(0) \leq 1$ **(a)**; Or $\rho_3 \leq \rho(0), \rho(1) \leq 1$ and $\rho(0) > \rho_r(0)$ **(b)**; Or $\rho_3 \leq \rho(0) \leq \rho_1, \rho_1 \leq \rho(1) \leq 1$ **(c)**; Or $\rho_3 \leq \rho(0), \rho(1) \leq 1$ and $\rho(0) < \rho_r(0)$ **(d)**.




FIG. 5: The shock position x_s decreases with parameters $\Omega_a, \alpha, \omega_b, \omega_s$, but increases with parameters $\Omega_d, \beta, \omega_f, \omega_h, c$. In the calculations, x_s is obtained by Eqs. (36) and (33).