

# An optimal control model of mosquito reduction management in a dengue endemic region

KARUNIA PUTRA WIJAYA<sup>a</sup>      THOMAS GÖTZ<sup>a</sup>  
EDY SOEWONO<sup>b</sup>

<sup>a</sup>Mathematical Institute, University of Koblenz  
56070 Koblenz, Germany

{karuniaputra,goetz}@uni-koblenz.de

<sup>b</sup>Department of Mathematics, Bandung Institute of Technology  
Bandung 40132, Indonesia  
esoewono@math.itb.ac.id

March 12, 2015

## Abstract

*Aedes aegypti* is known as the responsible vector in transmitting dengue flavivirus. Unavailability of medication to cure the transmission in human blood becomes a global health issue in recent decades. World epidemiologists are encouraged to focus on investigation toward an effective and inexpensive way to prevent dengue transmission, i.e. mosquito control. In this paper, we present a model depicting the dynamics of mosquito population based on indoor/outdoor life cycle classification. The basic mosquito offspring number is obtained and analysis of equilibria is shown. We bring along a discussion on application of optimal control model which engineers two simultaneous schemes. The first scheme is done by disseminating chemical like Temephos in spots where eggs and larvae develop, meanwhile the second scheme is done by conducting fumigation through areas where adult mosquitoes prevalently nest, indoor as well as outdoor. A version of gradient-based method is presented to set down a workflow in minimizing the objective functional with respect to control variable. Numerical results from analysis of the basic mosquito offspring number with constant control and from optimal control suggest that

one has to enhance the usage of fumigation rather than Temephos. It is also suggested that applying both control schemes simultaneously gives the most significant reduction to the population.

**Keywords:** mosquito population dynamics; mosquito reduction management; the basic mosquito offspring number; optimal control.

**MSC:** 49J15; 49M05; 92D30.

## 1 Introduction

It is known that *Aedes aegypti* mosquitoes be the primary transmitters of dengue fever in the world [2]. This species were predominantly tropical species and confined to coastal areas. Now they are widespread inland and cause deadly morbidities by means of dengue fever in mostly Southeast Asia, Africa, USA, Australia, Brazil, Argentina, Caribbean Islands, China, India, Japan, Portugal, etc [19, 9]. The areas where *Aedes aegypti* mosquitoes most probably prevalent in which have tremendous threat of dengue outbreaks are densely urbanized areas [3, 4]. Several studies confirmed that eradicating mosquito population had constricted the area of dengue endemicity throughout the globe, beside it helped to bring down the cost as regards a high number of surveillance. Even vector control has been considerably inexpensive, a government tendency to take out of concern the funding to attempt it continuously leads to a necessity of a more efficient and effective control management. In order the mosquito population to be precisely managed, one needs to model the dynamics of such population and further to investigate effectivity of a control scheme which is acted to it.

*Aedes aegypti* belongs to species which inhabit in domestic water containers: bath vessels, flowerpots, drums, tins, unused tyres, untreated swimming pools, or even in curved broads where it is possible for water to last on for long time [14]. Both male and female adult mosquitoes feed on nectar. Only females require additional blood sources to obtain nutrients before producing eggs and during eggs maturation. Note that a single female can lay down at one time about 100–200 immersing eggs in water [6]. Potential eggs can be produced by *Aedes aegypti* females up to five batches during their lifetime. An egg needs 2–5 days to mature its living embryo, depending on water temperature [13, 17]. In an advanced growth, each egg turns to a larva which posteriorly withstands for 5–10 days depending on water temperature and air humidity [11]. This living larva usually eats algae and microorganisms in

water surface, making each individual competes with its owns for logistics. A discussion on competitions amongst larvae can become important after the fact that in a joint container, *Aedes albopictus* larvae outcompete *Aedes aegypti* larvae, thus the winners develop at a faster rate [1]. In the prescribed range of lifetime, each larva undergoes four times skin exfoliation and turns to a pupa at the very end of the processes. An idle pupa needs to wait for 1–5 days before it metamorphoses into an adult [16]. In addition, the living adult can generally survive for 10 days, or in some extreme cases for 2–4 weeks [23].

Tabachnick et al. [20] found three polytypic origins where *Aedes aegypti* breeds: domestic (urban housing including its narrowing environment), sylvan (rural areas, some cases in tree holes and leaf axils), and peridomestic (artificial plantation areas such as coconut groves and farms). In general, separation of the origins as indoor and outdoor had also been highlighted. In [12], the authors notified that the number of *Aedes aegypti* differed based on indoor-outdoor classification and heterogeneity of containers. A brief corresponding result showed that *Aedes aegypti* species constituted as the most abundant in indoor containers compared to the other tested species. Some reference also mentioned that in indoor containers, competition amongst larvae had not always been the case since a particular tendency made *Aedes aegypti* grew more than *Aedes albopictus* [5].

By a basic idea of incorporating control measures to a mathematical model of mosquito population dynamics, several control schemes were tested toward fighting the spread of *Aedes aegypti*. We highlight, for instances, utilization of ultra-low-volume (ULV) insecticide [15] also combined with Temephos dissemination [22] and the sterile insect technique (SIT) [7, 18, 21, 10] as a Genetic-based Vector Control (GVC). However, in this paper we do not take the epidemiology of mosquito population or infection exposures amongst individuals into account. Therefore, a susceptible-infected segregation is no longer in use. We accentuate a model of mosquito population dynamics based on indoor-outdoor life cycle classification and introduction of control intervention as well as investigation to the best strategy for reducing the population with cost as cheap as possible. As control devices, we highlight utilization of Temephos (mainly used to kill larvae) and fumigation (used to kill adults by blocking respiration process) after the fact that they have already been well-known schemes in an integrated mosquito eradication programme.

We organize the rest of the paper as the following. In Section 2, we set up a mathematical model for depicting the dynamics of multi-age-segregated mosquito population based on indoor-outdoor life cycle classification. We add two control measures to the system in order to check how the popu-

lation trajectories response to such control. In Section 3, we examine the biological meaningfulness of the model based on positivity of the system's solution and existence and stability of equilibria. So far we use plausible constant value for the control. In Section 4, we discuss the optimal control model which generates a time-variant control solving the proposed optimization problem. In seeking an optimal solution, we first use the so-called indirect method to generate state-adjoint-gradient systems of equations and then utilize a gradient-based method to solve the generated systems in an algorithmic workflow. As the rest of the paper, in Section 5, we do numerical tests for bringing forward some visualizations of the model.

## 2 The mathematical model

To capture the dynamics of mosquito population, let us first classify the population based on age-segregation: egg  $\mathbf{E}$ , larva  $\mathbf{L}$  and adult  $\mathbf{A}$ . We out-concern the dynamics of pupa by our assumption that there will be no deaths and imbalance in- and outflow occurring amongst pupae, since then the population of pupa remains constant. Based on indoor-outdoor classification, the eggs and larvae differ from indoor  $\mathbf{E}_1, \mathbf{L}_1$  and outdoor  $\mathbf{E}_2, \mathbf{L}_2$ . No more segregation for the adults since every individual can fly to wherever it prevails, indoor as well as outdoor. We introduce the control measures  $\mathbf{u}_1$  and  $\mathbf{u}_2$  denoting the impact rate of utilization of Temephos dissemination and fumigation respectively. Let  $[0, T]$  be the range of observation time  $t$  and  $\mathbf{x} = (\mathbf{E}_1, \mathbf{E}_2, \mathbf{L}_1, \mathbf{L}_2, \mathbf{A})^\top \in L^2([0, T], \mathbb{R}^5)$  be the state variable. Our model is preliminarily exhibited by the following equation

$$\dot{\mathbf{x}}(t) = \mathbf{f}(\mathbf{x}(t), \mathbf{u}(t)), \quad \mathbf{x}(0) = \mathbf{x}_0 \succeq \mathbf{0}. \quad (1)$$

We assume that adult mosquitoes will select indoor breeding sites with the probability  $p$  and, therefore, select outdoor breeding sites with the probability  $1 - p$ . Alongside with introduction of the tendency-based probability, we denote  $\mu$  as the corresponding rate of the adults to lay eggs. It is assumed that natural deaths can occur in all classes, thus we denote  $\eta_{\{1, \dots, 5\}}$  as the corresponding rates. In average periods, both indoor and outdoor eggs metamorphose into corresponding larvae with the transition rates  $\alpha_{\{1, 2\}}$ . The same situations hold for respectively indoor and outdoor larvae, that they metamorphose into adults with the transition rates  $\beta_{\{1, 2\}}$ . As an underlying discussion, we introduce the two logistic coefficients  $\sigma_{\{1, 2\}}$  to accommodate phenomena of competition amongst larvae. Assume that all in- or outdoor water containers are treated as homogeneous such that  $\frac{\sigma_1}{\sigma_2}$  proportionates to

some constant  $M$ . The fact that suitable choice for  $M > 1$  leads to the situation where outdoor breeding sites have  $M$ -times larger carrying capacity than indoor. After all, the model (1) is unfolded as

$$\dot{\mathbf{E}}_1 = \mu p \mathbf{A} - (\alpha_1 + \eta_1) \mathbf{E}_1 - q \mathbf{u}_1 \mathbf{E}_1 \quad (2a)$$

$$\dot{\mathbf{E}}_2 = \mu(1 - p) \mathbf{A} - (\alpha_2 + \eta_2) \mathbf{E}_2 \quad (2b)$$

$$\dot{\mathbf{L}}_1 = \alpha_1 \mathbf{E}_1 - \sigma_1 \mathbf{L}_1^2 - (\beta_1 + \eta_3) \mathbf{L}_1 - \mathbf{u}_1 \mathbf{L}_1 \quad (2c)$$

$$\dot{\mathbf{L}}_2 = \alpha_2 \mathbf{E}_2 - \sigma_2 \mathbf{L}_2^2 - (\beta_2 + \eta_4) \mathbf{L}_2 \quad (2d)$$

$$\dot{\mathbf{A}} = \beta_1 \mathbf{L}_1 + \beta_2 \mathbf{L}_2 - \eta_5 \mathbf{A} - \mathbf{u}_2 \mathbf{A}. \quad (2e)$$

Two control measures are added into the model. Temephos is practically disseminated into indoor water containers to kill larvae and eggs. The weighting factor  $q \in [0, \frac{1}{2}]$  accounts for the fact that the dissemination has less impact on eggs. Meanwhile fumigation is directly targeted to adult mosquito population.

It is assumed that all the control  $\mathbf{u}$  belongs to a set of admissible control  $\mathcal{U} \subset L^2([0, T], \mathbb{R}_+^2)$ . Define

$$\mathcal{D} := \{\mathbf{x} : \dot{\mathbf{x}}(t) = \mathbf{f}(\mathbf{x}(t), \mathbf{u}(t)), t \in [0, T], \mathbf{x}(0) = \mathbf{x}_0 \succeq \mathbf{0}, \mathbf{u} \in \mathcal{U}\} \quad (3)$$

as the feasible set of solutions. It is intuitively believed that the higher  $\mathbf{u}$  plugged into the model, the lesser the number of all classes. On the other hand, the lesser  $\mathbf{u}$ , the more the population tends to explosion. The more  $\mathbf{u}$  means that a policy maker needs to spend more fund, otherwise there will be no significant reduction to the mosquito population size. Given positive trade-off constants  $\omega_{\mathbf{x}, \{1, \dots, 5\}}$  and  $\omega_{\mathbf{u}, \{1, 2\}}$ , the following objective functional accommodates the necessity of balancing situation between significant population reduction and limitation of fund:

$$J(\mathbf{u}) := \frac{1}{2T} \int_0^T \sum_{i=1}^5 \omega_{\mathbf{x}, i} \mathbf{x}_i^2(t) + \sum_{j=1}^2 \omega_{\mathbf{u}, j} \mathbf{u}_j^2(t) dt. \quad (4)$$

Now our optimization problem is stated as

$$(\text{OC}): \text{find } (\mathbf{x}, \mathbf{u}) \in \mathcal{D} \times \mathcal{U} \text{ such that } J(\mathbf{u}) \rightarrow \min. \quad (5)$$

### 3 Model analysis

We need to ensure that our model is biologically meaningful. The following theorem gives a primary meaningfulness of the model: whenever the initial condition is positive, then evolution of the solution points in forward time stays positive.

**Theorem 1** Consider the model (1) where  $\mathbf{u} \in \mathcal{U}$ . If  $\mathbf{x}_0 \succeq \mathbf{0}$  then  $\mathbf{x}(t) \succeq \mathbf{0}$  for all  $t > 0$ , or particularly on the observation range  $(0, T]$ .

**Proof** Following the steps in [22], we let  $\mathbf{n}$  be  $(5 \times 5)$ -matrix representing a collection of all normal vectors (by rows) to the boundary of nonnegative orthant  $\partial\mathbb{R}_+^5$ . Thus we have  $\mathbf{n} = -I_5$  where  $I_5$  denotes the identity matrix. To ensure that the trajectory of solution does not walk out of the nonnegative orthant, one only needs to check the trajectory of points in the boundary. Notice that at  $i$ -th boundary,  $\partial_i\mathbb{R}_+^5$ ,

$$[\mathbf{n}\mathbf{f}(\mathbf{x}, \mathbf{u})]_i|_{\mathbf{x} \in \partial_i\mathbb{R}_+^5, \mathbf{u} \in \mathcal{U}} \leq 0.$$

This means the direction of evolution of such points in a boundary is in counter-direction or at least perpendicular to the corresponding normal vector. Thus it follows that the solution must not leave  $\mathbb{R}_+^5$  for all  $t > 0$ .  $\square$

Let us first consider the autonomous system of nonlinear differential equations (1) with constant control  $\mathbf{u} \in \mathbb{R}_+^2$ . For abbreviate purpose, let  $s_1 = \mu p$ ,  $s_2 = \mu(1 - p)$ ,  $s_3 = \alpha_1$ ,  $s_4 = \alpha_2$ ,  $s_5 = \beta_1$ ,  $s_6 = \beta_2$  and  $d_1 = \alpha_1 + \eta_1 + q\mathbf{u}_1$ ,  $d_2 = \alpha_2 + \eta_2$ ,  $d_3 = \beta_1 + \eta_3 + \mathbf{u}_1$ ,  $d_4 = \beta_2 + \eta_4$ ,  $d_5 = \eta_5 + \mathbf{u}_2$ . In order to obtain equilibria of the system, we need to solve  $\mathbf{f}(\mathbf{x}, \mathbf{u}) = \mathbf{0}$ . It follows from this process, one holds

$$\mathbf{A}^* = \frac{s_5\mathbf{L}_1^* + s_6\mathbf{L}_2^*}{d_5} \quad (6a)$$

$$\mathbf{E}_1^* = \frac{s_1(s_5\mathbf{L}_1^* + s_6\mathbf{L}_2^*)}{d_1d_5} \quad (6b)$$

$$\mathbf{E}_2^* = \frac{s_2(s_5\mathbf{L}_1^* + s_6\mathbf{L}_2^*)}{d_2d_5} \quad (6c)$$

where  $(\mathbf{L}_1^*, \mathbf{L}_2^*)$  follows the leading equations

$$\mathbf{L}_1^2 + a_1\mathbf{L}_1 + b_1\mathbf{L}_2 = 0 \quad (6d)$$

$$\mathbf{L}_2^2 + a_2\mathbf{L}_2 + b_2\mathbf{L}_1 = 0. \quad (6e)$$

All the constants that belong to the last equations are given by

$$a_1 = \frac{d_3}{\sigma_1} - \frac{s_1s_3s_5}{d_1d_5\sigma_1}, b_1 = -\frac{s_1s_3s_6}{d_1d_5\sigma_1}, a_2 = \frac{d_4}{\sigma_2} - \frac{s_2s_4s_6}{d_2d_5\sigma_2}, b_2 = -\frac{s_2s_4s_5}{d_2d_5\sigma_2}$$

It follows from (6d)-(6e), three equilibria can simply be obtained

$$\begin{aligned} \mathbf{Q}_1 &= (0, 0, 0, 0, 0) \\ \mathbf{Q}_2 &= \left( -\frac{s_1s_5a_1}{d_1d_5}, -\frac{s_2s_5a_1}{d_2d_5}, -a_1, 0, -\frac{s_5a_1}{d_5} \right) \\ \mathbf{Q}_3 &= \left( -\frac{s_1s_6a_2}{d_1d_5}, -\frac{s_2s_6a_2}{d_2d_5}, 0, -a_2, -\frac{s_6a_2}{d_5} \right). \end{aligned}$$

Definition of  $\mathbf{Q}_2$  and  $\mathbf{Q}_3$  only makes sense in biological point of view if only  $a_1, a_2$  are negative. The definition simply means that after very long time, the class of either in- or outdoor larvae tends to extinction while the other classes stay alive. In biological context, such situation can hardly happen. In the next writing, we demonstrate that the choice of  $a_1, a_2$  to be positive leads to a more interesting discussion. First, we introduce a measure whose cubic value is given by

$$\mathcal{R}(\mathbf{u})^3 := \frac{s_1 s_3 s_5}{d_5 d_1 d_3} + \frac{s_2 s_4 s_6}{d_5 d_2 d_4}. \quad (7)$$

The following identity is found after some algebraic computations

$$\frac{b_1 b_2 - a_1 a_2}{d_3 d_4} \sigma_1 \sigma_2 = \mathcal{R}(\mathbf{u})^3 - 1. \quad (8)$$

Using unity as a threshold value, the following theorem justifies the stability of zero equilibrium  $\mathbf{Q}_1$  by considering the nominal of  $\mathcal{R}(\mathbf{u})$  relative to the threshold. Meanwhile, the next theorem shows the existence of a positive nontrivial equilibrium (also well-known as coexistence equilibrium).

**Theorem 2** *The zero equilibrium  $\mathbf{Q}_1$  is locally asymptotically stable if  $\mathcal{R}(\mathbf{u}) < 1$  and is unstable if  $\mathcal{R}(\mathbf{u}) > 1$ .*

**Proof** Checking the stability of the zero equilibrium in local view, it is similar to see behavior of solution of the linearized system (1) in the neighborhood of  $\mathbf{Q}_1$ , i.e.  $\dot{\mathbf{x}} = J(\mathbf{Q}_1, \mathbf{u})(\mathbf{x} - \mathbf{Q}_1)$ . In this case,  $J(\mathbf{Q}_1, \mathbf{u})$  stands for Jacobian of  $\mathbf{f}$  evaluated at  $\mathbf{Q}_1$ . Unfolding this  $J(\mathbf{Q}_1, \mathbf{u})$ , we get

$$J(\mathbf{Q}_1, \mathbf{u}) = \begin{bmatrix} -d_1 & 0 & 0 & 0 & s_1 \\ 0 & -d_2 & 0 & 0 & s_2 \\ s_3 & 0 & -d_3 & 0 & 0 \\ 0 & s_4 & 0 & -d_4 & 0 \\ 0 & 0 & s_5 & s_6 & -d_5 \end{bmatrix}.$$

$\mathbf{Q}_1$  is locally asymptotically stable if and only if all eigenvalues of  $J(\mathbf{Q}_1, \mathbf{u})$  have negative real part. To see this, a simple cofactorization method computes determinant of  $J(\mathbf{Q}_1, \mathbf{u}) - \lambda I_5$  as the sum of all its cofactors, given by negative of

$$(d_1 + \lambda)(d_2 + \lambda)(d_3 + \lambda)(d_4 + \lambda)(d_5 + \lambda) - s_2 s_4 s_6 (d_1 + \lambda)(d_3 + \lambda) - s_1 s_3 s_5 (d_2 + \lambda)(d_4 + \lambda).$$

Clearly  $\text{coeff}(\lambda^{\{3,4,5\}})$  are real positive, meanwhile

$$\begin{aligned} \text{coeff}(\lambda^0) &= d_1 d_2 d_3 d_4 d_5 - s_1 s_3 s_5 d_2 d_4 - s_2 s_4 s_6 d_1 d_3 \\ \text{coeff}(\lambda^1) &= d_1 d_2 d_3 d_4 + \cdots + d_2 d_3 d_4 d_5 - s_1 s_3 s_5 (d_2 + d_4) - s_2 s_4 s_6 (d_1 + d_3) \\ \text{coeff}(\lambda^2) &= d_1 d_2 d_3 + \cdots + d_3 d_4 d_5 - s_1 s_3 s_5 - s_2 s_4 s_6. \end{aligned}$$

If  $\mathcal{R}(\mathbf{u}) < 1$ , or equivalently  $\frac{s_1 s_3 s_5}{d_5 d_1 d_3} + \frac{s_2 s_4 s_6}{d_5 d_2 d_4} < 1$ , one can see that  $\mathbf{coeff}(\lambda^{\{0,1,2,3,4,5\}})$  are real positive. Since  $\lim_{\lambda \rightarrow \pm\infty} -\mathbf{det}(J(\mathbf{Q}_1, \mathbf{u}) - \lambda I_5) = \pm\infty$ ,  $-\mathbf{det}(J(\mathbf{Q}_1, \mathbf{u}) - \lambda I_5)|_{\lambda=0} > 0$  and

$$-\left. \frac{d\mathbf{det}(J(\mathbf{Q}_1, \mathbf{u}) - \lambda I_5)}{d\lambda} \right|_{\lambda>0} > 0, \quad -\left. \frac{d^2\mathbf{det}(J(\mathbf{Q}_1, \mathbf{u}) - \lambda I_5)}{d\lambda^2} \right|_{\lambda>0} > 0,$$

then  $\mathbf{det}(J(\mathbf{Q}_1, \mathbf{u}) - \lambda I_5)$  can not have zero that has positive real part. Moreover, all zeros of  $\mathbf{det}(J(\mathbf{Q}_1, \mathbf{u}) - \lambda I_5)$  have negative real part. If  $\mathcal{R}(\mathbf{u}) > 1$ , then  $\mathbf{coeff}(\lambda^0) < 0$ . Then for however the values of  $\mathbf{coeff}(\lambda^1)$  and  $\mathbf{coeff}(\lambda^2)$ , it follows that there must be at least one real positive zero. This means  $\mathbf{Q}_1$  is unstable. If  $\mathcal{R}(\mathbf{u}) = 1$ , then a similar step can prove that  $\mathbf{det}(J(\mathbf{Q}_1, \mathbf{u}) - \lambda I_5)$  has one zero with the value 0 and the other zeros remain with negative real part. In this case,  $\mathbf{Q}_1$  is considerably still stable but not asymptotical.  $\square$

**Theorem 3** *If  $a_1$  and  $a_2$  are positive, a unique coexistence equilibrium  $\mathbf{Q}_4$  exists if  $\mathcal{R}(\mathbf{u}) > 1$  and does not if  $\mathcal{R}(\mathbf{u}) \leq 1$ . If  $a_1$  and  $a_2$  are negative, then there always exists a coexistence equilibrium for any value of  $\mathcal{R}(\mathbf{u})$ . If else, then it always holds  $\mathcal{R}(\mathbf{u}) > 1$ , and therefore, there always exists a coexistence equilibrium.*

**Proof** One simple substitution in (6d)-(6e) makes the equilibrium state  $\mathbf{L}_1$  to follow

$$\mathbf{L}_1^4 + 2a_1\mathbf{L}_1^3 + (a_1^2 - b_1a_2)\mathbf{L}_1^2 + (b_1^2b_2 - a_1b_1a_2)\mathbf{L}_1 = 0.$$

As  $\mathbf{L}_1 = 0$  results to equilibria that has been discussed before, now let us consider the remaining cubic

$$P(\mathbf{L}_1) := \mathbf{L}_1^3 + 2a_1\mathbf{L}_1^2 + (a_1^2 - b_1a_2)\mathbf{L}_1 + (b_1^2b_2 - a_1b_1a_2).$$

Since coefficient of  $\mathbf{L}_1^3$  is  $1 > 0$ , then  $\lim_{\mathbf{L}_1 \rightarrow \pm\infty} P = \pm\infty$ . We prove this theorem by parity.

**Case 1**  $a_1$  and  $a_2$  are positive. In this case, three possibilities are carried out:  $\mathcal{R}(\mathbf{u}) > 1$ ,  $\mathcal{R}(\mathbf{u}) < 1$  and  $\mathcal{R}(\mathbf{u}) = 1$ . If  $\mathcal{R}(\mathbf{u}) > 1$ , then by identity (8) it holds that  $b_1b_2 > a_1a_2$ . Thus we have  $0 > b_1^2b_2 - a_1b_1a_2 = P(0)$  since  $b_1 < 0$ . It follows from the value of  $P(0)$  and  $\lim_{\mathbf{L}_1 \rightarrow \infty} P = \infty$  that  $P$  must have at least one positive root. Now we have to check that  $\mathbf{L}_2 = \frac{\mathbf{L}_1(\mathbf{L}_1 + a_1)}{-b_1} > 0$ . To see this, let  $\mathbf{L}_1 = \rho^2$  be the corresponding root, where  $\rho \in \mathbb{R} \setminus \{0\}$ . Extracting  $(\mathbf{L}_1 - \rho^2)$  out of the cubic, we get  $\mathbf{L}_1^2 + (2a_1 + \rho^2)\mathbf{L}_1 + \mathcal{N}$  where

$$\mathcal{N} := (a_1^2 - b_1a_2) + \rho^2(2a_1 + \rho^2).$$



Since  $\mathcal{N} = \frac{-(b_1^2 b_2 - a_1 b_1 a_2)}{\rho^2} > 0$  then together with the claim of uniqueness of positive root  $\mathbf{L}_1 = \rho^2$ , we need  $(2a_1 + \rho^2) > 0$  to ensure that the remaining quadratic function does not have real positive root. Since  $a_1 > 0$ , then  $(\rho^2 + a_1) > 0$ . Now we have proved that  $\mathbf{L}_2$  is positive, and by (6), the other states are also positive. If  $\mathcal{R}(\mathbf{u}) < 1$ , then  $P(0) > 0$ . Since  $\lim_{\mathbf{L}_1 \rightarrow -\infty} P = -\infty$ , this means there exists at least one negative root  $\mathbf{L}_1 = -\varepsilon^2$  of the cubic, where  $\varepsilon \in \mathbb{R} \setminus \{0\}$ . Now extract  $(\mathbf{L}_1 + \varepsilon^2)$  out of the cubic and we get the remaining quadratic function  $\mathbf{L}_1^2 + (2a_1 - \varepsilon^2)\mathbf{L}_1 + \mathcal{M}$  where

$$\mathcal{M} := (a_1^2 - b_1 a_2) - \varepsilon^2(2a_1 - \varepsilon^2).$$

It turns out that  $\mathcal{M} = \frac{b_1^2 b_2 - a_1 b_1 a_2}{\varepsilon^2} = \frac{P(0)}{\varepsilon^2} > 0$ . Thus further analysis confirms that a positive root exists whenever  $(2a_1 - \varepsilon^2)$  is strictly less than zero and the discriminant  $\Delta = (2a_1 - \varepsilon^2)^2 - 4\mathcal{M} \geq 0$ . One can see that  $\Delta = (2a_1 - \varepsilon^2)^2 - 4(a_1^2 - b_1 a_2) + 4\varepsilon^2(2a_1 - \varepsilon^2) = (2a_1 + 3\varepsilon^2)(2a_1 - \varepsilon^2) - 4(a_1^2 - b_1 a_2) < 0$  since  $a_1, a_2 > 0$  and  $b_1 < 0$ . This contradicts the necessity that  $\Delta \geq 0$ . Then there does not exist positive root of the cubic. At last, if  $\mathcal{R}(\mathbf{u}) = 1$ , then  $b_1^2 b_2 - a_1 b_1 a_2 = 0$  and we get the remaining quadratic  $\mathbf{L}_1^2 + 2a_1 \mathbf{L}_1 + (a_1^2 - b_1 a_2)$ . Since  $a_1, a_2 > 0$  and  $b_1 < 0$ , it is clear that the quadratic does not have any real positive root.

**Case 2**  $a_1$  and  $a_2$  are negative. If  $\mathcal{R}(\mathbf{u}) > 1$ , then it is easily seen that one positive root exists. Now we have to check that  $\mathbf{L}_2 = \frac{\mathbf{L}_1(\mathbf{L}_1 + a_1)}{-b_1} > 0$ . To see this, as it is previously done together with the claim of uniqueness of positive root  $\mathbf{L}_1 = \rho^2$ , we need  $(2a_1 + \rho^2) > 0$  to ensure that the remaining quadratic function does not have real positive root. Then  $(\rho^2 + a_1) > (2a_1 + \rho^2) > 0$  since  $a_1 < 0$ . Now we have proved that  $\mathbf{L}_2$  is positive, and therefore, the other states are also positive. If  $\mathcal{R}(\mathbf{u}) = 1$ , we have  $b_1^2 b_2 - a_1 b_1 a_2 = 0$  and the remaining quadratic function is  $\mathbf{L}_1^2 + 2a_1 \mathbf{L}_1 + (a_1^2 - b_1 a_2)$ . Since  $\Delta = 4a_1^2 - 4(a_1^2 - b_1 a_2) = 4b_1 a_2 > 0$  and  $2a_1 < 0$ , then a positive root exists whose uniqueness is confirmed by the value of  $(a_1^2 - b_1 a_2)$ . Thus we need to proof that  $\mathbf{L}_2 = \frac{\mathbf{L}_1(\mathbf{L}_1 + a_1)}{-b_1} > 0$ . Let  $\mathbf{L}_1 = \eta^2$  be such root, then the other root must be  $\mathbf{L}_1 = -(2a_1 + \eta^2)$  where  $(2a_1 + \eta^2) = -\frac{(a_1^2 - b_1 a_2)}{\eta^2}$ . However, the positive root is unique if  $(a_1^2 - b_1 a_2) \leq 0$ , then  $\eta^2 + a_1 > \eta^2 + 2a_1 = -\frac{(a_1^2 - b_1 a_2)}{\eta^2} \geq 0$ . This confirms that  $\mathbf{L}_2 > 0$ . If  $\mathcal{R}(\mathbf{0}) < 1$ , then as before, we have at least one negative root  $\mathbf{L}_1 = -\varepsilon^2$  and the remaining quadratic  $\mathbf{L}_1^2 + (2a_1 - \varepsilon^2)\mathbf{L}_1 + \mathcal{M}$ . Observe that  $\mathcal{M} > 0$  and  $(2a_1 - \varepsilon^2) < 0$  since  $a_1 < 0$ . Some factorization confirms that there exist two remaining positive roots  $\mathbf{L}_1^{(1,2)} = \{\rho^2, \xi^2\}$  where  $\xi^2 = -(2a_1 - \varepsilon^2 + \rho^2)$  and  $\rho^2 \xi^2 = \mathcal{M}$ . Then we have to check the result of  $\mathbf{L}_1^{(1,2)} + a$  since this confirms positivity of  $\mathbf{L}_2$ , or furthermore, existence of positive equilibrium. Since  $(\rho^2 + a_1) + (\xi^2 + a_1) = -(2a_1 - \varepsilon^2) + 2a_1 = \varepsilon^2 > 0$ ,

this ensures that at least one of the roots is greater than  $-a_1$ , since then at least one root makes  $\mathbf{L}_2$  positive.

**Case 3** Else. Since it always holds  $b_1 b_2 > a_1 a_2$ , then this implies  $\mathcal{R}(\mathbf{u}) > 1$  and  $P(0) = b_1^2 b_2 - a_1 b_1 a_2 < 0$ . Therefore, there is always (not necessarily unique) coexistence equilibrium. This completes the proof.  $\square$

All classes have responsibility to generate offsprings in sense that new-borns are impossible whenever one class in the system remains zero for all time. In line with this sense, we define the mosquito-free equilibrium (MFE) as a static condition when all classes can not reproduce for the next offsprings. This means MFE is equivalent to the zero equilibrium  $\mathbf{Q}_1$ . Following the next generation method [8], we define two matrices  $F(\mathbf{u}), V(\mathbf{u})$  where  $F(\mathbf{u})$  is a matrix with zeros in the main diagonal and  $V(\mathbf{u})$  is a positive diagonal matrix such that  $J(\mathbf{Q}_1, \mathbf{u}) = F(\mathbf{u}) - V(\mathbf{u})$ . Then we define the so-called *next generation matrix*  $\mathbf{G}(\mathbf{u}) = F(\mathbf{u})V(\mathbf{u})^{-1}$  which is nothing but

$$\mathbf{G}(\mathbf{u}) = \begin{pmatrix} 0 & 0 & 0 & 0 & \frac{s_1}{d_5} \\ 0 & 0 & 0 & 0 & \frac{s_2}{d_5} \\ \frac{s_3}{d_1} & 0 & 0 & 0 & 0 \\ 0 & \frac{s_4}{d_2} & 0 & 0 & 0 \\ 0 & 0 & \frac{s_5}{d_3} & \frac{s_6}{d_4} & 0 \end{pmatrix}. \quad (9)$$

$(i, j)$ -th element of this matrix represents the average number of new individuals in compartment  $i$  produced by a single individual from compartment  $j$  during compartment  $j$ 's average individual lifetime period. It is easy to see that the spectral radius of  $\mathbf{G}(\mathbf{u})$  is given by

$$\max\{|\rho| : \mathbf{u} \in \mathbb{R}_+^2, \det(\mathbf{G}(\mathbf{u}) - \rho I_5) = 0\} = \mathcal{R}(\mathbf{u}). \quad (10)$$

This  $\mathcal{R}(\mathbf{u})$  is often called as *the basic mosquito offspring number*. Formulation in (7) provides dependency of the basic mosquito offspring number on several parameters and the control  $\mathbf{u}$ . In Section 5, we check behaviour of the basic mosquito offspring number with respect to the control and to several unknown empirical parameters.

**Theorem 4** *The existing coexistence equilibrium  $\mathbf{Q}_4$  is globally asymptotically stable in the nonnegative orthant  $\mathbb{R}_+^5$  if  $\mathcal{R}(\mathbf{u}) > 1$ .*

**Remark** We omit writing the detail for the proof of the theorem. Furthermore, it requires the use of the so-called Krasovskii–LaSalle principle. As another reference, one can further see from numerical results that the solution of the model is monotonic and tends asymptotically to  $\mathbf{Q}_4$ , by supplemented condition that the set of parameters is chosen such that  $\mathcal{R}(\mathbf{u}) > 1$ .

## 4 Optimal control problem

Recall our optimization problem:

$$\text{find } (\mathbf{x}, \mathbf{u}) \in \mathcal{D} \times \mathcal{U} \text{ such that } J(\mathbf{u}) \rightarrow \min. \quad (\text{OC})$$

Denote by  $\mathcal{U}$  a set of admissible controls where it is assumed to be compact. Let  $\mathbf{g}$  be a functional such that  $J(\mathbf{u}) = \int_0^T \mathbf{g}(\mathbf{x}, \mathbf{u}) dt$ . Let  $\bar{\mathbf{u}}$  be an optimal control that solves (OC). Consider a small variation around the optimal control

$$\mathbf{u}^\epsilon(t) = \bar{\mathbf{u}}(t) + \epsilon \kappa(t), \quad |\epsilon| \ll 1. \quad (11)$$

Plugging this variation into (1) together with positive initial condition, we have the resulting perturbed state  $\mathbf{x}^\epsilon$ , where it holds

$$\dot{\mathbf{x}}^\epsilon - \dot{\bar{\mathbf{x}}} = \mathbf{f}(\mathbf{x}^\epsilon, \mathbf{u}^\epsilon) - \mathbf{f}(\bar{\mathbf{x}}, \bar{\mathbf{u}}). \quad (12)$$

Note that  $\mathbf{f} \in C^1(\mathbb{R}^5 \times \mathbb{R}^2, \mathbb{R}^5)$ , thus we have the following Taylor approach

$$\frac{d}{dt}(\mathbf{x}^\epsilon - \bar{\mathbf{x}}) = \mathbf{f}_{\mathbf{x}}(\bar{\mathbf{x}}, \bar{\mathbf{u}})(\mathbf{x}^\epsilon - \bar{\mathbf{x}}) + \epsilon \mathbf{f}_{\mathbf{u}}(\bar{\mathbf{x}}, \bar{\mathbf{u}})\kappa + \mathcal{O}(\epsilon^2). \quad (13)$$

Working at  $\mathcal{O}(\epsilon^2)$ , we let  $\varphi$  be the solution of the differential equation

$$\dot{\varphi} = \mathbf{f}_{\mathbf{x}}(\bar{\mathbf{x}}, \bar{\mathbf{u}})\varphi + \mathbf{f}_{\mathbf{u}}(\bar{\mathbf{x}}, \bar{\mathbf{u}})\kappa, \quad \varphi(0) = \mathbf{0}. \quad (14)$$

Thus some algebraic computations show that our perturbed state is given by

$$\mathbf{x}^\epsilon = \bar{\mathbf{x}} + \epsilon \varphi + \mathcal{O}(\epsilon^2). \quad (15)$$

Further it can be proved that the solution of (14) exists and therefore the term  $\frac{\partial \mathbf{x}^\epsilon}{\partial \epsilon}$  exists.

Let  $\mathbf{z} \in L^2([0, T], \mathbb{R}^5)$  be some dual variable. It follows that  $\int_0^T \frac{d}{dt}(\mathbf{z} \cdot \mathbf{x}^\epsilon) dt = \mathbf{z}(T) \cdot \mathbf{x}^\epsilon(T) - \mathbf{z}(0) \cdot \mathbf{x}_0$ . Append this to the objective functional and we get the following expression

$$J(\mathbf{u}^\epsilon) = \int_0^T \left[ \mathbf{g}(\mathbf{x}^\epsilon, \mathbf{u}^\epsilon) + \frac{d}{dt}(\mathbf{z} \cdot \mathbf{x}^\epsilon) \right] dt + \mathbf{z}(0) \cdot \mathbf{x}_0 - \mathbf{z}(T) \cdot \mathbf{x}^\epsilon(T). \quad (16)$$

Since  $\bar{\mathbf{u}}$  is a local minimizer of  $J$ , then derivative of  $J$  over  $\epsilon$  where  $\epsilon = 0$  exists and equals to zero, in other words  $\left. \frac{\partial J(\mathbf{u}^\epsilon)}{\partial \epsilon} \right|_{\epsilon=0} = 0$ . Steering (16) together with simplifying the result by factorization, we get

$$0 = \int_0^T (\mathbf{g}_{\mathbf{x}} + \mathbf{f}_{\mathbf{x}}^\top \mathbf{z} + \dot{\mathbf{z}}) \cdot \frac{\partial \mathbf{x}^\epsilon}{\partial \epsilon} \Big|_{\epsilon=0} dt + \int_0^T (\mathbf{g}_{\mathbf{u}} + \mathbf{f}_{\mathbf{u}}^\top \mathbf{z}) \cdot \kappa dt - \mathbf{z}(T) \cdot \frac{\partial \mathbf{x}^\epsilon}{\partial \epsilon} \Big|_{\epsilon=0}(T). \quad (17)$$

Note from (14)-(15) that variation of  $\kappa$  makes  $\frac{\partial \mathbf{x}^\epsilon}{\partial \epsilon}$  varying. Zeroing the right hand side of (17) together with taking  $\kappa = \mathbf{g}_u + \mathbf{f}_u^\top \mathbf{z}$ , we get the following theorem.

**Theorem 5** *Consider the optimization problem (OC). Let  $\bar{\mathbf{u}} \in \mathcal{U}$  be a local minimizer for  $J$  and  $\bar{\mathbf{x}} \in \mathcal{D}$  be the resulting state. Then there exists a dual variable  $\bar{\mathbf{z}} \in L^2([0, T], \mathbb{R}^5)$  such that the tuple  $(\bar{\mathbf{x}}, \bar{\mathbf{u}}, \bar{\mathbf{z}})$  satisfies the following system*

$$\dot{\mathbf{x}} = \frac{\partial \mathcal{H}}{\partial \mathbf{z}} \text{ with } \mathbf{x}(0) = \mathbf{x}_0 \succeq \mathbf{0}, \quad \dot{\mathbf{z}} = -\frac{\partial \mathcal{H}}{\partial \mathbf{x}}, \quad \frac{\partial \mathcal{H}}{\partial \mathbf{u}} = \mathbf{0}, \quad \mathbf{z}(T) = \mathbf{0} \quad (18)$$

for all  $t \in [0, T]$ . The functional  $\mathcal{H}(\mathbf{x}, \mathbf{u}, \mathbf{z}) := \mathbf{g}(\mathbf{x}, \mathbf{u}) + \langle \mathbf{z}, \mathbf{f}(\mathbf{x}, \mathbf{u}) \rangle$  is Hamiltonian functional, meanwhile all equations in (18) are respectively the state, adjoint, gradient equations and transversality condition.

The adjoint (with transversality condition) and gradient equations can now be unfolded as

$$\dot{\mathbf{z}}_1 = -\omega_{\mathbf{x},1} \mathbf{E}_1 + (\alpha_1 + q\mathbf{u}_1 + \eta_1)\mathbf{z}_1 - \alpha_1 \mathbf{z}_3, \quad \mathbf{z}_1(T) = \mathbf{0} \quad (19a)$$

$$\dot{\mathbf{z}}_2 = -\omega_{\mathbf{x},2} \mathbf{E}_2 + (\alpha_2 + \eta_2)\mathbf{z}_2 - \alpha_2 \mathbf{z}_4, \quad \mathbf{z}_2(T) = \mathbf{0} \quad (19b)$$

$$\dot{\mathbf{z}}_3 = -\omega_{\mathbf{x},3} \mathbf{L}_1 + (2\sigma_1 \mathbf{L}_1 + \beta_1 + \mathbf{u}_1 + \eta_3)\mathbf{z}_3 - \beta_1 \mathbf{z}_5, \quad \mathbf{z}_3(T) = \mathbf{0} \quad (19c)$$

$$\dot{\mathbf{z}}_4 = -\omega_{\mathbf{x},4} \mathbf{L}_2 + (2\sigma_2 \mathbf{L}_2 + \beta_2 + \eta_4)\mathbf{z}_4 - \beta_2 \mathbf{z}_5, \quad \mathbf{z}_4(T) = \mathbf{0} \quad (19d)$$

$$\dot{\mathbf{z}}_5 = -\omega_{\mathbf{x},5} \mathbf{A} + (\mathbf{u}_2 + \eta_5)\mathbf{z}_5 - \mu p \mathbf{z}_1 - \mu(1-p)\mathbf{z}_2, \quad \mathbf{z}_5(T) = \mathbf{0} \quad (19e)$$

and

$$\omega_{\mathbf{u},1} \mathbf{u}_1 - q \mathbf{E}_1 \mathbf{z}_1 - \mathbf{L}_3 \mathbf{z}_3 = 0 \quad (20a)$$

$$\omega_{\mathbf{u},2} \mathbf{u}_2 - \mathbf{A} \mathbf{z}_5 = 0. \quad (20b)$$

It is essential that  $\mathcal{U}$  is bounded as this supports meaningfulness in practical implementation of the control. For the sake of simplicity, we consider  $\mathcal{U} = L^2([0, T], [\mathbf{v}_1, \mathbf{w}_1] \times [\mathbf{v}_2, \mathbf{w}_2])$  and the according projection

$$\text{sat}(\mathbf{u}) = \max(\mathbf{v}, \min(\mathbf{w}, \mathbf{u})) \quad (21)$$

mapping any control  $\mathbf{u}$  into  $\mathcal{U}$ .

To solve the optimal control problem (OC) for bounded control  $\mathbf{u} \in \mathcal{U}$ , we first consider the unconstrained convex optimization problem

$$\text{find } (\mathbf{x}, \mathbf{u}) \in L^2 \times L^2 \text{ such that } J(\mathbf{u}) \rightarrow \min$$

and use the projection (21) to map the control into a convex set  $\mathcal{U}$ . Since  $\mathcal{U}$  is a constriction of  $L^2$ , the projection leads us to the optimal solution of (OC). After all, Algorithm 1 illustrates our scheme to solve (OC).

---

**Algorithm 1** Gradient-based method for solving (OC).

---

**Require:**  $k = 0$ , an initial guess for the control  $\mathbf{u}^k \in \mathcal{U}$ , an error tolerance  $\epsilon > 0$  and an initial step-length  $\lambda > 0$ .

- 1: Compute  $\mathbf{x}_{\mathbf{u}}^k \leftarrow \mathbf{x}^k(\cdot; \mathbf{u}^k)$  and  $\mathbf{z}_{\mathbf{u}}^k \leftarrow \mathbf{z}^k(\cdot; \mathbf{x}^k(\cdot; \mathbf{u}^k))$  consecutively from the state (with forward scheme) and adjoint equation (with backward scheme).
- 2: Compute the objective functional  $J(\mathbf{u}^k)$ .
- 3: Compute  $\bar{\mathbf{u}}^k(\mathbf{x}_{\mathbf{u}}^k, \mathbf{z}_{\mathbf{u}}^k)$  from the gradient equation.
- 4: Update  $\mathbf{u}^{k+1}(\lambda) \leftarrow \mathbf{u}^k + \lambda \bar{\mathbf{u}}^k$  and  $\mathbf{u}^{k+1} \leftarrow \text{sat}(\mathbf{u}^{k+1})$ . Compute  $\mathbf{x}_{\mathbf{u}}^{k+1}$  and  $\mathbf{z}_{\mathbf{u}}^{k+1}$ .
- 5: Compute  $J(\mathbf{u}^{k+1})$  and set  $\Delta J \leftarrow J(\mathbf{u}^{k+1}) - J(\mathbf{u}^k)$ .
- 6: **if**  $|\Delta J| < \epsilon$  **then**
- 7:   Stop. Set  $(\hat{\mathbf{x}}, \hat{\mathbf{c}}, \hat{J}) \leftarrow (\mathbf{x}_{\mathbf{u}}^{k+1}, \mathbf{u}^{k+1}, J(\mathbf{u}^{k+1}))$ .
- 8: **end if**
- 9: **while**  $\Delta J \geq 0$  **do**
- 10:   Update new  $\lambda \leftarrow \arg \min_{s \in [0, \lambda]} \phi(s) := J(\mathbf{u}^{k+1}(s))$  where  $\phi$  is a quadratic representative of  $J$  with respect to the step-length  $s$ . Note that the solution exists since  $\phi(0)$ ,  $\phi'(0)$  and  $\phi(\lambda)$  can be computed directly. Compute new  $\mathbf{u}^{k+1}(\lambda) \leftarrow \mathbf{u}^k + \lambda \bar{\mathbf{u}}^k$  and set  $\mathbf{u}^{k+1} \leftarrow \text{sat}(\mathbf{u}^{k+1})$ . Then compute new  $\mathbf{x}_{\mathbf{u}}^{k+1}$  and  $\mathbf{z}_{\mathbf{u}}^{k+1}$ .
- 11:   Compute  $J(\mathbf{u}^{k+1})$  and set  $\Delta J \leftarrow J(\mathbf{u}^{k+1}) - J(\mathbf{u}^k)$ .
- 12:   **if**  $|\Delta J| < \epsilon$  **then**
- 13:     Stop. Set  $(\hat{\mathbf{x}}, \hat{\mathbf{u}}, \hat{J}) \leftarrow (\mathbf{x}_{\mathbf{u}}^{k+1}, \mathbf{u}^{k+1}, J(\mathbf{u}^{k+1}))$ .
- 14:   **end if**
- 15: **end while**
- 16: Set  $k \leftarrow k + 1$ .
- 17: Go to Step 3.
- 18: **return** The tuple  $(\hat{\mathbf{x}}, \hat{\mathbf{u}}, \hat{J})$ .

---

## 5 Numerical results

Several discussions in this section aims at determining the impact of control intervention to the abundance of mosquito population in visual statements. We accentuate in this paper that significant reduction of the population evolution in forward time be a carry-over from only a cheap optimal control. Realistically, optimal control still suffers from practical drawbacks that one can not easily implement it in real situation. The reason is slightly supported by undisciplined executors in the field to implement the programme as it looks as in optimal control result. On the other hand, a constant control benefits from its easy-to-implement scheme as the executors only need to

deal with fixed-fund allotment problem with flat daily distribution. Another benefit from using constant control is that nominal of the basic mosquito offspring number can explicitly be computed. This computation educes some important statements regarding endemicity of the observed area. It has been preliminarily known that whenever the basic mosquito offspring number is less than unity, then two facts arise: the zero equilibrium is stable but the coexistence equilibrium does not exist. If this number is greater than unity, then the zero equilibrium is unstable and the coexistence equilibrium exists and is stable. This roughly means that the endemic situation will emerge and stay uninterruptible. Otherwise, constant control result always emerges with a higher cost compared to optimal control.

In order to do numerical tests, all the parameters involved in the model have to be represented in numerics. Table 1 gives an estimate for all parameters.

Param.	$M$	$p$	$\mu$	$\eta_1$	$\eta_2$	$\eta_3$	$\eta_4$	$\eta_5$	$q$
Unit	-	-	day <sup>-1</sup>	day <sup>-1</sup>	day <sup>-1</sup>	day <sup>-1</sup>	day <sup>-1</sup>	day <sup>-1</sup>	-
Est. val.	2	0.4	3.1	0.02	0.01	0.002	0.01	0.4	0.04

Param.	$\alpha_1$	$\alpha_2$	$\sigma_1$	$\sigma_2$	$\beta_1$	$\beta_2$
Unit	day <sup>-1</sup>	day <sup>-1</sup>	ind. <sup>-1</sup> × day <sup>-1</sup>	ind. <sup>-1</sup> × day <sup>-1</sup>	day <sup>-1</sup>	day <sup>-1</sup>
Est. val.	0.15	0.13	0.004	$\frac{\sigma_1}{M}$	0.08	0.05

Param.	$T$	$\omega_{\mathbf{x},\{1,\dots,4\}}$	$\omega_{\mathbf{x},5}$	$\omega_{\mathbf{u},\{1,2\}}$	$\mathbf{v}$	$\mathbf{w}$
Unit	day	-	-	-	day <sup>-1</sup>	day <sup>-1</sup>
Est. val.	150	1	2	$4 \times 10^4$	$\mathbf{0}$	[1, 1]

Table 1: Set of all parameters used in the model.

## 5.1 Constant control

To some reasons, the basic mosquito offspring number  $\mathcal{R}(\mathbf{u})$  reflects the hierarchy of endemicity of dengue. The higher  $\mathcal{R}(\mathbf{u})$ , the more rapid mosquito's growth, and therefore the higher the number of dengue incidences. In this case, we demonstrate primary impacts of using constant control to  $\mathcal{R}(\mathbf{u})$ .

Alongside with this illustration, we show a dependence of  $\mathcal{R}(\mathbf{0})$  to some unknown parameters that may be hard to find, or at least hard to estimate. Such parameters can be like  $p$  (the prevalence probability of mosquito adult to breed indoor) and  $\mu$  (the percapita birth rate of eggs). The following figures give the illustrations.




Figure 1: Contour of the basic mosquito offspring number  $\mathcal{R}(\mathbf{u})$  in  $(u_1, u_2)$ -plane. Figure 2: Contour of the basic mosquito offspring number  $\mathcal{R}(\mathbf{0})$  in  $(p, \mu)$ -plane.

From Fig. 1, some conclusions are highlighted. It is clear from the figure that  $u_2$  gives a more significant decrease to nominal of the basic mosquito offspring number compared to  $u_1$ . In constant control case, each value of control taken in the set of real positive numbers can roughly represent a negative return. A negative return  $u_1$  of 1 simply means that, 100% the number of individuals as in the previous step (in day) has to be killed in the current step by Temephos dissemination. In practice, this requirement seems hard to achieve due to several arising technical and spatial heterogeneity problems. From Fig. 1, a suitable choice of constant control pair in the range  $[0, 1] \times [0, 1]$  still gives insignificant reduction to the number of mosquitoes. In an extreme case, killing 100% the number of individuals as in the previous step in both Temephos dissemination and fumigation schemes still arises endemic situation in the observed area. However, to achieve the condition  $\mathcal{R}(\mathbf{u}) < 1$ , one has to produce a high value of negative return. For example, killing 40% (for indoor larvae and  $q$  times indoor eggs) and 180% (for adults) as in the previous step will produce  $\mathcal{R}(\mathbf{u}) \approx 0.9$ . With the same definition, killing respectively 60% and 120% leads to  $\mathcal{R}(\mathbf{u}) \lesssim 1$ , meaning that there

will be a possibility for the mosquito population to be completely eradicated for long time.

Meanwhile from Fig. 2, for whatever the values of  $(p, \mu)$  are taken in the range  $[0, 1] \times [1, 5]$ , the mosquitoes will never die out. The choice of all parameters in the model seems suitable with real situation in endemic region, that is, the basic mosquito offspring number with the absence of control must be greater than unity. This is another reason why we need control intervention. In the figure,  $\mu$  appears with similar sensitivity as  $\mathbf{u}_2$  in Fig. 1. It can be seen that  $\mu$  is very sensitive, as its slight changes lead to the significant differences of the hierarchy of endemicity. In reality,  $\mu$  can be a number that depends on meteorological parameters. It is believed that the more environmental condition sustain mosquitoes' life, the higher  $\mu$ . The result as in Fig. 2 can be another way to confirm that meteorology distribution gives a significant impact to the mosquito abundance in the field. The figure also tells us that, based on our model, to where adult mosquitoes prevalent to breed is not really a big deal.

## 5.2 Time-variant optimal control

Initially, we consider the trajectory of solution to the equation  $\dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{0})$ ,  $\mathbf{x}_0 \succeq \mathbf{0}$ , defined as  $\mathbf{x}(t; \mathbf{0})$ . It can numerically be shown that  $\mathbf{x}(t; \mathbf{0})$  is monotonic and tends to the coexistence equilibrium  $\mathbf{Q}_4$  for very long time. Set  $t_1 = 0$ ,  $t_2 = 50$ ,  $t_3 = 100$  and

$$\mathcal{D}_i := \{\mathbf{x} : \dot{\mathbf{x}} = \mathbf{f}(\mathbf{x}, \mathbf{u}), t \in [0, T], \mathbf{x}_0 = \mathbf{x}(t_i; \mathbf{0}), \mathbf{u} \in \mathcal{U}\}, \quad i = 1, 2, 3. \quad (22)$$

We divide our numerical scheme based on three scenarios:

(Sc-1) find  $(\mathbf{x}, \mathbf{u}) \in \mathcal{D}_1 \times \mathcal{U}$  such that  $J(\mathbf{u}) \rightarrow \min$ ,

(Sc-2) find  $(\mathbf{x}, \mathbf{u}) \in \mathcal{D}_2 \times \mathcal{U}$  such that  $J(\mathbf{u}) \rightarrow \min$ ,

(Sc-3) find  $(\mathbf{x}, \mathbf{u}) \in \mathcal{D}_3 \times \mathcal{U}$  such that  $J(\mathbf{u}) \rightarrow \min$ .

It is supposed that we want to check the dynamics of mosquito population in house-scale including its nearest neighborhood. On average proportion, we let  $\mathbf{x}(t_1; \mathbf{0}) = [8, 8, 6, 6, 5]$ . Thus we obtained from our preliminary simulation that  $\mathbf{x}(t_2; \mathbf{0}) = [200.41, 354.84, 73.82, 134.85, 30.56]$  and  $\mathbf{x}(t_3; \mathbf{0}) = [283.12, 514.21, 91.19, 168.22, 39.16]$  respectively.

One purpose of dividing initial condition for our simulations is to check which scenario arising with the cheapest cost, the one which is applied in  $T$  days after  $t = t_1$  (the earliest growth time),  $t = t_2$  (the peak of outbreak) or  $t = t_3$  (the population almost reach the coexistence equilibrium). It may also



be the case that the cost increases with respect to the magnitude of initial condition. Another purpose is to check which scenario giving the least total endpoints  $\|\mathbf{x}(T; \bar{\mathbf{u}})\|_1$  and if the optimal control results in positive value for  $\|\mathbf{x}(0; \bar{\mathbf{u}})\|_1 - \|\mathbf{x}(T; \bar{\mathbf{u}})\|_1$ , then there is a number  $n > 1$  such that continuation of optimal control scheme will completely eradicate mosquito population in  $nT$  days. Concerning definition of the total cost  $C(\mathbf{u})$ , we define a weighting factor  $A > 0$  such that

$$C(\mathbf{u}) := \frac{A}{2T} \int_0^T \sum_{i=1}^2 \omega_{\mathbf{x},i} \mathbf{u}_i^2 dt. \quad (23)$$

Numerical results from **Sc-1** are given as in Figs. 3-8.




Figure 3: Trajectory of indoor egg population.




Figure 4: Trajectory of outdoor egg population.

At first look, it is clear from Figs. 3-7 that the optimal control makes significant reduction to all classes of mosquito population. With the same scheme, we can find similar results for both **Sc-2** and **Sc-3**. It turns out that by using the data from Table 1, fumigation relatively needs to be enhanced rather than Temephos dissemination during application of the control. Figs. 1 and 8 tell us so. For further managerial reference, Table 2 represents the performance of optimal control in each scenario. We also highlight the optimal control performance when one scheme vanishes.

From the table, we conclude:

1. A complete control intervention using two schemes appears with the highest cost for each of three tested scenarios compared to the ones




Figure 5: Trajectory of indoor larva population.




Figure 6: Trajectory of outdoor larva population.

$C(\mathbf{u})$ , with $A = 1$	<b>Sc-1</b>	<b>Sc-2</b>	<b>Sc-3</b>
$\mathbf{u}_1 \neq 0, \mathbf{u}_2 = 0$	$7.4216 \times 10^2$	$2.9754 \times 10^2$	$2.2304 \times 10^2$
$\mathbf{u}_1 = 0, \mathbf{u}_2 \neq 0$	$6.6658 \times 10^3$	$9.5861 \times 10^3$	$1.0150 \times 10^4$
$\mathbf{u}_1 \neq 0, \mathbf{u}_2 \neq 0$	$1.4004 \times 10^4$	$1.6217 \times 10^4$	$1.6329 \times 10^4$
$\ \mathbf{x}(0; \bar{\mathbf{u}})\ _1, \ \mathbf{x}(T; \bar{\mathbf{u}})\ _1$	<b>Sc-1</b>	<b>Sc-2</b>	<b>Sc-3</b>
$\mathbf{u}_1 \neq 0, \mathbf{u}_2 = 0$	33, 839.57	794.50, 889.13	$1.09 \times 10^3$ , 901.57
$\mathbf{u}_1 = 0, \mathbf{u}_2 \neq 0$	33, 157.66	794.50, 154.73	$1.09 \times 10^3$ , 156.23
$\mathbf{u}_1 \neq 0, \mathbf{u}_2 \neq 0$	33, 31.62	794.50, 35.34	$1.09 \times 10^3$ , 37.52
$\frac{1}{T} \sum_{i=1}^5 \ \mathbf{x}_i(t; \bar{\mathbf{u}})\ _{L^1}$	<b>Sc-1</b>	<b>Sc-2</b>	<b>Sc-3</b>
$\mathbf{u}_1 \neq 0, \mathbf{u}_2 = 0$	610.9374	946.5915	$1.0169 \times 10^3$
$\mathbf{u}_1 = 0, \mathbf{u}_2 \neq 0$	196.5743	225.8798	235.1042
$\mathbf{u}_1 \neq 0, \mathbf{u}_2 \neq 0$	74.7872	119.5978	132.6077

Table 2: The performance of optimal control.

when one scheme vanishes. It is also noted that the cost slightly increases from **Sc-1** to **Sc-3**, meaning there is a positive correlation between initial condition taken for simulation and the cost. If we consider epidemiology of dengue amongst humans and mosquitoes, this fact can help us to decide when we should start to conduct eradication programme. It appears as well in the table that there is a huge difference between the cost for maintaining only Temephos dissemination and only fumigation. Mathematically, this phenomenon arises due to the choice of the trade-off coefficient  $\omega_{\mathbf{x}}$ . In this case, we assume that the




Figure 7: Trajectory of adult popula-  
tion.




Figure 8: Trajectory of optimal con-  
trol.

adults need to be accounted for more attention, hence we put a higher value for  $\omega_{\mathbf{x},5}$ . However, this does not always mean that taking more consideration to reduce adult trajectory requires a higher value of fumigation measure rather than Temephos dissemination measure. In our simulation, even if we take  $\omega_{\mathbf{x},5}$  similar to the others, the result is more or less similar to the one as in Fig. 8. A reader can play around with this coefficient in order to get the result which suits the real situation best.

2. From the table, it is seen that a complete combination of two control measures can extremely reduce the total endpoints of mosquito population. Maintaining only Temephos dissemination leads us to the worst case, i.e. there will be up to 800 total individuals remaining alive after  $T$  days of treatment. Meanwhile, maintaining only fumigation results in moderate total endpoints. If we compute  $\|\mathbf{x}(0; \bar{\mathbf{u}})\|_1 - \|\mathbf{x}(T; \bar{\mathbf{u}})\|_1$ , then we would highly recommend to apply both control schemes as an integrated programme.
3. We interpret  $\frac{1}{T} \sum_{i=1}^5 \|\mathbf{x}_i(t; \bar{\mathbf{u}})\|_{L^1}$  as the average number of total population size at each time during observation. The higher its value, the higher the number of dengue incidences that may probably occur. Again, maintaining only Temephos dissemination leads to the worst case. We highlight that the complete control with two measures gives the best reduction. Now from the table we know that a preventive act of starting the control programme when the number of individuals is

small and a simultaneous combination of both control measures will bring us the best results: least cost, fewest total endpoints and fewest number of individuals living during treatment. However, due to practical limitations, this ideal situation may not be achieved. Knowing the density of mosquito population in average house, we can refer to which column in the last three columns of Table 2 in order to give the best decision for a suitable control management.

## 6 Conclusion

We have an optimal control model for mosquito population dynamics with indoor-outdoor life cycle classification. This model initiates the depicting situation in most dengue endemic areas where mosquitoes originate from indoor as well as outdoor. Two schematic control measures are added to the model as regulators in reducing the evolution of mosquito population in forward time. The first measure represents a negative return that is earned by maintaining Temephos dissemination to some indoor spots where immature phases live. Temephos dissemination aims at killing larvae and a small percentage of eggs. Meanwhile the second measure is represented by a negative return earned from fumigation that targets directly to adults.

Some brief underlying results from the work in this paper are:

In constant control case, the proposed model enumerates a primary biological meaningfulness, i.e. the evolution of mosquito population in forward time results in positive values. With the given estimate of parameters and initial condition, it can numerically be seen that the evolution of total population remains bounded for all  $t > 0$  by the positive number  $\max\{\|\mathbf{x}(0; \mathbf{u})\|_1, \|\mathbf{Q}_4\|_1\}$ .

If the basic mosquito offspring number is less than unity, then two conditions arise: (i) the zero equilibrium is locally asymptotically stable and (ii) the coexistence equilibrium does not exist.

If the basic mosquito offspring number is greater than unity, then three conditions arise: (i) the zero equilibrium is unstable, (ii) the coexistence equilibrium exists and (iii) is globally asymptotically stable in nonnegative orthant.

The optimal control results as displayed in Table 2 can be used as a reference in decision-making process. A brief conclusion states that: in an endemic situation, the best mosquito control impacts are produced if the control is started when the number of individuals is as small as possible, also with combination of the two simultaneous schemes proposed in this paper.

Both in constant control and optimal control cases, one has to enhance

implementation of fumigation rather than Temephos dissemination.

## Acknowledgement

The first and third authors acknowledge the financial support from Indonesia Endowment Fund for Education (LPDP) and Indonesia Directorate of Higher Education (DIKTI) on behalf of ITB Research Grant 2011. The authors are grateful to Prof. Mick Roberts (Massey University, NZ) and Prof. Neville Fowkes (University of Western Australia) and also the handling editor for their constructive comments and recommendations.

## References

- [1] Barrera R. Competition and resistance to starvation in larvae of container-inhabiting *Aedes* mosquitoes. *Ecological Entomology*, **21**: 117–127, 1996.
- [2] Black, W.C., Bennett, K.E., Gorrochoteki-Escalante, N., Barillas-Murry, C.V., Fernandez-Salas, I., de Lourdes Munoz, M., Farfan-Ale, J.A., Olson, K.W., Beaty, B.J. Flavivirus susceptibility in *Aedes aegypti*. *Archive of Medical Research*, **33**(4): 379–388, 2002.
- [3] Braks, M.A.H., Honório, N.A., Lourenço-de-Oliveira, R., Juliano, S.A., Lounibos, L.P. Convergent habitat segregation of *Aedes aegypti* and *Aedes albopictus* (Diptera: Culicidae) in southeastern Brazil and Florida. *Journal of Medical Entomology*, **40**: 785–794, 2003.
- [4] Carbajo, A.E., Curto S.I., Schweigmann, N.J. Spatial distribution pattern of oviposition in the mosquito *Aedes aegypti* in relation to urbanization in Buenos Aires: southern fringe bionomics of an introduced vector. *Medical and Veterinary Entomology*, **20**: 209–218, 2006.
- [5] Chan, K.L., Ho, B.C., Chan, Y.C. *Aedes aegypti* (L.) and *Aedes albopictus* (Skuse) in Singapore City. *Bulletin of World Health Organization*, **44**(5): 629–633, 1971.
- [6] Clemons, A., Haugen, M., Flannery, E., Tomchaney, M., Kast, K., Jacowski, C., Le, C., Mori, A., Holland, W.S., Sarro, J., Severson, D.W., Duman-Scheel, M. *Aedes aegypti*: An emerging model for vector mosquito development. *Cold Spring Harb Protoc*, **2010**(10):1091–1099, 2010.

- [7] Costello, W.G., Taylor, H.M. Mathematical models of the sterile male technique of insect control, in: A. Charnes, W.R. Lynn (eds.). *Mathematical Analysis of Decision Problems in Ecology, Lecture Notes Biomathematics*, Springer-Verlag, Berlin, **5**: 318–359, 1975.
- [8] Driessche, V.d., Watmough, J. Reproduction numbers and sub-threshold endemic equilibria for compartmental models of disease transmission. *Mathematical Biosciences*, **180**: 29–44, 2002.
- [9] Evenhuis, N.L., Gon III, S.M. "22. Family Culicidae". In Neal L. Evenhuis. *Catalog of the Diptera of the Australasian and Oceanian regions. Bishop Museum*: 191–218, 2007.
- [10] Fister, K.R., McCarthy, M.L., Oppenheimer, S.F., Collins, C. Optimal control of insects through sterile insect release and habitat modification. *Mathematical Biosciences*, **244(2)**: 201–212, 2013.
- [11] Gomes, A.d-C., Gotlieb, S.L.D., Marques, C.C.d-A., de Paula, M.B., Marques, G.R.A.M. Duration of larval and pupal development stages of *Aedes albopictus* in natural and artificial containers. *Rev Saúde Pública*, **29**: 15–19, 1995.
- [12] Idowu, O., Adeleke, M., Aina, T.M. Evaluation of indoor breeding activities of mosquitos during the dry season in Abeokuta, Southwestern Nigeria. *Journal of Environmental Health Research*, **12(1)**: 25–29, 2012.
- [13] Judson, C.L., Hokama, Y., Haydockt, I. The psychology of hatching of aedine mosquito eggs: some larval responses to the hatching stimulus. *Journal of Insect Physiology*, **11**: 1169–1177, 1965.
- [14] Lee, D.J., Hiks, M.M., Griffiths, M., Debenham, M.L., Bryan, J.H., Russel, R.C., Geary, M., Marks, E.N. *The Culicidae of the Australasian region*, **4**. Australian Government Publishing Service, Canberra, 1987.
- [15] Luz, P.M., Codeco, C.T., Medlock, J., Struchiner, C.J., Valle, D., Galvani, A.P. Impact of insecticide interventions on the abundance and resistance profile of *Aedes aegypti*. *Epidemiology and Infection*, **137(8)**: 1203–1215, 2012.
- [16] Martins, A.J., Ribeiro, C.D.e., Bellinato, D.F., Peixoto, A.A., Valle, D., Lima, J.B.P. Effect of insecticide resistance on development, longevity and reproduction of field or laboratory selected *Aedes aegypti* populations. *PLoS ONE*, **7(3)**: e31889, 2012.

- [17] Mohammed, A., Chadee, D.D. Effects of different temperature regimens on the development of *Aedes aegypti* (l.) (diptera: Culicidae) mosquitoes. *Acta Tropica*, **119**: 38–43, 2011.
- [18] Plant, R.E., Mangel, M. Modeling and simulation in agricultural pest management. *SIAM Revision*, **29**: 235–261, 1987.
- [19] Powell, J.R., Tabachnick, W.J. History of domestication and spread of *Aedes aegypti* - A Review. *Memórias do Instituto Oswaldo Cruz*, **108(1)**: 11–17, 2013.
- [20] Tabachnick, W.J., Munstermann, L.E., Powell, J.R. Genetic distinctness of sympatric forms of *Aedes aegypti* in East Africa. *Evolution*, **33**: 287–295, 1978.
- [21] Thomé, R.C.A., Yang, H.M., Esteva, L. Optimal control of *Aedes aegypti* mosquitoes by the sterile insect technique and insecticide. *Mathematical Biosciences*, **223**: 12–23, 2010.
- [22] Wijaya, K.P., Götz, T., Soewono, E., Nuraini, N. Temephos spraying and thermal fogging efficacy on *Aedes aegypti* in homogeneous urban residences. *ScienceAsia*, **39S**: 48–56, 2013.
- [23] Zettel, C., Kaufman, P. *Yellow fever mosquito Aedes aegypti*. University of Florida, Institute of Food and Agricultural Sciences, retrieved 27-08-2010.

For more information, please visit the journal website:  
<http://www.worldscinet.com/ijb.html>