

Quasispecies dynamics on a network of interacting genotypes and idiotypes: Formulation of the model

Valmir C. Barbosa,¹ Raul Donangelo,^{2,3} and Sergio R. Souza^{2,4}

¹*Programa de Engenharia de Sistemas e Computação,
COPPE, Universidade Federal do Rio de Janeiro,*

Caixa Postal 68511, 21941-972 Rio de Janeiro - RJ, Brazil

²*Instituto de Física, Universidade Federal do Rio de Janeiro,
Caixa Postal 68528, 21941-972 Rio de Janeiro - RJ, Brazil*

³*Instituto de Física, Facultad de Ingeniería, Universidad de la República,
Julio Herrera y Reissig 565, 11.300 Montevideo, Uruguay*

⁴*Instituto de Física, Universidade Federal do Rio Grande do Sul,
Caixa Postal 15051, 91501-970 Porto Alegre - RS, Brazil*

A quasispecies is the stationary state of a set of interrelated genotypes that evolve according to the usual principles of selection and mutation. Quasispecies studies have invariably concentrated on the possibility of errors during genotype replication and their role in promoting either the survival or the demise of the quasispecies. In a previous work [V. C. Barbosa, R. Donangelo, and S. R. Souza, *J. Theor. Biol.* **312**, 114 (2012)], we introduced a network model of quasispecies dynamics, based on a single probability parameter (p) and capable of addressing several plausibility issues of previous models. Here we extend that model by pairing its network with another one aimed at modeling the dynamics of the immune system when confronted with the quasispecies. The new network is based on the idiotypic-network model of immunity and, together with the previous one, constitutes a network model of interacting genotypes and idiotypes. The resulting model requires further parameters and as a consequence leads to a vast phase space. We have focused on a particular niche in which it is possible to observe the trade-offs involved in the quasispecies' survival or destruction. Within this niche, we give simulation results that highlight some key preconditions for quasispecies survival. These include a minimum initial abundance of genotypes relative to that of the idiotypes and a minimum value of p . The latter, in particular, is to be contrasted with the stand-alone quasispecies network of our previous work, in which arbitrarily low values of p constitute a guarantee of quasispecies survival.

PACS numbers: 87.23.Kg, 89.75.Fb, 02.10.Ox, 02.50.-r

I. INTRODUCTION

The immune system is one of the central regulatory systems of the body, being responsible for the detection and eventual removal of both external agents that may be potentially harmful and body cells whose behavior may have become abnormal. These targets of the immune system are generically referred to as antigens, in allusion to the fact that they trigger the system's response and ultimate production of antibodies or activation of destroying cells. In order to be able to respond appropriately to a potentially large variety of antigens, the immune system is highly adaptive and along its existence evolves from an initial state of innate immunity through a series of states of acquired immunity. Modeling the dynamics that gives rise to this type of learning has been challenging for many decades and has elicited the appearance of at least two main explanatory frameworks. One of them is the clonal-selection theory [1, 2], which postulates the preferential survival of those cell types that are more effective for a certain class of antigens. This theory has been successful in many respects but has failed in others, e.g., explaining the existence of any level of innate immunity, which by definition exists in the absence of any antigens.

The other leading framework to model the dynamics of immunity is that of the so-called idiotypic network

[3]. The main idea is that the molecular structures capable of being recognized by the immune system, known as epitopes, are found not only in antigens but also in the receptors of the immune-system cells whose task is to recognize those antigens. That is, not only can these cells recognize antigens, they can also recognize one another in much the same way. Readily, the existence of such a network of epitope types, known as idiotypes, has the potential to explain not only how immunity is acquired but also how it can exist before antigens are ever encountered. The idiotypic-network theory has enjoyed both enthusiasm and skepticism along the years, the latter owing mainly to the many difficulties associated with confirming it experimentally. Many of its elements, however, are present in models of a hybrid nature, particularly in those that aim to characterize those phenomena, such as autoimmunity, that are essentially of a systemic nature (cf., e.g., [4–6] and references therein).

One common element of all mathematical models of the immune system is the overly simplistic manner in which the system's interaction with antigens is handled. Typically the immune-system model involves a set of time-dependent equations describing the behavior of certain quantities (e.g., the abundance of a given cell type) and including independent terms to account for the presence of antigens. Such terms can be adjusted to account

for the simultaneous presence of several antigens, but in general independently of one another. The drawback, of course, is that in important cases such as infections by some viral species, the many different viral strains that may be concomitantly present mutate into one another frequently and confront the immune system in ever-changing ways.

Here we address the problem of how antigens and entities of the immune system interact with one another when both sides already display complex dynamic behavior even when left to themselves. We introduce a random-graph model of this interaction that takes into account not only the interaction itself but also the dynamics of genotype mutation on the antigen side and of idio- type recognition on the immune-system side. Our model can be viewed as comprising two subnetworks, one whose nodes represent genotypes that mutate into one another as they replicate, the other whose nodes represent idiotypes that stimulate (are recognized by) one another and proliferate as a consequence. The two subnetworks are put together by the addition of new edges to give idiotypes the further stimuli provided by the genotypes.

In building the two subnetworks we have drawn on previous ideas dealing with graph-theoretic representations of both interacting genotypes and idiotypes. On the genotype side the network we use is the same we introduced recently [7] to model the dynamics of the so-called quasispecies. A quasispecies is the stationary state of a set of genotypes that mutate into one another while replicating without recombination based on fitnesses that do not depend on genotype abundance (cf., e.g., [8–12] and references therein). Our model is based on a single probability parameter, p , that regulates both graph connectivity and the occurrence of mutations, and can also successfully account for the well-known transition from adaptation to degeneracy when mutations become too frequent. For the idio- type side we use a similar network while incorporating some of the essential premises of the idio- typic-network theory. Like our quasispecies network, our idio- typic network is based on a single probability parameter, r , whose function is to regulate both graph connectivity and the occurrence of stimulation/recognition of idiotypes by one another.

Each node in our model is a binary sequence, i.e., a sequence of 0's and 1's. This representation is suitable for both genotypes and idiotypes, since it can easily accommodate the types of interaction that are necessary in each subnetwork. Specifically, on the quasispecies side mutations are bit alterations and are more likely to occur between two genotypes that do not differ at too many loci. On the idio- typic side, by contrast, stimulation occurs as a function of how well the two sequences involved can be said to be complementary to each other on a locus-wise basis, being more likely as complementarity increases. This stems directly from the physicochemical nature of the molecular coupling that, in the immune system, leads to stimulation. As the quasispecies and idio-

typic networks are combined together into the full model, this same notion of complementarity is used to account for how genotypes stimulate idiotypes, again based on probability r .

Before proceeding, we note that, by virtue of its underlying random-graph model and of the agent interactions this model represents, the subject we study in the present work is related to various other topics of interest. One of them is the wider discipline of complex networks, which in little more than a decade has uncovered many common underpinnings to a great variety of natural and technological systems of interacting agents [13–15]. Another of these topics is that of evolutionary games, which encompass the so-called predator-prey systems and, essentially, are characterized by the mediation of abundance-dependent fitnesses. The reader is referred to [16, 17], for example, as well as to references therein.

In a similar vein, note that the essence of our model is that it portrays the interaction of antagonistic agent populations as they adapt to each other while competing for supremacy. Thus, while in the present study we instantiate such agents by interrelated genotypes invading the body and idiotypes defending it, taking a broader view may come to benefit several other areas. One area with obvious conceptual ties to our invader-defender setting is that of computer security, and in fact proposals have been put forward that are based on immune-related notions [18, 19]. This has also been the case of other areas with less obvious but equally relevant connections to our study, such as combinatorial optimization (cf. [20] for an application to the prediction of protein structure), user profiling for adaptive information filtering in online systems [21], and classification of textual documents [22]. Useful reviews on how these and other areas have come to be influenced by immune-inspired abstractions can be found in [23, 24].

The remainder of the paper is organized in the following manner. First we introduce the model's details in Section II. Then we discuss some aspects of the computational methodology we have used and present our results, all in Section III. These results elucidate the role played by initial conditions, as well as by one of our main parameters, the probability p , in promoting the survival or the demise of the quasispecies as a function of how genotypes interact with one another and with the idiotypes. These issues are discussed in Section IV, which is followed by conclusions in Section V.

II. MODEL

We consider 2^{L+1} binary sequences of length L and group them into sets A and B , each set comprising all 2^L distinct sequences of length L . Each sequence in set A stands for a genotype, each sequence in set B for an idio- type. Our model is based on a directed random graph D of node set $A \cup B$. For $i \in A \cup B$, we use I_i to denote the set of in-neighbors of node i and O_i to denote its set

of out-neighbors. Graph D has self-loops at all nodes, so it holds that both $i \in I_i$ and $i \in O_i$. We describe the edge set of graph D in three separate stages: first the edges involving nodes in A exclusively, then edges involving nodes in B exclusively, then the edges used to interconnect the two halves.

A. Quasispecies network

For $i, j \in A$ (i.e., i and j are both genotypes), possibly identical, an edge exists directed from i to j with probability $p_{ij} = p^{H_{ij}}$, where p is a probability parameter and H_{ij} is the Hamming distance between i and j . If the edge does exist then it is possible for genotype i to mutate into genotype j during replication, provided $i \neq j$. The probability that such mutation occurs is q_{ij} , assumed proportional to p_{ij} in such a way that $\sum_{j \in O_i} q_{ij} = 1$, where q_{ii} is the probability that genotype i remains unchanged during replication. Note that larger Hamming distances entail smaller connection probabilities and, consequently, smaller mutation probabilities as well.

Letting X_i denote the time-dependent abundance of genotype i , we write

$$\dot{X}_i = \sum_{j \in I_i} f_j q_{ji} X_j, \quad (1)$$

where f_j is the fitness of genotype j and reflects its replication rate. We assume throughout that $f_j = 2^{-d_j}$, where d_j is the number of 1's in genotype j . That is, a genotype's fitness decays exponentially with its Hamming distance to the genotype having no 1's (the fittest genotype, or wild type). Equation (1) is the well-known quasispecies equation [16, 25], now written for graph D as in the uniform case of [7].

B. Idiotypic network

For $i, j \in B$ (i.e., i and j are both idiotypes), again possibly identical, an edge exists directed from i to j with probability $r_{ij} = r^{L-H_{ij}}$, where r is another probability parameter. The existence of this edge indicates that idiomotype i stimulates idiomotype j during the proliferation phase of the idiotypic dynamics, provided $i \neq j$. The probability that this stimulation occurs is s_{ij} , assumed proportional to r_{ij} in such a way that $\sum_{j \in O_i} s_{ij} = 1$, where s_{ii} is the probability that idiomotype i stimulates no other idiomotype during proliferation. Now larger Hamming distances entail larger connection and stimulation probabilities, which indeed are expected to be larger if the idiotypes involved are more complementary to each other (i.e., differ at more loci).

Letting X_i denote the time-dependent abundance of idiomotype i , and assuming that X_i grows in proportion by

some rate $\lambda > 0$ to the total stimulus received by i , yields

$$\dot{X}_i = \lambda \sum_{j \in I_i} s_{ji} X_j. \quad (2)$$

This constitutes a very simple model of the idiotypic network in the absence of antigens.

C. Genotype-idiotype interaction network

We are now in position to describe the entirety of graph D , of node set $A \cup B$, which is intended to model the interaction of the quasispecies and idiotypic networks, of node sets A and B , respectively. We do this by allowing the existence of edges directed from nodes in A to nodes in B , indicating that idiotypes are stimulated not only by one another but also by the genotypes against which they are supposed to constitute a defense. This, in effect, brings antigens into the idiotypic dynamics and lets graph D work as a model of how invading genotypes and defending idiotypes interact with one another. Adding these edges to the network has the potential of enlarging the O_i sets for $i \in A$ and the I_i sets for $i \in B$. The former has the immediate consequence of requiring that we adjust the quasispecies-network constraint $\sum_{j \in O_i} q_{ij} = 1$ to $\sum_{j \in O_i \cap A} q_{ij} = 1$. As for the latter, it remains for us to specify how a genotype $i \in A$ stimulates an idiomotype $j \in B$.

We begin by assuming that the length- L sequences representing genotypes and idiotypes are all relative to the same support, i.e., they all share the same representation space, $\{0, 1\}^L$. In other words, the Hamming distance is well defined also between a genotype and an idiomotype. Our approach is then to handle idiomotype stimulation independently of whether it is effected by a genotype or another idiomotype. That is, we let the edge from genotype $i \in A$ to idiomotype $j \in B$ exist with the same probability r_{ij} as above. When it does exist, stimulation too occurs with the same probability s_{ij} as above, so every node $i \in A$ is now subject to the additional constraint that $\sum_{j \in O_i \cap B} s_{ij} = 1$. Equation (2), therefore, remains unchanged.

Once coupled in this way, the 2^{L+1} differential equations given in Eqs. (1) and (2) mandate an unbounded exponential growth of both genotype and idiomotype abundances from any nontrivial initial values. We therefore lack further coupling in order for the possibility of genotype removal by the idiotypes to be explicitly taken into account. We achieve this by rewriting Eq. (1) as

$$\dot{X}_i = \sum_{j \in I_i} f_j q_{ji} X_j - \mu \sum_{j \in O_i \cap B} s_{ij} X_j, \quad (3)$$

where $\mu > 0$ is a rate parameter. This modification to Eq. (1) lets the abundance of genotype i be decreased at a rate that is proportional to how strongly i stimulates each idiomotype j .

Further modifications to all equations might still be considered for the removal of genotypes as they mutate into other genotypes or the removal of idiotypes as they stimulate (and are therefore recognized and sought for destruction by) other idiotypes. We approach this last step by henceforth considering relative, rather than absolute, abundances. Proceeding in this way is equivalent to imposing that the various abundances sum up to a constant at all times, which automatically leads to the appearance of terms that account for the desired sources of both genotype and idiotypic removal.

For $i \in A \cup B$, let $x_i = X_i / \sum_{k \in A \cup B} X_k$ be the relative abundance of genotype or idiotypic i . Thus, $\sum_{i \in A \cup B} x_i = 1$ at all times. It follows that

$$\begin{aligned} \dot{x}_i &= \frac{\dot{X}_i}{\sum_{k \in A \cup B} X_k} - x_i \frac{\sum_{k \in A \cup B} \dot{X}_k}{\sum_{k \in A \cup B} X_k} \\ &= \frac{\dot{X}_i}{\sum_{k \in A \cup B} X_k} - x_i(\phi - \mu\psi + \lambda), \end{aligned} \quad (4)$$

where $\phi = \sum_{k \in A} f_k x_k$ and $\psi = \sum_{k \in B} x_k \sum_{\ell \in I_k \cap A} s_{\ell k}$. In this expression for ψ , the rightmost summation represents the stimulatory influence of all genotypes upon idiotypic k and is here referred to as that idiotypic's proliferability. Note, additionally, that $\phi / \sum_{k \in A} x_k$ and $\psi / \sum_{k \in B} x_k$ are, respectively, the average genotype fitness and idiotypic proliferability.

Equation (4) yields the final expressions for \dot{x}_i :

$$\dot{x}_i = \sum_{j \in I_i} f_j q_{ji} x_j - \mu \sum_{j \in O_i \cap B} s_{ij} x_j - x_i(\phi - \mu\psi + \lambda) \quad (5)$$

for $i \in A$ and

$$\dot{x}_i = \lambda \sum_{j \in I_i} s_{ji} x_j - x_i(\phi - \mu\psi + \lambda) \quad (6)$$

for $i \in B$.

D. Expected connectivity

In graph D , a genotype i may have out-neighbors both in set A (other genotypes) and in set B (idiotypes). If one ignores all self-loops, then the expected number of out-neighbors of the former type can be obtained by considering every other genotype j and counting the edge from i to j with weight $p^{H_{ij}}$ [26]. That is, a genotype's expected number of out-neighbors that are genotypes other than itself is

$$\sum_{h=1}^L \binom{L}{h} p^h = (1+p)^L - 1. \quad (7)$$

As for a genotype's expected number of out-neighbors that are idiotypes, we have

$$\sum_{h=0}^L \binom{L}{h} r^{L-h} = (1+r)^L. \quad (8)$$

An idiotypic, on the other hand, can only have other idiotypes as out-neighbors. Its number of out-neighbors that are also idiotypes, ignoring self-loops, is

$$\sum_{h=1}^L \binom{L}{h} r^{L-h} = (1+r)^L - r^L. \quad (9)$$

III. RESULTS

We begin by revisiting Eq. (5) and recognizing that it is possible for \dot{x}_i to be negative when $x_i = 0$, thus leading the implicit constraint that $x_i \geq 0$ for all $i \in A \cup B$ at all times to be violated. We fix this by noting that it suffices that this equation's predecessor for absolute abundances, Eq. (3), be modified to

$$\dot{X}_i = \sum_{j \in I_i} f_j q_{ji} X_j - \mu H(X_i) \sum_{j \in O_i \cap B} s_{ij} X_j, \quad (10)$$

where $H(z)$ is the Heaviside step function, adapted to yield 1 if and only if $z > 0$ (yield 0, otherwise). This modification forbids \dot{X}_i to be negative when $X_i = 0$ and affects both Eqs. (5) and (6), since the expression for ψ is itself affected.

The actual expressions we use are then as follows, where all appearances of the H function have been shifted by a suitably small $\delta > 0$ to avoid numerical instabilities. For

$$\psi = \sum_{k \in B} x_k \sum_{\ell \in I_k \cap A} s_{\ell k} H(x_\ell - \delta), \quad (11)$$

we have

$$\dot{x}_i = \sum_{j \in I_i} f_j q_{ji} x_j - \mu H(x_i - \delta) \sum_{j \in O_i \cap B} s_{ij} x_j - x_i(\phi - \mu\psi + \lambda) \quad (12)$$

for $i \in A$ and

$$\dot{x}_i = \lambda \sum_{j \in I_i} s_{ji} x_j - x_i(\phi - \mu\psi + \lambda) \quad (13)$$

for $i \in B$.

While Eqs. (12) and (13) provide instantaneous values of \dot{x}_i that are correct in the sense of never allowing x_i to fall below δ , actually solving the equations requires further control. Specifically, when the value of x_i is to be updated, say at time t , one must ensure that the time step to be used, Δt , necessarily entails $x_i + \dot{x}_i \Delta t \geq \delta$. Thus, before stepping time when $\dot{x}_i < 0$ one must first discover the time $T > t$ at which this derivative will lead to $x_i = \delta$ [that is, one must solve $x_i + \dot{x}_i(T - t) = \delta$ for T] and then choose $\Delta t < T - t$. What we do is solve $x_i + \dot{x}_i(T - t) = \delta/10$ instead (again, to avoid numerical instabilities), which implies that $x_i > \delta/10$ at all times. Consequently, even though Eqs. (12) and (13) ensure $\dot{x}_i \geq 0$ already for $x_i = \delta$, during solution it is still

possible for x_i to fall below δ , though remaining strictly above $\delta/10$. We use $\delta = 10^{-10}$ throughout.

We give results for $L = 10$ (i.e., for 1024 genotypes and 1024 idiotypes) in Figs. 1–6. These results reflect our exploration of a specific parameter niche in which it is possible to observe a rich variety of behaviors for the entire quasispecies (Figs. 1–3) and, particularly, for the wild type (Figs. 4–6). We follow [7] and let x_1 denote the relative abundance of the wild type. Moreover, we let the relative abundance of genotypes be denoted by x_A , i.e.,

$$x_A = \sum_{i \in A} x_i. \quad (14)$$

We use $x_A(0)$ to denote the initial value of x_A .

All figures give results based on at least 10^4 instances of graph D . Solving Eqs. (12) and (13) for each instance starts at uniform relative abundances for genotypes and likewise for idiotypes. That is, initially $x_i = x_A(0)/2^L$ for $i \in A$ and $x_i = [1 - x_A(0)]/2^L$ for $i \in B$. All 2^{L+1} equations are then time-stepped through $t = 50$, which empirically we found to suffice for all relative abundances to reach a stationary state.

IV. DISCUSSION

The three plots in Fig. 1 show the distribution of the genotypes' relative abundance, x_A , as t is varied from $t = 0$ through $t = 30$. All cases correspond to $p = r = 0.1$, meaning that on average both the genotype and the idiotypic subnetworks have densities of the same order of magnitude, and so does the set of edges directed from the genotype subnetwork to the idiotypic subnetwork to account for the removal of genotypes by idiotypes [cf. Eqs. (7)–(9)]. It also means that the dynamics of genotype mutation and of idiotypic stimulation are based on the same underlying probability. The three cases also have in common that $\lambda = \mu = 0.1$, so the rate at which idiotypes proliferate due to stimulation by other idiotypes and the rate at which genotypes are eliminated as they stimulate idiotypes are the same. What distinguishes the three cases from one another is the initial relative abundance of genotypes, $x_A(0)$, which is varied from 0.14 in panel (a) through 0.16 in panel (c).

These three values of $x_A(0)$ were singled out, despite being so close to one another, because they allow us to qualitatively zoom in on what appears to be a transition from a regime in which the genotypes may either disappear or endure in the long run (i.e., the quasispecies may perish or survive) to another in which they endure almost certainly. In fact, examining the three panels of Fig. 1 reveals that, even though in all three cases the genotypes are driven toward a sharp decrease in relative abundance up to about $t = 2$, regardless of the particular instance of graph D on which the dynamics is taking place, thereafter network structure begins to matter and does so in a manner that depends on what the relative genotype abundance was to begin with.




FIG. 1: Evolution of the probability distribution of x_A from $x_A(0) = 0.14$ (a), $x_A(0) = 0.15$ (b), and $x_A(0) = 0.16$ (c), for $p = r = 0.1$ and $\lambda = \mu = 0.1$. Data are based on at least 10^4 independent instances of graph D and are log-binned to the base 1.2. Probabilities are given according to the gray scale on the right of each panel, where darker is closer to 1.

For the lowest of the three values used in the figure [$x_A(0) = 0.14$], and notwithstanding the fact that there exist network topologies continuing to push the genotypes toward the demise of the quasispecies, there are also cases in which the underlying network topology supports the survival of the quasispecies to the point




FIG. 2: Evolution of the probability distribution of x_A from $x_A(0) = 0.10$ for $p = r = 0.1$. Panel (a) has $\lambda = 0.01$ with $\mu = 0.1$; panel (b) has $\lambda = 0.1$ with $\mu = 0.01$. Data are based on at least 10^4 independent instances of graph D and are log-binned to the base 1.2. Probabilities are given according to the gray scale on the right of each panel, where darker is closer to 1.

that it seriously threatens the idiotypic population, i.e., x_A approaches 1. This turn of events becomes significantly more pronounced as $x_A(0)$ is slightly increased, quickly reaching the situation in which almost no network topology supports the destruction of the quasispecies [for $x_A(0) = 0.16$].

Interestingly, a similar situation occurs when $x_A(0)$ remains fixed, along with p and r , while λ and μ are made to vary relative to each other. This is shown in Fig. 2, in whose panels we have $p = r = 0.1$ and $x_A(0) = 0.1$ but λ and μ are given values one order of magnitude apart: $\lambda = 0.01$ and $\mu = 10\lambda$ in panel (a), $\lambda = 0.1$ and $\mu = \lambda/10$ in panel (b). Thus, the ratio λ/μ increases by two orders of magnitude from one scenario to the next. In the first of these scenarios it is uncertain whether the quasispecies will be led to extinction, whereas in the second it almost certainly survives. The ratio λ/μ indicates how much more responsive the immune system is in reorganizing itself to respond to the threat of the genotypes than it is in actually destroying genotypes. What we see in the figure




FIG. 3: (Color online) Probability distribution of a genotype's number of out-neighbors in set A , ignoring self-loops. Data are given for $p = 0.1$ and are based on 10^4 independent instances of graph D . As defined, the probability that an edge exists between two genotypes in D depends on the Hamming distance between them. If this probability were independent of the two genotypes, the resulting distribution would be the Poisson distribution, here shown as a solid line for the same mean of 1.59 that is given for $p = 0.1$ by Eq. (7). If z is the expected number of out-neighbors given by this equation, then the Poisson distribution of mean z requires that edges exist with the fixed probability $z/(2^L - 1) = [(1+p)^L - 1]/(2^L - 1)$, which approaches the probabilities we use only as p nears 1.

is that a heavy imbalance toward the former almost certainly leads to dominance by the attacking quasispecies.

Throughout the transition depicted in Figs. 1 and 2, the bifurcation that the genotype population undergoes at about $t = 2$ clearly depends on the particular instance of graph D being used. We study the role of network topology on the quasispecies by initially continuing to focus on the $p = r = 0.1$ setting that is common to both Figs. 1 and 2. In this setting, the stationary-state distribution of a genotype's number of out-neighbors is the one shown in Fig. 3, where only out-neighbors in set A are counted (i.e., out-neighbors that are genotypes as well, so only the value of p matters) and self-loops are ignored (so that 0 is a possibility). This distribution is clearly concentrated on the lowest numbers of out-neighbors and in this range it seems to be possible to approximate it by the Poisson distribution of mean given by Eq. (7). However, the Poisson distribution arises only when edges exist with the same probability for all node pairs (as in the original Erdős-Rényi model [27] and its directed variant [26]), and in fact we see in the figure that the two distributions differ markedly for the higher numbers of out-neighbors.

The distribution shown in Fig. 3 indicates that a randomly chosen genotype is likely to be able to mutate only into a small number of other genotypes. If idiotypes were altogether absent this would entail excellent survival chances for the quasispecies, meaning in particular




FIG. 4: (Color online) Stationary-state relative abundance of the wild type (x_1) as a function of its number of out-neighbors in set A , ignoring the self-loop. Data are given for $p = r = 0.1$ and $\lambda = \mu = 0.1$, and are based on 10^5 independent instances of graph D .




FIG. 5: (Color online) Stationary-state relative abundance of the wild type (x_1) as a function of its number of out-neighbors in set A , ignoring the self-loop. Data are given for $p = r = 0.1$ and $\lambda = \mu = 0.1$, and are based on either 10^5 (filled symbols) or 10^6 (empty symbols) independent instances of graph D .

that the wild type would be able to climb from its initial relative abundance of $1/2^L \approx 10^{-3}$ toward a stationary-state relative abundance of the order of 10^{-1} [7]. In the presence of idiotypes, however, one must look at how network topology affects the quasispecies from a closer perspective, paying special attention to how the wild type's number of out-neighbors affects its stationary-state relative abundance, x_1 .

Figure 4 shows the average value of x_1 as a function of the wild type's number of out-neighbors in set A and ignoring the self-loop. All data in this figure are still




FIG. 6: (Color online) Stationary-state relative abundance of the wild type (x_1) as a function of its number of out-neighbors in set A , ignoring the self-loop. Data are given for $\lambda = \mu = 0.1$ and $x_A(0) = 0.1$, and are based on 10^5 independent instances of graph D when $p = 0.1$, 10^6 instances when $p = 0.01$.

relative to $p = r = 0.1$ and $\lambda = \mu = 0.1$. Each plot corresponds to a different value of the initial genotype population, beginning at $x_A(0) = 0.095$ and proceeding through $x_A(0) = 0.15$ (i.e., up to inside the transition qualitatively depicted in Fig. 1). In those instances of graph D in which the wild type has no out-neighbors, the value of x_1 is influenced from outside exclusively by mutations into the wild type and by the action of the idiotypes. Under these circumstances, it is clear from the figure that the wild type recovers from its initial situation of uniform dilution with respect to the other genotypes, i.e., from an initial relative abundance of $x_A(0)/2^L \approx 10^{-4}$. Instances with larger numbers of out-neighbors at the wild type, on the other hand, reflect the effect of mutations of the wild type into other genotypes as well, leading to progressively smaller values of x_1 until a precipitous drop to some value between $\delta/10$ and δ is reached (cf. Section III), indicating the total absence of D instances in which the wild type has any of the corresponding numbers out-neighbors. Some of this figure's plots are repeated in Fig. 5 alongside new plots for the same values of $x_A(0)$ but ten times as many instances of graph D , which postpones the aforementioned drop as D instances appear in which the wild type has a larger number of out-neighbors. Taken together, the two figures seem to support the presence of an exponential decay of x_1 with the wild type's number of out-neighbors. However, as already suggested by the data in Fig. 1, such decay is ever slower as $x_A(0)$ increases, so the wild type is ever more resilient to the possibility of mutation into a larger number of genotypes.

Varying the value of p and r relative to each other can provide further insight into wild-type survival, since these are the parameters regulating edge density and also mutation or stimulation on those edges that do exist in a

particular instance of graph D . We give data for this in Fig. 6, which continues to be relative to $\lambda = \mu = 0.1$ but now focuses exclusively on the $x_A(0) = 0.1$ case of Figs. 4 and 5. Note, first, that using $p = 0.1$ with $r = 0.01$ has no relevant effect with respect to the $p = r = 0.1$ case. The reasons for this can be grasped from Eqs. (8) and (9): According to these equations, decreasing r from 0.1 to 0.01, though entailing a difference by one order of magnitude in the value of this probability, impacts the expected number of out-neighbors a genotype or an idio type has inside set B much more modestly. In fact, these variations are in both cases from about 2.59 to about 1.01 and seem to affect genotype-idio type interaction very little.

On the other hand, using $p = 0.01$ with $r = 0.1$, while affecting only the quasispecies side of the network in comparison to the $p = r = 0.1$ case, leads to very different results. By Eq. (7), now a genotype's expected number of out-neighbors in set A drops from 1.59 to 0.1, therefore by one order of magnitude as well. Genotypes are then very sparsely interconnected to one another, but in spite of this we see in Fig. 6 that the wild type manages to survive for those instances of graph D in which it has a very small number of out-neighbors. The pattern of survival is significantly different from the previous one, which may at first seem as a surprise given that the pull of idio types on genotypes continues to be the same and all that has changed is the density in genotype interconnection and the probabilities that mutations occur. However, the resulting expected sparsity does not rule out those relatively rare D instances in which the wild type is fed by mutations from at least one other genotype. Combined with the very low number of genotypes into which the wild type itself may mutate, this unexpected existence of edges incoming to the wild type ensures survival.

It follows from these analyses of the data in Fig. 6 that, as in the case of the isolated quasispecies network [7], probability p is instrumental in determining the fate of the wild type. Contrasting with that case, however, the wild type is no longer guaranteed to survive for arbitrarily low values of p . Instead, it seems that some minimum value is required for the quasispecies to resist the action of the idio types, i.e., for the wild type to escape being diluted into the remainder of the quasispecies.

V. CONCLUSION

Quasispecies studies have invariably concentrated on characterizing two distinct regimes for the evolution of genotypes through time, one in which the wild type survives in the quasispecies, another in which the wild type becomes diluted and no more abundant than any of the quasispecies' other genotypes. Distinguishing between the two regimes has been a matter of selecting the right perspective from which to model replication errors, or mutations. In our own previous model, for example, the single parameter p can be used to characterize quasispecies survival or demise: Increasing p progressively

leads the wild type to a situation in which its relative abundance cannot rise above those of the other, less fit genotypes [7]. This focus on the two extremal regimes of survival and demise has been motivated by the theory's purported use in the modeling of viral populations, along with its appeal as a potential aid in the discovery of therapies [10, 11].

However, in our view the picture has clearly been incomplete, particularly when viral pathogens are the motivation, because the quasispecies' interaction with the host's immune system seems to have been left completely aside. The present work constitutes an attempt to take the immune system into account and study its effect on quasispecies behavior. Our model uses the same network as [7] to represent the interacting genotypes and interconnects it to another network built on top of the so-called idio types, some of the fundamental motifs of the host's immune response. Like the quasispecies network, this new, idio typic network is self-adapting (based on the idea of complementarity among idio types) and potentially effective in destroying genotypes (based on the idea that genotypes and idio types are mutually complementary to some degree).

The resulting network is a continuous-time dynamical system whose variables, representing the relative abundances of genotypes and idio types, are coupled with one another according to a random-graph model. The system's behavior depends on four parameters (the probabilities p and r and the rates λ and μ), as well as on the initial relative abundance of genotypes, $x_A(0)$, which together give rise to a phase space of vast proportions. In view of this, we have concentrated on analyzing a specific niche inside which both survival and destruction of the quasispecies can be examined side by side.

Our exploration of this particular niche has highlighted the existence of two main factors influencing quasispecies survival. The first one is $x_A(0)$ itself, or equivalently the interplay between λ and μ . If genotypes are reasonably abundant in the beginning with respect to idio types, or if idio types eliminate genotypes much more slowly than the idio types adapt their abundances to meet the challenge of the mutating genotypes (i.e., $\mu \ll \lambda$), then it is possible for the quasispecies to survive. The second factor is the topology of the quasispecies network, governed by the p parameter. Unlike the case of the isolated quasispecies network, in which very low values of p are practically a guarantee that the wild type will nearly dominate the quasispecies even from an initial situation of total dilution, the presence of the idio typic network makes things quite different. Specifically, a non-negligible value of p seems necessary to ensure that the quasispecies network is sufficiently dense, and mutations sufficiently likely, for genotypes to change into one another and help the wild type evade the action of the idio typic network.

Further research will concentrate on revisiting Eqs. (2) and (3) with the goal of modeling the action on the immune system of those quasispecies, such as that of HIV, that work toward depleting the system's supply of idio-

types. While the required changes may turn out to be conceptually simple, it seems unavoidable that at least one new rate parameter will be needed, thus enlarging the model's phase space even further. Finding an appropriate parameter niche to explore, however, may provide interesting insight into how genotype mutation, idiotypic stimulation, and idiotypic destruction affect one another.

Acknowledgments

We acknowledge partial support from CNPq, CAPES, a FAPERJ BBP grant, and the joint PRONEX initiative of CNPq/FAPERJ under contract No. 26-111.443/2010.

-
- [1] F. M. Burnet, *The Clonal Selection Theory of Acquired Immunity* (Cambridge University Press, Cambridge, UK, 1959).
 - [2] D. R. Forsdyke, *FASEB J.* **9**, 164 (1995).
 - [3] N. K. Jerne, *Ann. Inst. Pasteur Imm.* **C125**, 373 (1974).
 - [4] L. E. Flores, E. J. Aguilar, V. C. Barbosa, and L. A. V. Carvalho, *J. Theor. Biol.* **229**, 311 (2004).
 - [5] U. Behn, *Immunol. Rev.* **216**, 142 (2007).
 - [6] A. Madi, D. Y. Kenett, S. Bransburg-Zabary, Y. Merbl, F. J. Quintana, A. I. Tauber, I. R. Cohen, and E. Ben-Jacob, *PLoS ONE*, **6**, e17445 (2011).
 - [7] V. C. Barbosa, R. Donangelo, and S. R. Souza, *J. Theor. Biol.* **312**, 114 (2012).
 - [8] M. Eigen, *Naturwissenschaften* **58**, 465 (1971).
 - [9] M. Eigen and P. Schuster, *Naturwissenschaften* **64**, 541 (1977).
 - [10] E. Domingo, *Contrib. Sci.* **5**, 161 (2009).
 - [11] A. S. Luring and R. Andino, *PLoS Pathog.* **6**, e1001005 (2010).
 - [12] A. Más, C. López-Galíndez, I. Cacho, J. Gómez, and M. A. Martínez, *J. Mol. Biol.* **397**, 865 (2010).
 - [13] S. Bornholdt and H. G. Schuster, eds., *Handbook of Graphs and Networks* (Wiley-VCH, Weinheim, Germany, 2003).
 - [14] M. Newman, A.-L. Barabási, and D. J. Watts, eds., *The Structure and Dynamics of Networks* (Princeton University Press, Princeton, NJ, 2006).
 - [15] B. Bollobás, R. Kozma, and D. Miklós, eds., *Handbook of Large-Scale Random Networks* (Springer, Berlin, Germany, 2009).
 - [16] M. A. Nowak, *Evolutionary Dynamics* (Harvard University Press, Cambridge, MA, 2006).
 - [17] J. Knebel, T. Krüger, M. F. Weber, and E. Frey, *Phys. Rev. Lett.* **110**, 168106 (2013).
 - [18] P. K. Harmer, P. D. Williams, G. H. Gunsch, and G. B. Lamont, *IEEE T. Evol. Comput.* **6**, 252 (2002).
 - [19] M. Drozda, S. Schaust, and S. Schildt, *Nat. Comput.* **10**, 243 (2011).
 - [20] V. Cutello, G. Nicosia, M. Pavone, and J. Timmis, *IEEE T. Evol. Comput.* **11**, 101 (2007).
 - [21] N. Nanas, M. Vavalis, and A. De Roeck, *Swarm Intell.* **4**, 275 (2010).
 - [22] A. Abi-Haidar and L. M. Rocha, *Evol. Intell.* **4**, 69 (2011).
 - [23] S. M. Garrett, *Evol. Comput.* **13**, 145 (2005).
 - [24] J. Timmis, A. Hone, T. Stibor, and E. Clark, *Theor. Comput. Sci.* **403**, 11 (2008).
 - [25] C. K. Biebricher and M. Eigen, in *Quasispecies: Concept and Implications for Virology*, edited by E. Domingo (Springer, Berlin, Germany, 2006), vol. 299 of *Current Topics in Microbiology and Immunology*, pp. 1–31.
 - [26] R. M. Karp, *Random Struct. Algor.* **1**, 73 (1990).
 - [27] P. Erdős and A. Rényi, *Publ. Math. (Debrecen)* **6**, 290 (1959).