



COMP220: Graphics & Simulation

# 9: Newtonian mechanics

# Learning outcomes

- ▶ **Recall** the definitions of key concepts such as position, velocity, acceleration, force, friction and restitution
- ▶ **Solve** simple mathematical problems involving these key concepts
- ▶ **Write** programs which feature realistic physics simulations

# Calculus



# Isaac Newton (1643-1727)



# Isaac Newton (1643-1727)

- Invented **calculus**



# Isaac Newton (1643-1727)

- ▶ Invented **calculus**
  - ▶ Study of **rates of change**



# Isaac Newton (1643-1727)

- ▶ Invented **calculus**
  - ▶ Study of **rates of change**
- ▶ Developed **laws of motion**



# Isaac Newton (1643-1727)

- ▶ Invented **calculus**
  - ▶ Study of **rates of change**
- ▶ Developed **laws of motion**
  - ▶ “The” laws of motion until 20th Century (Einstein’s theory of relativity, quantum mechanics)





# Isaac Newton (1643-1727)

- ▶ Invented **calculus**
  - ▶ Study of **rates of change**
- ▶ Developed **laws of motion**
  - ▶ “The” laws of motion until 20th Century (Einstein’s theory of relativity, quantum mechanics)
  - ▶ Still useful for motion of “everyday” objects (size above quantum scale, speed much lower than speed of light)



# Isaac Newton (1643-1727)



- ▶ Invented **calculus**
  - ▶ Study of **rates of change**
- ▶ Developed **laws of motion**
  - ▶ “The” laws of motion until 20th Century (Einstein’s theory of relativity, quantum mechanics)
  - ▶ Still useful for motion of “everyday” objects (size above quantum scale, speed much lower than speed of light)
- ▶ Developed **laws of gravitation**

# Isaac Newton (1643-1727)



- ▶ Invented **calculus**
  - ▶ Study of **rates of change**
- ▶ Developed **laws of motion**
  - ▶ “The” laws of motion until 20th Century (Einstein’s theory of relativity, quantum mechanics)
  - ▶ Still useful for motion of “everyday” objects (size above quantum scale, speed much lower than speed of light)
- ▶ Developed **laws of gravitation**
  - ▶ Realised that falling objects and orbiting celestial bodies are governed by the same principles

# Isaac Newton (1643-1727)



- ▶ Invented **calculus**
  - ▶ Study of **rates of change**
- ▶ Developed **laws of motion**
  - ▶ “The” laws of motion until 20th Century (Einstein’s theory of relativity, quantum mechanics)
  - ▶ Still useful for motion of “everyday” objects (size above quantum scale, speed much lower than speed of light)
- ▶ Developed **laws of gravitation**
  - ▶ Realised that falling objects and orbiting celestial bodies are governed by the same principles
- ▶ Many other contributions to mathematics and physics

# Rates of change

# Rates of change


- ▶ Consider a quantity that **changes over time**

# Rates of change

- ▶ Consider a quantity that **changes over time**
- ▶ Rate of change =  $\frac{\text{change in quantity}}{\text{change in time}}$

# Rates of change

- ▶ Consider a quantity that **changes over time**
- ▶ Rate of change =  $\frac{\text{change in quantity}}{\text{change in time}}$




Same as the **gradient** of a graph  
(from GCSE maths):

$$\text{gradient} = \frac{\text{change in } y}{\text{change in } x}$$



# Rates of change

- ▶ Consider a quantity that **changes over time**
- ▶ Rate of change =  $\frac{\text{change in quantity}}{\text{change in time}}$




Same as the **gradient** of a graph  
(from GCSE maths):

$$\text{gradient} = \frac{\text{change in } y}{\text{change in } x}$$

- ▶ The **derivative** of a quantity  $x$  with respect to time  $t$  is **the rate of change** of  $x$  with respect to  $t$

# Rates of change

- ▶ Consider a quantity that **changes over time**
- ▶ Rate of change =  $\frac{\text{change in quantity}}{\text{change in time}}$




Same as the **gradient** of a graph  
(from GCSE maths):

$$\text{gradient} = \frac{\text{change in } y}{\text{change in } x}$$

- ▶ The **derivative** of a quantity  $x$  with respect to time  $t$  is **the rate of change** of  $x$  with respect to  $t$
- ▶ Denoted  $\frac{dx}{dt}$

# Rates of change

- ▶ Consider a quantity that **changes over time**
- ▶ Rate of change =  $\frac{\text{change in quantity}}{\text{change in time}}$



Same as the **gradient** of a graph  
(from GCSE maths):

$$\text{gradient} = \frac{\text{change in } y}{\text{change in } x}$$

- ▶ The **derivative** of a quantity  $x$  with respect to time  $t$  is **the rate of change** of  $x$  with respect to  $t$
- ▶ Denoted  $\frac{dx}{dt}$
- ▶ The mathematical process of finding  $\frac{dx}{dt}$  given  $x$  is called **differentiation**

# Derivatives – example

# Derivatives – example

- ▶ A car is driving along a straight road at a constant speed

# Derivatives – example

- ▶ A car is driving along a straight road at a constant speed
- ▶ In half an hour, it covers a distance of 20 miles

# Derivatives – example

- ▶ A car is driving along a straight road at a constant speed
- ▶ In half an hour, it covers a distance of 20 miles
- ▶ Its average speed is  $\frac{20 \text{ miles}}{0.5 \text{ hours}} = 40 \text{ miles per hour}$

# Derivatives – example

- ▶ A car is driving along a straight road at a constant speed
- ▶ In half an hour, it covers a distance of 20 miles
- ▶ Its average speed is  $\frac{20 \text{ miles}}{0.5 \text{ hours}} = 40 \text{ miles per hour}$
- ▶ In other words...



# Derivatives – example

- ▶ A car is driving along a straight road at a constant speed
- ▶ In half an hour, it covers a distance of 20 miles
- ▶ Its average speed is  $\frac{20 \text{ miles}}{0.5 \text{ hours}} = 40 \text{ miles per hour}$
- ▶ In other words...
  - ▶ **Distance travelled** is a quantity varying with time

# Derivatives – example

- ▶ A car is driving along a straight road at a constant speed
- ▶ In half an hour, it covers a distance of 20 miles
- ▶ Its average speed is  $\frac{20 \text{ miles}}{0.5 \text{ hours}} = 40 \text{ miles per hour}$
- ▶ In other words...
  - ▶ **Distance travelled** is a quantity varying with time
  - ▶ We call the rate of change of this quantity **speed**

# Derivatives – example

- ▶ A car is driving along a straight road at a constant speed
- ▶ In half an hour, it covers a distance of 20 miles
- ▶ Its average speed is  $\frac{20 \text{ miles}}{0.5 \text{ hours}} = 40 \text{ miles per hour}$
- ▶ In other words...
  - ▶ **Distance travelled** is a quantity varying with time
  - ▶ We call the rate of change of this quantity **speed**
  - ▶ If  $x$  is distance travelled and  $t$  is time, then we have

$$\frac{dx}{dt} = \frac{20}{0.5} = 40$$

# Integration

# Integration

- ▶ Given  $\frac{dx}{dt}$ , find  $x$

# Integration

- ▶ Given  $\frac{dx}{dt}$ , find  $x$
- ▶  $x$  is the **integral** of  $\frac{dx}{dt}$

# Integration

- ▶ Given  $\frac{dx}{dt}$ , find  $x$
- ▶  $x$  is the **integral** of  $\frac{dx}{dt}$
- ▶ The process of finding  $x$  is called **integration**, the opposite of differentiation

# Integration

- ▶ Given  $\frac{dx}{dt}$ , find  $x$
- ▶  $x$  is the **integral** of  $\frac{dx}{dt}$
- ▶ The process of finding  $x$  is called **integration**, the opposite of differentiation
- ▶ We are interested in **numerical integration**



# Integration

- ▶ Given  $\frac{dx}{dt}$ , find  $x$
- ▶  $x$  is the **integral** of  $\frac{dx}{dt}$
- ▶ The process of finding  $x$  is called **integration**, the opposite of differentiation
- ▶ We are interested in **numerical integration**
  - ▶ I.e. integration by computer calculation, not by mathematician with pen and paper...

# Euler method

# Euler method

- ▶ If we know values of  $x$  and  $\frac{dx}{dt}$  at time  $t$ , we can **estimate** the value of  $x$  at time  $t + h$

# Euler method

- ▶ If we know values of  $x$  and  $\frac{dx}{dt}$  at time  $t$ , we can **estimate** the value of  $x$  at time  $t + h$
- ▶ Formula:

$$x(t + h) \approx x(t) + h \times \frac{dx}{dt}(t)$$

# Euler method

- ▶ If we know values of  $x$  and  $\frac{dx}{dt}$  at time  $t$ , we can **estimate** the value of  $x$  at time  $t + h$

- ▶ Formula:

$$x(t + h) \approx x(t) + h \times \frac{dx}{dt}(t)$$

- ▶  $\frac{dx}{dt}$  is rate of change, i.e. how much  $x$  changes by if  $t$  changes by 1

# Euler method


- ▶ If we know values of  $x$  and  $\frac{dx}{dt}$  at time  $t$ , we can **estimate** the value of  $x$  at time  $t + h$

- ▶ Formula:


$$x(t + h) \approx x(t) + h \times \frac{dx}{dt}(t)$$

- ▶  $\frac{dx}{dt}$  is rate of change, i.e. how much  $x$  changes by if  $t$  changes by 1
- ▶ So  $h \times \frac{dx}{dt}$  is how much  $x$  changes by if  $t$  changes by  $h$

# Euler method




# Euler method



- If  $\frac{dx}{dt}$  does not change between  $t$  and  $t + h$ , this gives the **exact** answer; otherwise there will be an **error**




# Euler method



- ▶ If  $\frac{dx}{dt}$  does not change between  $t$  and  $t + h$ , this gives the **exact** answer; otherwise there will be an **error**
- ▶ If  $h$  is small enough, the error should also be small...

# Euler method



- ▶ If  $\frac{dx}{dt}$  does not change between  $t$  and  $t + h$ , this gives the **exact** answer; otherwise there will be an **error**
- ▶ If  $h$  is small enough, the error should also be small...
- ▶ There are more advanced forms of numerical integration which give smaller errors

# Calculus with vectors

# Calculus with vectors

- Can talk about rate of change of vectors as well

# Calculus with vectors

- ▶ Can talk about rate of change of vectors as well
- ▶ If  $x$  is an  $n$ -vector, then so is  $\frac{dx}{dt}$

# Calculus with vectors

- ▶ Can talk about rate of change of vectors as well
- ▶ If  $x$  is an  $n$ -vector, then so is  $\frac{dx}{dt}$
- ▶ Each component of  $\frac{dx}{dt}$  is the rate of change of the corresponding component of  $x$

# Basic mechanics



# Point masses



# Point masses

- For now we assume everything is a **point mass**, i.e. ignore the actual shape and size of objects

# Point masses

- ▶ For now we assume everything is a **point mass**, i.e. ignore the actual shape and size of objects
- ▶ **Mass** is measured in **kilograms**

# Point masses

- ▶ For now we assume everything is a **point mass**, i.e. ignore the actual shape and size of objects
- ▶ **Mass** is measured in **kilograms**
- ▶ Not to be confused with **weight** (GCSE physics!)

# Position, velocity and acceleration

# Position, velocity and acceleration

- ▶ **Position** describes an object's location in space

# Position, velocity and acceleration

- ▶ **Position** describes an object's location in space
  - ▶ Usually expressed as a 3-vector relative to the **origin**

# Position, velocity and acceleration

- ▶ **Position** describes an object's location in space
  - ▶ Usually expressed as a 3-vector relative to the **origin**
  - ▶ Measured in **metres**

# Position, velocity and acceleration

- ▶ **Position** describes an object's location in space
  - ▶ Usually expressed as a 3-vector relative to the **origin**
  - ▶ Measured in **metres**
- ▶ **Velocity** is **rate of change of position**



# Position, velocity and acceleration

- ▶ **Position** describes an object's location in space
  - ▶ Usually expressed as a 3-vector relative to the **origin**
  - ▶ Measured in **metres**
- ▶ **Velocity** is **rate of change of position**
  - ▶ Measured in **metres per second** ( $\text{ms}^{-1}$ )

# Position, velocity and acceleration

- ▶ **Position** describes an object's location in space
  - ▶ Usually expressed as a 3-vector relative to the **origin**
  - ▶ Measured in **metres**
- ▶ **Velocity** is **rate of change of position**
  - ▶ Measured in **metres per second** ( $\text{ms}^{-1}$ )
  - ▶ Velocity is a **vector**

# Position, velocity and acceleration

- ▶ **Position** describes an object's location in space
  - ▶ Usually expressed as a 3-vector relative to the **origin**
  - ▶ Measured in **metres**
- ▶ **Velocity** is **rate of change of position**
  - ▶ Measured in **metres per second** ( $\text{ms}^{-1}$ )
  - ▶ Velocity is a **vector**
  - ▶ Speed is a **scalar** (a number), the magnitude of velocity

# Position, velocity and acceleration

- ▶ **Position** describes an object's location in space
  - ▶ Usually expressed as a 3-vector relative to the **origin**
  - ▶ Measured in **metres**
- ▶ **Velocity** is **rate of change of position**
  - ▶ Measured in **metres per second** ( $\text{ms}^{-1}$ )
  - ▶ Velocity is a **vector**
  - ▶ Speed is a **scalar** (a number), the magnitude of velocity
  - ▶ speed : velocity :: distance : position

# Position, velocity and acceleration

- ▶ **Position** describes an object's location in space
  - ▶ Usually expressed as a 3-vector relative to the **origin**
  - ▶ Measured in **metres**
- ▶ **Velocity** is **rate of change of position**
  - ▶ Measured in **metres per second** ( $\text{ms}^{-1}$ )
  - ▶ Velocity is a **vector**
  - ▶ Speed is a **scalar** (a number), the magnitude of velocity
  - ▶ speed : velocity :: distance : position
- ▶ **Acceleration** is **rate of change of velocity**

# Position, velocity and acceleration

- ▶ **Position** describes an object's location in space
  - ▶ Usually expressed as a 3-vector relative to the **origin**
  - ▶ Measured in **metres**
- ▶ **Velocity** is **rate of change of position**
  - ▶ Measured in **metres per second** ( $\text{ms}^{-1}$ )
  - ▶ Velocity is a **vector**
  - ▶ Speed is a **scalar** (a number), the magnitude of velocity
  - ▶ speed : velocity :: distance : position
- ▶ **Acceleration** is **rate of change of velocity**
  - ▶ Measured in **metres per second per second** ( $\text{ms}^{-2}$ )

# Newton's Laws of Motion

# Newton's Laws of Motion

An object remains at rest or moves at constant velocity  
unless acted upon by an external force



# Newton's Laws of Motion

An object remains at rest or moves at constant velocity unless acted upon by an external force

$F = ma$ : The sum of forces acting upon an object is equal to its mass multiplied by its acceleration

# Newton's Laws of Motion

An object remains at rest or moves at constant velocity unless acted upon by an external force

$F = ma$ : The sum of forces acting upon an object is equal to its mass multiplied by its acceleration

When one body exerts a force on another, the second body exerts an equal and opposite force on the first

# Force

# Force

- ▶ Measured in **Newtons** (N)

# Force

- ▶ Measured in **Newtons** (N)
- ▶  $F = ma$ : 1N of force causes a 1kg object to accelerate by  $1\text{ms}^{-2}$

# Force

- ▶ Measured in **Newton**s (N)
- ▶  $F = ma$ : 1N of force causes a 1kg object to accelerate by  $1\text{ms}^{-2}$
- ▶ Forces occur when objects **interact**

# Force

- ▶ Measured in **Newtons** (N)
- ▶  $F = ma$ : 1N of force causes a 1kg object to accelerate by  $1\text{ms}^{-2}$
- ▶ Forces occur when objects **interact**
- ▶ E.g. gravity, air resistance, friction

# Force

- ▶ Measured in **Newtons** (N)
- ▶  $F = ma$ : 1N of force causes a 1kg object to accelerate by  $1\text{ms}^{-2}$
- ▶ Forces occur when objects **interact**
- ▶ E.g. gravity, air resistance, friction
- ▶ E.g. reaction force: stops objects from passing through each other



# Force

- ▶ Measured in **Newtons** (N)
- ▶  $F = ma$ : 1N of force causes a 1kg object to accelerate by  $1\text{ms}^{-2}$
- ▶ Forces occur when objects **interact**
- ▶ E.g. gravity, air resistance, friction
- ▶ E.g. reaction force: stops objects from passing through each other
- ▶ E.g. applied forces: car engine, rocket engine, launched projectile, human muscle, ...

# Simulating Newtonian physics

# Simulating Newtonian physics

- ▶ Each object needs to store its **position** and **velocity**

# Simulating Newtonian physics

- ▶ Each object needs to store its **position** and **velocity**
- ▶ On each frame:

# Simulating Newtonian physics

- ▶ Each object needs to store its **position** and **velocity**
- ▶ On each frame:
  - ▶ Apply **numerical integration** to determine the new position from the current velocity

# Simulating Newtonian physics

- ▶ Each object needs to store its **position** and **velocity**
- ▶ On each frame:
  - ▶ Apply **numerical integration** to determine the new position from the current velocity
  - ▶ Calculate the **forces** acting upon the object and use these to calculate **acceleration**

$$F = ma \quad \implies \quad a = \frac{F}{m}$$

# Simulating Newtonian physics

- ▶ Each object needs to store its **position** and **velocity**
- ▶ On each frame:
  - ▶ Apply **numerical integration** to determine the new position from the current velocity
  - ▶ Calculate the **forces** acting upon the object and use these to calculate **acceleration**

$$F = ma \quad \implies \quad a = \frac{F}{m}$$

- ▶ Apply **numerical integration** again to determine the new velocity from the current acceleration

# Gravity



# Gravity

- ▶ Gravity pulls **all objects** with mass **towards each other**

# Gravity

- ▶ Gravity pulls **all objects** with mass **towards each other**
- ▶ Gravitational force is tiny unless one or both objects has a huge mass (e.g. a planet...)

# Gravity

- ▶ Gravity pulls **all objects** with mass **towards each other**
- ▶ Gravitational force is tiny unless one or both objects has a huge mass (e.g. a planet...)
- ▶ Near the surface of a planet, gravity pulls objects **downwards** (i.e. towards the centre of the planet) with a force called **weight**

# Gravity

- ▶ Gravity pulls **all objects** with mass **towards each other**
- ▶ Gravitational force is tiny unless one or both objects has a huge mass (e.g. a planet...)
- ▶ Near the surface of a planet, gravity pulls objects **downwards** (i.e. towards the centre of the planet) with a force called **weight**
- ▶  $w = mg$ , where  $w$  is weight,  $m$  is mass and  $g$  is the **gravitational constant**

# Gravity

- ▶ Gravity pulls **all objects** with mass **towards each other**
- ▶ Gravitational force is tiny unless one or both objects has a huge mass (e.g. a planet...)
- ▶ Near the surface of a planet, gravity pulls objects **downwards** (i.e. towards the centre of the planet) with a force called **weight**
- ▶  $w = mg$ , where  $w$  is weight,  $m$  is mass and  $g$  is the **gravitational constant**
- ▶ On Earth,  $g \approx 9.81$  (often rounded to  $g = 10$ )

# Gravity

$$F = ma$$

$$F = w = mg$$

$$\Rightarrow mg = ma$$

$$\Rightarrow g = a$$

# Gravity

$$F = ma$$

$$F = w = mg$$

$$\implies mg = ma$$

$$\implies g = a$$

- So gravity applies **the same** acceleration ( $9.81 \text{ ms}^{-2}$  downwards) to all objects **regardless** of weight!

# Gravity

$$F = ma$$

$$F = w = mg$$


$$\implies mg = ma$$

$$\implies g = a$$


- ▶ So gravity applies **the same** acceleration ( $9.81 \text{ ms}^{-2}$  downwards) to all objects **regardless** of weight!
- ▶ Famous experiment: in a **vacuum** (no air resistance), a bowling ball falls at the **same speed** as a feather



# Basic collision response




# Basic collision response




- For an **elastic collision**, the component of velocity parallel to the **surface normal** is **reversed**

# Basic collision response




- ▶ For an **elastic collision**, the component of velocity parallel to the **surface normal** is **reversed**
- ▶ E.g. if the surface is the  $xz$  plane, flip the  $y$  component

# Basic collision response



- ▶ For an **elastic collision**, the component of velocity parallel to the **surface normal** is **reversed**
- ▶ E.g. if the surface is the  $xz$  plane, flip the  $y$  component
- ▶ For an **inelastic collision**, some velocity is lost

# Basic collision response



- ▶ For an **elastic collision**, the component of velocity parallel to the **surface normal** is **reversed**
- ▶ E.g. if the surface is the  $xz$  plane, flip the  $y$  component
- ▶ For an **inelastic collision**, some velocity is lost
- ▶ Flip the  $y$  component and multiply it by something between 0 and 1

# Sprint review

