



FALMOUTH
UNIVERSITY



COMP110: Principles of Computing

11: Numerical Methods

Worksheets

- ▶ Worksheet 7: due **today**
- ▶ Worksheet 8: due **next Monday**

Scientific notation



Integer powers

Let $a \neq 0$ be a real number, and let $b > 0$ be an integer

Integer powers

Let $a \neq 0$ be a real number, and let $b > 0$ be an integer

$$a^b = \underbrace{a \times a \times \cdots \times a}_{b \text{ times}}$$

Integer powers

Let $a \neq 0$ be a real number, and let $b > 0$ be an integer

$$a^b = \underbrace{a \times a \times \cdots \times a}_{b \text{ times}}$$

$$a^0 = 1$$

Integer powers

Let $a \neq 0$ be a real number, and let $b > 0$ be an integer

$$a^b = \underbrace{a \times a \times \cdots \times a}_{b \text{ times}}$$

$$a^0 = 1$$

$$a^{-b} = \frac{1}{\underbrace{a \times a \times \cdots \times a}_{b \text{ times}}}$$

Powers of 10

Powers of 10

$$10^6 = 1 \underbrace{000\,000}_{6 \text{ zeroes}}$$

Powers of 10

$$10^6 = 1 \underbrace{000\,000}_{6 \text{ zeroes}}$$

$$10^1 = 10$$

Powers of 10

$$10^6 = 1 \underbrace{000\,000}_{6 \text{ zeroes}}$$

$$10^1 = 10$$

$$10^0 = 1$$

Powers of 10

$$10^6 = 1 \underbrace{000\,000}_{6 \text{ zeroes}}$$

$$10^1 = 10$$

$$10^0 = 1$$

$$10^{-1} = 0.1$$

Powers of 10

$$10^6 = 1 \underbrace{000\,000}_{6 \text{ zeroes}}$$

$$10^1 = 10$$

$$10^0 = 1$$

$$10^{-1} = 0.1$$

$$10^{-6} = 0.\underbrace{000\,001}_{5 \text{ zeroes}}$$

Multiplying by powers of 10

Multiplying by powers of 10

Multiplying by 10^n is the same as moving the decimal point n places to the **right** (adding zeroes if necessary)

Multiplying by powers of 10

Multiplying by 10^n is the same as moving the decimal point n places to the **right** (adding zeroes if necessary)

$$3.14159 \times 10^2 = 314.159$$

Multiplying by powers of 10

Multiplying by 10^n is the same as moving the decimal point n places to the **right** (adding zeroes if necessary)

$$3.14159 \times 10^2 = 314.159$$

$$27 \times 10^3 = 27\,000$$

Multiplying by powers of 10

Multiplying by 10^n is the same as moving the decimal point n places to the **right** (adding zeroes if necessary)

$$3.14159 \times 10^2 = 314.159$$

$$27 \times 10^3 = 27\,000$$

Similarly if n is negative, the decimal point moves to the **left**

Multiplying by powers of 10

Multiplying by 10^n is the same as moving the decimal point n places to the **right** (adding zeroes if necessary)

$$3.14159 \times 10^2 = 314.159$$

$$27 \times 10^3 = 27\,000$$

Similarly if n is negative, the decimal point moves to the **left**

$$123.45 \times 10^{-2} = 1.2345$$

Scientific notation

Scientific notation

- ▶ A way of writing **very large** and **very small** numbers

Scientific notation

- ▶ A way of writing **very large** and **very small** numbers
- ▶ $a \times 10^b$, where

Scientific notation

- ▶ A way of writing **very large** and **very small** numbers
- ▶ $a \times 10^b$, where
 - ▶ a ($1 \leq |a| < 10$) is the **mantissa**

Scientific notation

- ▶ A way of writing **very large** and **very small** numbers
- ▶ $a \times 10^b$, where
 - ▶ a ($1 \leq |a| < 10$) is the **mantissa**
 - ▶ (a is a positive or negative number with a single non-zero digit before the decimal point)

Scientific notation

- ▶ A way of writing **very large** and **very small** numbers
- ▶ $a \times 10^b$, where
 - ▶ a ($1 \leq |a| < 10$) is the **mantissa**
 - ▶ (a is a positive or negative number with a single non-zero digit before the decimal point)
 - ▶ b (an integer) is the **exponent**

Scientific notation

- ▶ A way of writing **very large** and **very small** numbers
- ▶ $a \times 10^b$, where
 - ▶ a ($1 \leq |a| < 10$) is the **mantissa**
 - ▶ (a is a positive or negative number with a single non-zero digit before the decimal point)
 - ▶ b (an integer) is the **exponent**
- ▶ E.g. 1 light year = 9.461×10^{15} metres

Scientific notation

- ▶ A way of writing **very large** and **very small** numbers
- ▶ $a \times 10^b$, where
 - ▶ a ($1 \leq |a| < 10$) is the **mantissa**
 - ▶ (a is a positive or negative number with a single non-zero digit before the decimal point)
 - ▶ b (an integer) is the **exponent**
- ▶ E.g. 1 light year = 9.461×10^{15} metres
- ▶ E.g. Planck's constant = 6.626×10^{-34} joules

Scientific notation

- ▶ A way of writing **very large** and **very small** numbers
- ▶ $a \times 10^b$, where
 - ▶ a ($1 \leq |a| < 10$) is the **mantissa**
 - ▶ (a is a positive or negative number with a single non-zero digit before the decimal point)
 - ▶ b (an integer) is the **exponent**
- ▶ E.g. 1 light year = 9.461×10^{15} metres
- ▶ E.g. Planck's constant = 6.626×10^{-34} joules
- ▶ Socrative FALCOMPED

Scientific notation in code

Scientific notation in code

Instead of writing $\times 10$, write e (no spaces)

Scientific notation in code

Instead of writing $\times 10$, write `e` (no spaces)

```
double lightYear = 9.461e15;  
double plancksConstant = 6.626e-34;
```

Floating point numbers



Floating point numbers

Floating point numbers

- ▶ Similar to scientific notation, but **base 2** (binary)

Floating point numbers

- ▶ Similar to scientific notation, but **base 2** (binary)
- ▶ $a \times 10^b$, where

Floating point numbers

- ▶ Similar to scientific notation, but **base 2** (binary)
- ▶ $a \times 10^b$, where
 - ▶ a ($1 \leq |a| < 2$) is the **mantissa**

Floating point numbers

- ▶ Similar to scientific notation, but **base 2** (binary)
- ▶ $a \times 10^b$, where
 - ▶ a ($1 \leq |a| < 2$) is the **mantissa**
 - ▶ (a is a positive or negative number with a single 1 before the “decimal point”)

Floating point numbers

- ▶ Similar to scientific notation, but **base 2** (binary)
- ▶ $a \times 10^b$, where
 - ▶ a ($1 \leq |a| < 2$) is the **mantissa**
 - ▶ (a is a positive or negative number with a single 1 before the “decimal point”)
 - ▶ b (an integer) is the **exponent**

Storing the mantissa

Storing the mantissa

- ▶ A positive or negative number with a single 1 before the “decimal point”

Storing the mantissa

- ▶ A positive or negative number with a single 1 before the “decimal point”
- ▶ Sign is stored as a single bit: 0 = +, 1 = -

Storing the mantissa

- ▶ A positive or negative number with a single 1 before the “decimal point”
- ▶ Sign is stored as a single bit: 0 = +, 1 = -
- ▶ We know there’s a 1 before the point so no need to store it — just store the binary digits after the point

Storing the exponent

Storing the exponent

- ▶ An integer — can be positive, negative or zero

Storing the exponent

- ▶ An integer — can be positive, negative or zero
- ▶ NB we don't use 2's complement to store it!

Storing the exponent

- ▶ An integer — can be positive, negative or zero
- ▶ NB we don't use 2's complement to store it!
- ▶ Instead it is stored in binary as a positive integer with a **bias** added on

Storing the exponent

- ▶ An integer — can be positive, negative or zero
- ▶ NB we don't use 2's complement to store it!
- ▶ Instead it is stored in binary as a positive integer with a **bias** added on
- ▶ (This is so that exponents can be efficiently compared (less/greater than) — 2's complement would be less efficient for this)

Exponent bias

Exponent bias

- ▶ E.g. if the bias is 127:

Exponent bias

- E.g. if the bias is 127:

| An exponent of... | ... is stored as |
|-------------------|------------------|
| -126 | 00000001 (1) |
| ⋮ | ⋮ |
| -1 | 01111110 (126) |
| 0 | 01111111 (127) |
| 1 | 10000000 (128) |
| ⋮ | ⋮ |
| 127 | 11111110 (254) |

Exponent bias

- ▶ E.g. if the bias is 127:

| An exponent of... | ... is stored as |
|-------------------|------------------|
| -126 | 00000001 (1) |
| ⋮ | ⋮ |
| -1 | 01111110 (126) |
| 0 | 01111111 (127) |
| 1 | 10000000 (128) |
| ⋮ | ⋮ |
| 127 | 11111110 (254) |

- ▶ (Exponents of 00000000 and 11111111 have special meaning — more on this later)

IEEE 754 floating point formats

| Type | Sign | Exponent | Mantissa | Total |
|--------------------|-------|-----------------------|----------|---------|
| Single precision | 1 bit | 8 bits bias 127 | 23 bits | 32 bits |
| Double precision | 1 bit | 11 bits bias 1023 | 52 bits | 64 bits |
| Extended precision | 1 bit | 15 bits bias 16383 | 64 bits | 80 bits |

IEEE 754 floating point formats

IEEE 754 floating point formats

- ▶ C# (and many other languages) have `float` and `double` types for single and double precision respectively

IEEE 754 floating point formats

- ▶ C# (and many other languages) have `float` and `double` types for single and double precision respectively
- ▶ Literals are interpreted as `float` if they end in `f`, otherwise `double`

IEEE 754 floating point formats

- ▶ C# (and many other languages) have **float** and **double** types for single and double precision respectively
- ▶ Literals are interpreted as **float** if they end in `f`, otherwise **double**
 - ▶ `3.14f` is a **float**

IEEE 754 floating point formats

- ▶ C# (and many other languages) have **float** and **double** types for single and double precision respectively
- ▶ Literals are interpreted as **float** if they end in `f`, otherwise **double**
 - ▶ `3.14f` is a **float**
 - ▶ `3.14` is a **double**

IEEE 754 floating point formats

- ▶ C# (and many other languages) have `float` and `double` types for single and double precision respectively
- ▶ Literals are interpreted as `float` if they end in `f`, otherwise `double`
 - ▶ `3.14f` is a `float`
 - ▶ `3.14` is a `double`
- ▶ Python's `float` type is double precision as standard

IEEE 754 floating point formats

- ▶ C# (and many other languages) have `float` and `double` types for single and double precision respectively
- ▶ Literals are interpreted as `float` if they end in `f`, otherwise `double`
 - ▶ `3.14f` is a `float`
 - ▶ `3.14` is a `double`
- ▶ Python's `float` type is double precision as standard
- ▶ Extended precision is not usually used in programs, but is used internally on Intel CPUs

Example

Example

What is the value stored in the following IEEE single-precision floating point number?

01000000110100000000000000000000

Example

Example

01000000110100000000000000000000

Example

010000001101000000000000000000000

- Sign bit is 0

Example

010000001101000000000000000000000

- ▶ Sign bit is 0
- ▶ Therefore the number is positive

Example

Example

0 10000001 101000000000000000000000

Example

0 10000001 101000000000000000000000

- ▶ Exponent is 10000001

Example

0 10000001 101000000000000000000000

- ▶ Exponent is 10000001
- ▶ This is 129 in binary

Example

0 10000001 101000000000000000000000

- ▶ Exponent is 10000001
- ▶ This is 129 in binary
- ▶ Exponent is stored with a bias of 127, therefore the actual exponent is $129 - 127 = 2$

Example

Example

01000000110100000000000000000000

Example

01000000110100000000000000000000

- Mantissa is 101000...

Example

01000000110100000000000000000000

- ▶ Mantissa is 101000...
- ▶ Remember we only store the digits after the “decimal point”, so the mantissa is actually 1.101000...

Example

01000000110100000000000000000000

- ▶ Mantissa is 101000...
- ▶ Remember we only store the digits after the “decimal point”, so the mantissa is actually 1.101000...
- ▶ The exponent is 2, so we move the point 2 places to the right: 110.1000...

Example

01000000110100000000000000000000

- ▶ Mantissa is 101000...
- ▶ Remember we only store the digits after the “decimal point”, so the mantissa is actually 1.101000...
- ▶ The exponent is 2, so we move the point 2 places to the right: 110.1000...
- ▶ $4 + 2 + \frac{1}{2} = 6.5$

Special floating point numbers

Special floating point numbers

- ▶ Zero is represented by a mantissa and exponent of all 0s

Special floating point numbers

- ▶ Zero is represented by a mantissa and exponent of all 0s
- ▶ An exponent of all 1s is used to represent infinity or NaN

Special floating point numbers

- ▶ Zero is represented by a mantissa and exponent of all 0s
- ▶ An exponent of all 1s is used to represent infinity or NaN
 - ▶ `float.PositiveInfinity` Or `double.PositiveInfinity`: a number which is greater than every other number

Special floating point numbers

- ▶ Zero is represented by a mantissa and exponent of all 0s
- ▶ An exponent of all 1s is used to represent infinity or NaN
 - ▶ `float.PositiveInfinity` Or `double.PositiveInfinity`: a number which is greater than every other number
 - ▶ `float.NegativeInfinity` Or `double.NegativeInfinity`: a number which is less than every other number

Special floating point numbers

- ▶ Zero is represented by a mantissa and exponent of all 0s
- ▶ An exponent of all 1s is used to represent infinity or NaN
 - ▶ `float.PositiveInfinity` Or `double.PositiveInfinity`: a number which is greater than every other number
 - ▶ `float.NegativeInfinity` Or `double.NegativeInfinity`: a number which is less than every other number
 - ▶ `float.NaN` Or `double.NaN`: “Not A Number” — `<`, `>`, `==` always return `false`

Special floating point numbers

- ▶ Zero is represented by a mantissa and exponent of all 0s
- ▶ An exponent of all 1s is used to represent infinity or NaN
 - ▶ `float.PositiveInfinity` Or `double.PositiveInfinity`: a number which is greater than every other number
 - ▶ `float.NegativeInfinity` Or `double.NegativeInfinity`: a number which is less than every other number
 - ▶ `float.NaN` Or `double.NaN`: "Not A Number" — `<`, `>`, `==` always return `false`
- ▶ Can check for these with `float.IsInfinity`, `double.IsNaN`, etc.

Special floating point numbers

- ▶ Zero is represented by a mantissa and exponent of all 0s
- ▶ An exponent of all 1s is used to represent infinity or NaN
 - ▶ `float.PositiveInfinity` Or `double.PositiveInfinity`: a number which is greater than every other number
 - ▶ `float.NegativeInfinity` Or `double.NegativeInfinity`: a number which is less than every other number
 - ▶ `float.NaN` Or `double.NaN`: “Not A Number” — `<`, `>`, `==` always return `false`
- ▶ Can check for these with `float.IsInfinity`, `double.IsNaN`, etc.
- ▶ Infinities and NaNs sometimes arise from calculations (e.g. dividing by zero)

Numerical precision



Precision of floating point numbers

Precision of floating point numbers

- ▶ Precision **varies** by **magnitude**

Precision of floating point numbers

- ▶ Precision **varies** by **magnitude**
- ▶ Numbers near 0 can be stored more accurately than numbers further from 0

Precision of floating point numbers

- ▶ Precision **varies** by **magnitude**
- ▶ Numbers near 0 can be stored more accurately than numbers further from 0
- ▶ Analogy: in scientific notation with 3 decimal places

Precision of floating point numbers

- ▶ Precision **varies** by **magnitude**
- ▶ Numbers near 0 can be stored more accurately than numbers further from 0
- ▶ Analogy: in scientific notation with 3 decimal places
 - ▶ Around 3.142×10^0 : can represent a difference of 0.001

Precision of floating point numbers

- ▶ Precision **varies** by **magnitude**
- ▶ Numbers near 0 can be stored more accurately than numbers further from 0
- ▶ Analogy: in scientific notation with 3 decimal places
 - ▶ Around 3.142×10^0 : can represent a difference of 0.001
 - ▶ Around 3.142×10^3 : can represent a difference of 1

Precision of floating point numbers

- ▶ Precision **varies** by **magnitude**
- ▶ Numbers near 0 can be stored more accurately than numbers further from 0
- ▶ Analogy: in scientific notation with 3 decimal places
 - ▶ Around 3.142×10^0 : can represent a difference of 0.001
 - ▶ Around 3.142×10^3 : can represent a difference of 1
 - ▶ Around 3.142×10^6 : can represent a difference of 1000

Range of floating point numbers

| Type | Smallest value (closest to 0) | Largest value (furthest from 0) |
|------------------|----------------------------------|------------------------------------|
| Single precision | $\pm 1.175 \times 10^{-38}$ | $\pm 3.403 \times 10^{38}$ |
| Double precision | $\pm 2.225 \times 10^{-308}$ | $\pm 1.798 \times 10^{308}$ |

Rounding errors

Rounding errors

- ▶ Many numbers cannot be represented exactly in IEEE float

Rounding errors

- ▶ Many numbers cannot be represented exactly in IEEE float
 - ▶ Similar to how decimal notation cannot exactly represent $\frac{1}{3} = 0.333333 \dots$ or $\frac{1}{7} = 0.142857 \dots$

Rounding errors

- ▶ Many numbers cannot be represented exactly in IEEE float
 - ▶ Similar to how decimal notation cannot exactly represent $\frac{1}{3} = 0.333333 \dots$ or $\frac{1}{7} = 0.142857 \dots$
- ▶ Decimal: can represent $\frac{a}{b}$ exactly iff $b = 2^m 5^n$

Rounding errors

- ▶ Many numbers cannot be represented exactly in IEEE float
 - ▶ Similar to how decimal notation cannot exactly represent $\frac{1}{3} = 0.333333 \dots$ or $\frac{1}{7} = 0.142857 \dots$
- ▶ Decimal: can represent $\frac{a}{b}$ exactly iff $b = 2^m 5^n$
- ▶ Binary: can represent $\frac{a}{b}$ exactly iff $b = 2^n$

Rounding errors

- ▶ Many numbers cannot be represented exactly in IEEE float
 - ▶ Similar to how decimal notation cannot exactly represent $\frac{1}{3} = 0.3333333 \dots$ or $\frac{1}{7} = 0.142857 \dots$
- ▶ Decimal: can represent $\frac{a}{b}$ exactly iff $b = 2^m 5^n$
- ▶ Binary: can represent $\frac{a}{b}$ exactly iff $b = 2^n$
- ▶ In particular, IEEE float can't represent $\frac{1}{10} = 0.1$ exactly!

Rounding errors

- ▶ Many numbers cannot be represented exactly in IEEE float
 - ▶ Similar to how decimal notation cannot exactly represent $\frac{1}{3} = 0.3333333 \dots$ or $\frac{1}{7} = 0.142857 \dots$
- ▶ Decimal: can represent $\frac{a}{b}$ exactly iff $b = 2^m 5^n$
- ▶ Binary: can represent $\frac{a}{b}$ exactly iff $b = 2^n$
- ▶ In particular, IEEE float can't represent $\frac{1}{10} = 0.1$ exactly!
- ▶ This can lead to **rounding errors** with some calculations

Testing for equality

Testing for equality

- ▶ Due to rounding errors, using `==` or `!=` with floating point numbers is almost always a bad idea

Testing for equality

- ▶ Due to rounding errors, using `==` or `!=` with floating point numbers is almost always a bad idea
- ▶ E.g. in most languages, `0.1 + 0.2 == 0.3` evaluates to `false`!

Testing for equality

- ▶ Due to rounding errors, using `==` or `!=` with floating point numbers is almost always a bad idea
- ▶ E.g. in most languages, `0.1 + 0.2 == 0.3` evaluates to **false!**
- ▶ Better to check for **approximate equality**: calculate the difference between the numbers, and check that it's smaller than some threshold

Testing for equality

- ▶ Due to rounding errors, using `==` or `!=` with floating point numbers is almost always a bad idea
- ▶ E.g. in most languages, `0.1 + 0.2 == 0.3` evaluates to **false!**
- ▶ Better to check for **approximate equality**: calculate the difference between the numbers, and check that it's smaller than some threshold
- ▶ E.g. Unity has `Mathf.Approximately` which does exactly this

Decimal type

Decimal type

- ▶ C# has a `decimal` type

Decimal type

- ▶ C# has a `decimal` type
- ▶ Uses base 10 rather than base 2, so avoids some of the surprises of IEEE float

Decimal type

- ▶ C# has a `decimal` type
- ▶ Uses base 10 rather than base 2, so avoids some of the surprises of IEEE float
- ▶ ... however not natively supported by the CPU, hence much slower than `float`/`double`