

COMP110: Principles of Computing

12: Further Computational Mathematics for Games

Worksheet E

- ► Assembly programming (TIS-100)
- ▶ Due week 13 (after Christmas see timetable)

Quiz E

- ► There is no Quiz E
- ► But complete quizzes A–D if you haven't already!

Final worksheet submission

- Check MyFalmouth for the deadline
- Download all five of your worksheet forks as zips
- Extract them into five separate folders
- ▶ Re-compress the five folders into a single zip file
- ▶ Upload this zip file to LearningSpace



Recursion and induction

$$\sum_{i=1}^n i = \frac{1}{2}n(n+1)$$

$$\sum_{i=1}^n i = \frac{1}{2}n(n+1)$$

$$\sum_{i=1}^{n} i = \frac{1}{2} n(n+1)$$

►
$$n = 1$$
: $1 = \frac{1}{2} \times 1 \times 2$



$$\sum_{i=1}^{n} i = \frac{1}{2} n(n+1)$$

►
$$n = 1$$
: $1 = \frac{1}{2} \times 1 \times 2$

►
$$n = 2$$
: $1 + 2 = \frac{1}{2} \times 2 \times 3 = 3$

$$\sum_{i=1}^{n} i = \frac{1}{2} n(n+1)$$

▶
$$n = 1$$
: $1 = \frac{1}{2} \times 1 \times 2$

►
$$n = 2$$
: $1 + 2 = \frac{1}{2} \times 2 \times 3 = 3$

►
$$n = 3$$
: $1 + 2 + 3 = \frac{1}{2} \times 3 \times 4 = 6$

$$\sum_{i=1}^n i = \frac{1}{2}n(n+1)$$

▶
$$n = 1$$
: $1 = \frac{1}{2} \times 1 \times 2$

►
$$n = 2$$
: $1 + 2 = \frac{1}{2} \times 2 \times 3 = 3$

►
$$n = 3$$
: $1 + 2 + 3 = \frac{1}{2} \times 3 \times 4 = 6$

٠.,



► We can verify the formula for individual values of *n*

- We can verify the formula for individual values of n
- ► How do we **prove** it for **all** *n*?

- We can verify the formula for individual values of n
- ► How do we prove it for all n?
- We can use proof by induction

Base case

Base case

►
$$n = 1: 1 = \frac{1}{2} \times 1 \times 2$$

Base case

►
$$n = 1$$
: $1 = \frac{1}{2} \times 1 \times 2$

Inductive assumption

Base case

►
$$n = 1$$
: $1 = \frac{1}{2} \times 1 \times 2$

Inductive assumption
$$\sum_{k=1}^{k-1} \frac{1}{k} \frac$$

Base case

►
$$n = 1: 1 = \frac{1}{2} \times 1 \times 2$$

Inductive assumption

Base case

►
$$n = 1$$
: $1 = \frac{1}{2} \times 1 \times 2$

Inductive assumption

$$\blacktriangleright \sum_{i=1}^{k} i = \left(\sum_{i=1}^{k-1} i\right) + k$$

Base case

►
$$n = 1$$
: $1 = \frac{1}{2} \times 1 \times 2$

Inductive assumption

$$\blacktriangleright \sum_{i=1}^{k} i = \left(\sum_{i=1}^{k-1} i\right) + k$$

$$ightharpoonup = \frac{1}{2}(k-1)k + k$$
 (by inductive assumption)

Base case

►
$$n = 1$$
: $1 = \frac{1}{2} \times 1 \times 2$

Inductive assumption

$$\blacktriangleright \sum_{i=1}^{k} i = \left(\sum_{i=1}^{k-1} i\right) + k$$

$$ightharpoonup = rac{1}{2}(k-1)k + k$$
 (by inductive assumption)

$$\blacktriangleright = \frac{1}{2}k^2 - \frac{1}{2}k + k$$

Base case

►
$$n = 1$$
: $1 = \frac{1}{2} \times 1 \times 2$

Inductive assumption

$$\blacktriangleright \sum_{i=1}^{k} i = \left(\sum_{i=1}^{k-1} i\right) + k$$

$$ightharpoonup = rac{1}{2}(k-1)k + k$$
 (by inductive assumption)

$$\blacktriangleright = \frac{1}{2}k^2 - \frac{1}{2}k + k$$

$$\blacktriangleright = \frac{1}{2}k^2 + \frac{1}{2}k$$

Base case

►
$$n = 1$$
: $1 = \frac{1}{2} \times 1 \times 2$

Inductive assumption

$$\blacktriangleright \sum_{i=1}^{k} i = \left(\sum_{i=1}^{k-1} i\right) + k$$

$$ightharpoonup = rac{1}{2}(k-1)k + k$$
 (by inductive assumption)

$$\blacktriangleright = \frac{1}{2}k^2 - \frac{1}{2}k + k$$

$$\blacktriangleright = \frac{1}{2}k^2 + \frac{1}{2}k$$

$$\blacktriangleright = \frac{1}{2}k(k-1)$$

Base case

►
$$n = 1$$
: $1 = \frac{1}{2} \times 1 \times 2$

Inductive assumption

$$\blacktriangleright \ \sum_{i=1}^{k-1} i = \frac{1}{2}(k-1)k$$

Therefore

$$\blacktriangleright \sum_{i=1}^{k} i = \left(\sum_{i=1}^{k-1} i\right) + k$$

$$ightharpoonup = rac{1}{2}(k-1)k + k$$
 (by inductive assumption)

$$\blacktriangleright = \frac{1}{2}k^2 - \frac{1}{2}k + k$$

$$\blacktriangleright = \frac{1}{2}k^2 + \frac{1}{2}k$$

$$\blacktriangleright = \frac{1}{2}k(k-1)$$

So if the formula works for n = k - 1, then it works for n = k

► We know:

- ▶ We know:
 - ▶ The formula works for n = 1

- ▶ We know:
 - ▶ The formula works for n = 1
 - ▶ If the formula works for n = k 1, then it works for n = k

- ▶ We know:
 - ▶ The formula works for n = 1
 - ▶ If the formula works for n = k 1, then it works for n = k
- ▶ The formula works for n = 1

- ▶ We know:
 - ▶ The formula works for n = 1
 - ▶ If the formula works for n = k 1, then it works for n = k
- ▶ The formula works for n = 1
- ▶ Therefore the formula works for n = 1 + 1 = 2

- ▶ We know:
 - ▶ The formula works for n = 1
 - ▶ If the formula works for n = k 1, then it works for n = k
- ▶ The formula works for n = 1
- ▶ Therefore the formula works for n = 1 + 1 = 2
- ▶ Therefore the formula works for n = 2 + 1 = 3

- ▶ We know:
 - ▶ The formula works for n = 1
 - ▶ If the formula works for n = k 1, then it works for n = k
- ▶ The formula works for n = 1
- ▶ Therefore the formula works for n = 1 + 1 = 2
- ▶ Therefore the formula works for n = 2 + 1 = 3
- ▶ Therefore the formula works for n = 3 + 1 = 4

- ▶ We know:
 - ▶ The formula works for n = 1
 - ▶ If the formula works for n = k 1, then it works for n = k
- ▶ The formula works for n = 1
- ▶ Therefore the formula works for n = 1 + 1 = 2
- ▶ Therefore the formula works for n = 2 + 1 = 3
- ▶ Therefore the formula works for n = 3 + 1 = 4
- ٠...

Completing the proof

- ▶ We know:
 - ▶ The formula works for n = 1
 - ▶ If the formula works for n = k 1, then it works for n = k
- ▶ The formula works for n = 1
- ▶ Therefore the formula works for n = 1 + 1 = 2
- ▶ Therefore the formula works for n = 2 + 1 = 3
- ▶ Therefore the formula works for n = 3 + 1 = 4
- ٠...
- Therefore the formula works for all positive integers n

Exercise

Prove

$$\sum_{i=1}^{n} i^2 = \frac{n(n+1)(2n+1)}{6}$$

▶ I want to prove something for all n

- ▶ I want to prove something for all n
- ► Given k, if I had already proved n = k 1 then I could prove n = k

- ▶ I want to prove something for all n
- Fiven k, if I had already proved n = k 1 then I could prove n = k
- ▶ I can also prove n = 1

- ► I want to prove something for all n
- Given k, if I had already proved n = k − 1 then I could prove n = k
- ▶ I can also prove n = 1
- Therefore by induction I can prove the result for all n

▶ I want to solve a problem

- ▶ I want to solve a problem
- If I already had a function to solve smaller instances of the problem, I could use it to write my function



- ▶ I want to solve a problem
- If I already had a function to solve smaller instances of the problem, I could use it to write my function
- ▶ I can solve the smallest possible problem

- I want to solve a problem
- If I already had a function to solve smaller instances of the problem, I could use it to write my function
- I can solve the smallest possible problem
- Therefore I can write a recursive function

Exercise

- Write a pseudocode function to calculate the total size of all files in a directory and its subdirectories
- You may use the following functions in your pseudocode:
 - ListDir(directory): return a list of names of all files and folders in the given directory
 - GETSIZE(filename): return the size, in bytes, of the given file
 - IsDIR(name), IsFILE(name): determine whether the given name refers to a file or a directory

procedure CALCDIRSIZE(directory)
...

▷ return total size in bytes
end procedure