

# DPLYR Evaluate Function over N Individuals with Individual Specific and Shared Arrays of Parameters

Fan Wang

2020-04-01

## Contents

Row Input Functions . . . . .	1
-------------------------------	---

### Row Input Functions

Go back to [fan's REconTools Package](#), [R4Econ Repository](#), or [Intro Stats with R Repository](#).

We want evaluate nonlinear function  $f(Q\_i, y\_i, ar\_x, ar\_y, c, d)$ , where  $c$  and  $d$  are constants, and  $ar\_x$  and  $ar\_y$  are arrays, both fixed.  $x\_i$  and  $y\_i$  vary over each row of matrix. We would like to evaluate this nonlinear function concurrently across  $N$  individuals. The eventual goal is to find the  $i$  specific  $Q$  that solves the nonlinear equations.

This is a continuation of [R use Apply, Sapply and dplyr Mutate to Evaluate one Function Across Rows of a Matrix](#)

**Set up Input Arrays** There is a function that takes  $M = Q + P$  inputs, we want to evaluate this function  $N$  times. Each time, there are  $M$  inputs, where all but  $Q$  of the  $M$  inputs, meaning  $P$  of the  $M$  inputs, are the same. In particular,  $P = Q * N$ .

$$M = Q + P = Q + Q * N$$

```
# it_child_count = N, the number of children
it_N_child_cnt = 5
# it_heter_param = Q, number of parameters that are heterogeneous across children
it_Q_hetpa_cnt = 2

# P fixed parameters, nN is N dimensional, nP is P dimensional
ar_nN_A = seq(-2, 2, length.out = it_N_child_cnt)
ar_nN_alpha = seq(0.1, 0.9, length.out = it_N_child_cnt)
ar_nP_A_alpha = c(ar_nN_A, ar_nN_alpha)
ar_nN_N_choice = seq(1, it_N_child_cnt) / sum(seq(1, it_N_child_cnt))

# N by Q varying parameters
mt_nN_by_nQ_A_alpha = cbind(ar_nN_A, ar_nN_alpha, ar_nN_N_choice)
# Show
kable(mt_nN_by_nQ_A_alpha) %>%
  kable_styling_fc_wide()
```

**Testing Function** Test non-linear Equation.

ar_nN_A	ar_nN_alpha	ar_nN_N_choice
-2	0.1	0.0666667
-1	0.3	0.1333333
0	0.5	0.2000000
1	0.7	0.2666667
2	0.9	0.3333333

```

# Test Parameters
fl_N_agg = 100
fl_rho = -1
fl_N_q = ar_nN_N_choice[4]*fl_N_agg
ar_A_alpha = mt_nN_by_nQ_A_alpha[4,]
# Apply Function
ar_p1_s1 = exp((ar_A_alpha[1] - ar_nN_A)*fl_rho)
ar_p1_s2 = (ar_A_alpha[2]/ar_nN_alpha)
ar_p1_s3 = (1/(ar_nN_alpha*fl_rho - 1))
ar_p1 = (ar_p1_s1*ar_p1_s2)^ar_p1_s3
ar_p2 = fl_N_q^((ar_A_alpha[2]*fl_rho-1)/(ar_nN_alpha*fl_rho-1))
ar_overall = ar_p1*ar_p2
fl_overall = fl_N_agg - sum(ar_overall)
print(fl_overall)

```

```
## [1] -598.2559
```

Implement the non-linear problem's evaluation using apply over all  $N$  individuals.

```

# Define Implicit Function
ffi_nonlin_dplyrdo <- function(fl_A, fl_alpha, fl_N, ar_A, ar_alpha, fl_N_agg, fl_rho){
  # ar_A_alpha[1] is A
  # ar_A_alpha[2] is alpha

  # # Test Parameters
  # fl_N = 100
  # fl_rho = -1
  # fl_N_q = 10

  # Apply Function
  ar_p1_s1 = exp((fl_A - ar_A)*fl_rho)
  ar_p1_s2 = (fl_alpha/ar_alpha)
  ar_p1_s3 = (1/(ar_alpha*fl_rho - 1))
  ar_p1 = (ar_p1_s1*ar_p1_s2)^ar_p1_s3
  ar_p2 = fl_N^((fl_alpha*fl_rho-1)/(ar_alpha*fl_rho-1))
  ar_overall = ar_p1*ar_p2
  fl_overall = fl_N_agg - sum(ar_overall)

  return(fl_overall)
}

```

```

# Parameters
fl_rho = -1

# Evaluate Function
print(ffl_nonlin_dplyrdo(mt_nN_by_nQ_A_alpha[1,1],
                        mt_nN_by_nQ_A_alpha[1,2],
                        mt_nN_by_nQ_A_alpha[1,3]*fl_N_agg,
                        ar_nN_A, ar_nN_alpha, fl_N_agg, fl_rho))

## [1] 81.86645

for (i in seq(1,dim(mt_nN_by_nQ_A_alpha)[1])){
  fl_eval = ffl_nonlin_dplyrdo(mt_nN_by_nQ_A_alpha[i,1],
                              mt_nN_by_nQ_A_alpha[i,2],
                              mt_nN_by_nQ_A_alpha[i,3]*fl_N_agg,
                              ar_nN_A, ar_nN_alpha, fl_N_agg, fl_rho)

  print(fl_eval)
}

## [1] 81.86645
## [1] 54.48885
## [1] -65.5619
## [1] -598.2559
## [1] -3154.072

# Convert Matrix to Tibble
ar_st_col_names = c('fl_A', 'fl_alpha', 'fl_N')
tb_nN_by_nQ_A_alpha <- as_tibble(mt_nN_by_nQ_A_alpha) %>% rename_all(~c(ar_st_col_names))

# Define Implicit Function
ffl_nonlin_dplyrdo <- function(fl_A, fl_alpha, fl_N, ar_A, ar_alpha, fl_N_agg, fl_rho){

  # Test Parameters
  # ar_A = ar_nN_A
  # ar_alpha = ar_nN_alpha
  # fl_N = 100
  # fl_rho = -1
  # fl_N_q = 10

  # Apply Function
  ar_p1_s1 = exp((fl_A - ar_A)*fl_rho)
  ar_p1_s2 = (fl_alpha/ar_alpha)
  ar_p1_s3 = (1/(ar_alpha*fl_rho - 1))
  ar_p1 = (ar_p1_s1*ar_p1_s2)^ar_p1_s3
  ar_p2 = (fl_N*fl_N_agg)^((fl_alpha*fl_rho-1)/(ar_alpha*fl_rho-1))
  ar_overall = ar_p1*ar_p2
  fl_overall = fl_N_agg - sum(ar_overall)

  return(fl_overall)
}

# fl_A, fl_alpha are from columns of tb_nN_by_nQ_A_alpha
tb_nN_by_nQ_A_alpha = tb_nN_by_nQ_A_alpha %>% rowwise() %>%

```

```

mutate(dplyr_eval = ffi_nonlin_dplyrdo(fl_A, fl_alpha, fl_N,
                                       ar_nN_A, ar_nN_alpha,
                                       fl_N_agg, fl_rho))
# Show
kable(tb_nN_by_nQ_A_alpha) %>%
  kable_styling_fc_wide()

```

fl__A	fl__alpha	fl__N	dplyr__eval
-2	0.1	0.06666667	81.86645
-1	0.3	0.13333333	54.48885
0	0.5	0.20000000	-65.56190
1	0.7	0.26666667	-598.25595
2	0.9	0.33333333	-3154.07226

Evaluate Nonlinear Function using dplyr mutate