

R Summarize a Quantitative/Continuous Variable with Categorical Groups

Fan Wang

2020-04-01

Contents

Histogram 1

Histogram

Generate Test Score Dataset

Go to the [RMD](#), [R](#), [PDF](#), or [HTML](#) version of this file. Go back to [fan's REconTools Package](#), [R4Econ Repository \(bookdown site\)](#), or [Intro Stats with R Repository](#).

- r generate text string as csv
- r tibble matrix hand input

First, we will generate a test score dataset, directly from string. Below we type line by line a dataset with four variables in comma separated (csv) format, where the first row includes the variables names. These texts could be stored in a separate file, or they could be directly included in code and read in as csv

```
ar_test_scores_ec3 <- c(107.72,101.28,105.92,109.31,104.27,110.27,91.92846154,81.8,109.0071429,103.07,99.07,100.27,104.27,90.23615385,77.8,103.4357143,97.07,93.07)
ar_test_scores_ec1 <- c(101.72,101.28,99.92,103.31,100.27,104.27,90.23615385,77.8,103.4357143,97.07,93.07)
mt_test_scores <- cbind(ar_test_scores_ec1, ar_test_scores_ec3)
ar_st_varnames <- c('course_total_ec1p', 'course_total_ec3p')
tb_final_twovar <- as_tibble(mt_test_scores) %>% rename_all(~c(ar_st_varnames))
summary(tb_final_twovar)
```

A Dataset with only Two Continuous Variable

```
## course_total_ec1p course_total_ec3p
## Min. : 40.48 Min. : 44.23
## 1st Qu.: 76.46 1st Qu.: 79.91
## Median : 86.35 Median : 89.28
## Mean : 83.88 Mean : 87.90
## 3rd Qu.: 95.89 3rd Qu.:100.75
## Max. :104.27 Max. :112.22
```

```
ff_summ_percentiles(df = tb_final_twovar, bl_statsasrows = TRUE, col2varname = FALSE)
```

```
## # A tibble: 17 x 3
```

	stats	course.total.ec1p	course.total.ec3p
	<chr>	<chr>	<chr>
## 1 n	46	46	
## 2 NAobs	0	0	
## 3 ZEROobs	0	0	

```
## 4 mean      83.87572      87.90239
## 5 sd        15.87272      16.76041
## 6 cv         0.1892409     0.1906706
## 7 min        40.475       44.225
## 8 p01        42.14434     45.82202
## 9 p05        56.9650      57.1575
## 10 p10       63.05462     66.07500
## 11 p25       76.45616     79.90500
## 12 p50       86.35236     89.27923
## 13 p75       " 95.89054"   100.75250
## 14 p90       100.8137     106.8200
## 15 p95       102.9125     109.2343
## 16 p99       103.8946     111.3439
## 17 max       104.2700     112.2225
```

```
ar_final_scores <- c(94.28442509,95.68817475,97.25219512,77.89268293,95.08795497,93.27380863,92.3,84.25)
mt_test_scores <- cbind(seq(1,length(ar_final_scores)), ar_final_scores)
ar_st_varnames <- c('index', 'course_final')
tb_onevar <- as_tibble(mt_test_scores) %>% rename_all(~c(ar_st_varnames))
summary(tb_onevar)
```

A Dataset with one Continuous Variable and Histogram

```
##      index      course_final
## Min.   : 1.0   Min.     : 2.293
## 1st Qu.:12.5   1st Qu.: 76.372
## Median :24.0   Median : 86.959
## Mean   :24.0   Mean    : 82.415
## 3rd Qu.:35.5   3rd Qu.: 94.686
## Max.   :47.0   Max.     :100.898
```

```
ff_summ_percentiles(df = tb_onevar, bl_statsasrows = TRUE, col2varname = FALSE)
```

```
## # A tibble: 17 x 3
##   stats      course.final index
##   <chr>      <chr>      <chr>
## 1 n          47          47
## 2 NAobs      0           0
## 3 ZEROobs    0           0
## 4 mean       82.41501    24.00000
## 5 sd         18.35476    13.71131
## 6 cv          0.2227113    0.5713046
## 7 min         2.292683    1.000000
## 8 p01         18.67401    " 1.46000"
## 9 p05         49.72075    " 3.30000"
## 10 p10        66.28051    " 5.60000"
## 11 p25        76.37177    12.50000
## 12 p50        86.95932    24.00000
## 13 p75        94.68619    35.50000
## 14 p90        97.52332    42.40000
## 15 p95        99.47459    44.70000
## 16 p99       100.5244    " 46.5400"
## 17 max       100.898      " 47.000"
```

```

#load in data empirically by hand
txt_test_data <- "init_prof, later_prof, class_id, exam_score
'SW', 'SW', 1, 102
'SW', 'SW', 1, 102
'SW', 'SW', 1, 101
'SW', 'SW', 1, 100
'SW', 'SW', 1, 100
'SW', 'SW', 1, 99
'SW', 'SW', 1, 98.5
'SW', 'SW', 1, 98.5
'SW', 'SW', 1, 97
'SW', 'SW', 1, 95
'SW', 'SW', 1, 94
'SW', 'SW', 1, 91
'SW', 'SW', 1, 91
'SW', 'SW', 1, 90
'SW', 'SW', 1, 89
'SW', 'SW', 1, 88.5
'SW', 'SW', 1, 88
'SW', 'SW', 1, 87
'SW', 'SW', 1, 87
'SW', 'SW', 1, 87
'SW', 'SW', 1, 86
'SW', 'SW', 1, 86
'SW', 'SW', 1, 84
'SW', 'SW', 1, 82
'SW', 'SW', 1, 78.5
'SW', 'SW', 1, 76
'SW', 'SW', 1, 72
'SW', 'SW', 1, 70.5
'SW', 'SW', 1, 67.5
'SW', 'SW', 1, 67.5
'SW', 'SW', 1, 67
'SW', 'SW', 1, 63.5
'SW', 'SW', 1, 60
'SW', 'SW', 1, 59
'SW', 'SW', 1, 44.5
'SW', 'SW', 1, 44
'SW', 'SW', 1, 42.5
'SW', 'SW', 1, 40.5
'SW', 'SW', 1, 40.5
'SW', 'SW', 1, 36.5
'SW', 'SW', 1, 35.5
'SW', 'SW', 1, 21.5
'SW', 'SW', 1, 4
'MP', 'MP', 2, 105
'MP', 'MP', 2, 103
'MP', 'MP', 2, 102
'MP', 'MP', 2, 101
'MP', 'MP', 2, 101
'MP', 'MP', 2, 100.5
'MP', 'MP', 2, 100

```

```

'MP', 'MP', 2, 99
'MP', 'MP', 2, 97
'MP', 'MP', 2, 97
'MP', 'MP', 2, 97
'MP', 'MP', 2, 97
'MP', 'MP', 2, 96
'MP', 'MP', 2, 95
'MP', 'MP', 2, 91
'MP', 'MP', 2, 89
'MP', 'MP', 2, 85
'MP', 'MP', 2, 84
'MP', 'MP', 2, 84
'MP', 'MP', 2, 84
'MP', 'MP', 2, 83.5
'MP', 'MP', 2, 82.5
'MP', 'MP', 2, 81.5
'MP', 'MP', 2, 80.5
'MP', 'MP', 2, 80
'MP', 'MP', 2, 77
'MP', 'MP', 2, 77
'MP', 'MP', 2, 75
'MP', 'MP', 2, 75
'MP', 'MP', 2, 71
'MP', 'MP', 2, 70
'MP', 'MP', 2, 68
'MP', 'MP', 2, 63
'MP', 'MP', 2, 56
'MP', 'MP', 2, 56
'MP', 'MP', 2, 55.5
'MP', 'MP', 2, 49.5
'MP', 'MP', 2, 48.5
'MP', 'MP', 2, 47.5
'MP', 'MP', 2, 44.5
'MP', 'MP', 2, 34.5
'MP', 'MP', 2, 29.5
'CA', 'MP', 3, 103
'CA', 'MP', 3, 103
'CA', 'MP', 3, 101
'CA', 'MP', 3, 96.5
'CA', 'MP', 3, 93.5
'CA', 'MP', 3, 93
'CA', 'MP', 3, 93
'CA', 'MP', 3, 92
'CA', 'MP', 3, 90
'CA', 'MP', 3, 90
'CA', 'MP', 3, 89
'CA', 'MP', 3, 86.5
'CA', 'MP', 3, 84.5
'CA', 'MP', 3, 83
'CA', 'MP', 3, 83
'CA', 'MP', 3, 82
'CA', 'MP', 3, 78
'CA', 'MP', 3, 75

```

'CA', 'MP', 3, 74.5
 'CA', 'MP', 3, 70
 'CA', 'MP', 3, 54.5
 'CA', 'MP', 3, 52
 'CA', 'MP', 3, 50
 'CA', 'MP', 3, 42
 'CA', 'MP', 3, 36.5
 'CA', 'MP', 3, 28
 'CA', 'MP', 3, 26
 'CA', 'MP', 3, 11
 'CA', 'SN', 4, 103
 'CA', 'SN', 4, 103
 'CA', 'SN', 4, 102
 'CA', 'SN', 4, 102
 'CA', 'SN', 4, 101
 'CA', 'SN', 4, 100
 'CA', 'SN', 4, 98
 'CA', 'SN', 4, 98
 'CA', 'SN', 4, 98
 'CA', 'SN', 4, 95
 'CA', 'SN', 4, 95
 'CA', 'SN', 4, 92.5
 'CA', 'SN', 4, 92
 'CA', 'SN', 4, 91
 'CA', 'SN', 4, 90
 'CA', 'SN', 4, 85.5
 'CA', 'SN', 4, 84
 'CA', 'SN', 4, 82.5
 'CA', 'SN', 4, 81
 'CA', 'SN', 4, 77.5
 'CA', 'SN', 4, 77
 'CA', 'SN', 4, 72
 'CA', 'SN', 4, 71.5
 'CA', 'SN', 4, 69
 'CA', 'SN', 4, 68.5
 'CA', 'SN', 4, 68
 'CA', 'SN', 4, 67
 'CA', 'SN', 4, 65.5
 'CA', 'SN', 4, 62.5
 'CA', 'SN', 4, 62
 'CA', 'SN', 4, 61.5
 'CA', 'SN', 4, 61
 'CA', 'SN', 4, 57.5
 'CA', 'SN', 4, 54
 'CA', 'SN', 4, 52.5
 'CA', 'SN', 4, 51
 'CA', 'SN', 4, 50.5
 'CA', 'SN', 4, 50
 'CA', 'SN', 4, 49
 'CA', 'SN', 4, 43
 'CA', 'SN', 4, 39.5
 'CA', 'SN', 4, 32.5
 'CA', 'SN', 4, 25.5

```
'CA', 'SN', 4, 18"
```


```
csv_test_data = read.csv(text=txt_test_data, header=TRUE)
ar_st_varnames <- c('first_half_professor', 'second_half_professor', 'course_id', 'exam_score')
tb_test_data <- as_tibble(csv_test_data) %>% rename_all(~c(ar_st_varnames))
summary(tb_test_data)
```

A Dataset with Multiple Variables

```
## first_half_professor second_half_professor course_id exam_score
## 'CA':72 'MP':70 Min. :1.000 Min. : 4.00
## 'MP':42 'SN':44 1st Qu.:1.000 1st Qu.: 60.00
## 'SW':43 'SW':43 Median :2.000 Median : 82.00
## Mean :2.465 Mean : 75.08
## 3rd Qu.:4.000 3rd Qu.: 94.00
## Max. :4.000 Max. :105.00
```


Test Score Distributions

```
ggplot(tb_final_twovar, aes(x=ar_test_scores_ec3)) +
  geom_histogram(bins=25) +
  labs(title = paste0('Sandbox: Final Distribution (Econ 2370, FW)'),
       caption = 'FW Section, formula: 0.3*exam1Perc + 0.3*exam2Perc + 0.42*HWtotalPerc + 0.03*AttendancePerc + 0.03*ExtraCreditPerc',
       theme_bw())
```



Histogram

```
ggplot(tb_test_data, aes(x=exam_score)) +  
  geom_histogram(bins=16) +  
  labs(title = paste0('Exam Distribution'),  
        caption = 'All Sections') +  
  theme_bw()
```



All Sections