

Randomly Draw Subsets of Rows from Matrix in R

Fan Wang

2020-04-19

Contents

1 Drawly Random Rows	1
1.1 Draw Random Subset of Sample	1
1.2 Random Subset of Panel	2

1 Drawly Random Rows

Go to the [RMD](#), [R](#), [PDF](#), or [HTML](#) version of this file. Go back to [fan's REconTools](#) Package, [R Code Examples](#) Repository ([bookdown site](#)), or [Intro Stats with R](#) Repository ([bookdown site](#)).

1.1 Draw Random Subset of Sample

- r random discrete

We have a sample of N individuals in some dataframe. Draw without replacement a subset $M < N$ of rows.

```
# parameters, it_M < it_N
it_N <- 10
it_M <- 5

# Draw it_m from indexed list of it_N
set.seed(123)
ar_it_rand_idx <- sample(it_N, it_M, replace=FALSE)

# dataframe
df_full <- as_tibble(matrix(rnorm(4,mean=0,sd=1), nrow=it_N, ncol=4)) %>% rowid_to_column(var = "ID")

# random Subset
df_rand_sub_a <- df_full[ar_it_rand_idx,]

# Random subset also
df_rand_sub_b <- df_full[sample(dim(df_full)[1], it_M, replace=FALSE),]

# Print
# Display
kable(df_full) %>% kable_styling_fc()

kable(df_rand_sub_a) %>% kable_styling_fc()

kable(df_rand_sub_b) %>% kable_styling_fc()
```

ID	V1	V2	V3	V4
1	0.1292877	0.4609162	0.1292877	0.4609162
2	1.7150650	-1.2650612	1.7150650	-1.2650612
3	0.4609162	0.1292877	0.4609162	0.1292877
4	-1.2650612	1.7150650	-1.2650612	1.7150650
5	0.1292877	0.4609162	0.1292877	0.4609162
6	1.7150650	-1.2650612	1.7150650	-1.2650612
7	0.4609162	0.1292877	0.4609162	0.1292877
8	-1.2650612	1.7150650	-1.2650612	1.7150650
9	0.1292877	0.4609162	0.1292877	0.4609162
10	1.7150650	-1.2650612	1.7150650	-1.2650612

ID	V1	V2	V3	V4
3	0.4609162	0.1292877	0.4609162	0.1292877
10	1.7150650	-1.2650612	1.7150650	-1.2650612
2	1.7150650	-1.2650612	1.7150650	-1.2650612
8	-1.2650612	1.7150650	-1.2650612	1.7150650
6	1.7150650	-1.2650612	1.7150650	-1.2650612

ID	V1	V2	V3	V4
5	0.1292877	0.4609162	0.1292877	0.4609162
3	0.4609162	0.1292877	0.4609162	0.1292877
9	0.1292877	0.4609162	0.1292877	0.4609162
1	0.1292877	0.4609162	0.1292877	0.4609162
4	-1.2650612	1.7150650	-1.2650612	1.7150650

1.2 Random Subset of Panel

There are N individuals, each could be observed M times, but then select a subset of rows only, so each person is randomly observed only a subset of times. Specifically, there are 3 unique students with student ids, and the second variable shows the random dates in which the student showed up in class, out of the 10 classes available.

```
# Define
it_N <- 3
it_M <- 10
svr_id <- 'student_id'

# dataframe
set.seed(123)
df_panel_rand <- as_tibble(matrix(it_M, nrow=it_N, ncol=1)) %>%
  rowid_to_column(var = svr_id) %>%
  uncount(V1) %>%
  group_by(!sym(svr_id)) %>% mutate(date = row_number()) %>%
  ungroup() %>% mutate(in_class = case_when(rnorm(n(), mean=0, sd=1) < 0 ~ 1, TRUE ~ 0)) %>%
  filter(in_class == 1) %>% select(!sym(svr_id), date) %>%
  rename(date_in_class = date)

# Print
kable(df_panel_rand) %>% kable_styling_fc()
```

student_id	date_in_class
1	1
1	2
1	8
1	9
1	10
2	5
2	8
2	10
3	1
3	2
3	3
3	4
3	5
3	6
3	9